

# UTILITY PATENT APPLICATION TRANSMITTAL

Only for new nonprovisional applications under 37 CFR 1.53(b)

Attorney Docket No. BB-1036-B

First Named Inventor or Application Identifier

John Browne

Express Mail Label No. EE353045470US

Express Mailing Date December 23, 1998

## APPLICATION ELEMENTS

See MPEP chapter 600 concerning utility patent application contents.

ADDRESS TO:

Assistant Commissioner for Patents  
Box Patent Application  
Washington, DC 20231

1. ☒ Fee (Authority to charge deposit account below.)  
(Submit an original, and a duplicate for fee processing)
2. ☒ Specification [Total Pages 164]  
(preferred arrangement set forth below)
- Descriptive title of the invention
  - Cross References to Related Applications (if needed)
  - Statement Regarding Fed sponsored R & D (if needed)
  - Reference to Microfiche Appendix (if filed)
  - Background of the Invention
  - Brief Summary of the Invention
  - Brief Description of the Drawings (if filed)
  - Detailed Description
  - Claim(s)
  - Abstract of the Disclosure

6. ☐ Microfiche Computer Program (Appendix)
7. ☐ Nucleotide and/or Amino Acid Sequence Submission  
(if applicable, all necessary)
- a. ☐ Computer Readable Copy
  - b. ☐ Paper Copy (identical to computer copy)
  - c. ☐ Statement verifying identity of above copies

## ACCOMPANYING APPLICATION PARTS

3. ☐ Drawing(s) (35 USC 113) [Total Sheets 0]
4. ☒ Oath or Declaration [Total Pages 2]
- a. ☐ Newly executed (original or copy)
  - b. ☒ Copy from a prior application (37 CFR 1.63(d))  
(for continuation/divisional with Box 14 completed)
  - i. ☐ **DELETION OF INVENTORS**  
Signed Statement below at 15 deleting inventor(s) named in the prior application, see 37 CFR 1.63(d)(2) and 1.33(b).
5. ☒ Incorporation by Reference (useable if Box 4b is checked)  
The entire disclosure of the prior application, from which a copy of the oath or declaration is supplied under Box 4b, is considered as being part of the disclosure of the accompanying application and is hereby incorporated by reference therein.

8. ☐ Power of Attorney
9. ☒ Information Disclosure Statement (IDS)/Cover Letter plus PTO-1449 ☐ Copies of IDS Citations
10. ☒ Preliminary Amendment
11. ☒ Return Receipt Postcard (MPEP 503)  
(Should be specifically itemized)
12. ☐ Certified Copy of Priority Document(s)  
(if foreign priority is claimed)
13. ☒ Other:  
Request for Transfer of Computer Readable Form From Previously Filed Application

14. If a CONTINUING APPLICATION, check appropriate box and supply the requisite information:

☒ Continuation ☐ Divisional ☐ Continuation-in-part (CIP) of prior Application No.: 08/244,205

15. ☐ **DELETION OF INVENTOR(S) STATEMENT:** This application is being filed by less than all the inventors named in the prior application. In accordance with 37 CFR 1.63(d)(2) and 1.33(b), the Assistant Commissioner is requested to delete the name(s) of the following person or persons who are not inventors of the invention being claimed in this application:

16. ☒ Amend the specification by inserting before the first line the sentence:  
-- This is a continuation of Application No. 08/244,205 filed August 26, 1994, now allowed, which is a 371 of PCT/US92/10284, filed December 3, 1992, now abandoned, which is a Continuation-in-Part of Application No. 07/804,259 filed December 4, 1991, now abandoned. --

17. ☐ Cancel in this application original claims \_\_\_\_ of the prior application before calculating the filing. (At least one original independent claim must be retained for filing purposes.)

18. ☐ Priority of foreign Application No. \_\_\_\_\_ filed on \_\_\_\_\_ in \_\_\_\_\_

\_\_\_\_\_ is claimed under 35 U.S.C. 119.  
(country)

CLAIMS	(1) FOR	(2) NUMBER FILED	(3) NUMBER EXTRA	(4) RATE	(5) CALCULATIONS
	TOTAL CLAIMS (37 CFR 1.16(c))	21 - 20 =	1	x \$ 18 =	\$ 18.00
	INDEPENDENT CLAIMS (37 CFR 1.16(b))	7 - 3 =	4	x \$ 78 =	\$ 312.00
	MULTIPLE DEPENDENT CLAIM(S) (if applicable)			+ \$ 260 =	260.00
				BASIC FEE (37 CFR 1.16(a))	+ \$ 760.00
				TOTAL =	\$ 1350.00

19. The Commissioner is hereby authorized to credit overpayments or charge the following fees to Deposit Account No. 04-1928:

a. ☒ Fees required under 37 CFR 1.16.

b. ☒ Fees required under 37 CFR 1.17.

20. ☐ Other:

21. CORRESPONDENCE ADDRESS					
NAME	Lynne M. Christenbury				
ADDRESS	E. I. du Pont de Nemours and Company				
	Legal - Patents				
	1007 Market Street				
CITY	Wilmington	STATE	Delaware	ZIP CODE	19898
COUNTRY	U.S.A.	TELEPHONE	302-992-5481	FAX	302-892-7949

22. SIGNATURE OF ATTORNEY OR AGENT REQUIRED		
NAME	Lynne M. Christenbury	REG. NO.: 30,971
SIGNATURE	<i>Lynne M. Christenbury</i>	
DATE	December 23, 1998	

PATENT  
IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

IN THE APPLICATION OF:

JOHN BROWSE  
LUIS PEREZ GRAU  
ANTHONY J. KINNEY  
JOHN W. PIERCE, JR.  
ANNA M. WIERZBICKI  
NARENDRA S. YADAV

CASE NO.: BB-1036-B

APPLN. NO.: UNKNOWN

GROUP ART UNIT: UNKNOWN

FILED: CONCURRENTLY HERewith

EXAMINER: UNKNOWN

FOR: FATTY ACID DESATURASE GENES  
FROM PLANTS

Date: DECEMBER 23, 1998

Assistant Commissioner for Patents  
Washington, DC 20231

Sir:

**Preliminary Amendment**

This is submitted to facilitate prosecution of the above-identified application.

**In the Claims**

Kindly cancel claims 2-19.

Kindly add the following new claims:

--20. An isolated nucleic acid fragment which is useful in antisense inhibition or sense suppression of endogenous desaturase activity in a transformed plant wherein said isolated nucleic acid fragment hybridizes to one of the nucleotide sequences set forth in SEQ ID NOS:1, 3, 5, 7, 9, 11, and 15 or any portion thereof under one of the following sets of conditions:

(a) hybridization in 50 mM Tris, pH 7.6, 6X SSC, 5X Denhardt's, 0.5% sodium dodecyl sulfate (SDS), 100 µg denatured calf thymus DNA at 50°C overnight and wash with 6X SSC, 0.5% SDS at room temperature for 15 min, repeat with 2X

**CERTIFICATE OF MAILING BY "EXPRESS MAIL"**

"EXPRESS MAIL" MAILING LABEL NUMBER EE353045470US

I hereby certify that this paper or fee is being deposited with the United States Postal Service as "Express Mail Post Office to Addressee" Service under 37 C.F.R. 1.10 in an envelope addressed to: Assistant Commissioner for Patents, Washington, DC 20231 on this date.

Diane M. Dick  
Diane M. Dick

12/23/98  
Date

SSC, 0.5% SDS at 45°C for 30 min, then repeat twice with 0.2X SSC, 0.5% SDS at 50 °C for 30 min each;

(b) hybridization in 50 mM Tris, pH 7.6, 6X SSC, 5X Denhardt's, 0.5% sodium dodecyl sulfate (SDS), 100 µg denatured calf thymus DNA at 50°C overnight and wash with 6X SSC, 0.5% SDS at room temperature for 15 min, repeat with 2X SSC, 0.5% SDS at 45°C for 30 min, then repeat twice with 0.2X SSC, 0.5% SDS at 60 °C for 30 min each;

(c) hybridization in 50 mM Tris-HCl, pH 7.5, 1M NaCl, 1% sodium dodecyl sulfate (SDS), 5% dextran sulfate and 0.1 mg/mL denatured salmon sperm DNA at 50 °C for eighteen hours and wash twice at room temperature with 2X SSPE, 1% SDS for 5 min, then washing for 5 min at 50 °C in 0.2X SSPE, 1% SDS;

(d) hybridization in 50 mM Tris-HCl, pH 7.5, 1M NaCl, 1% sodium dodecyl sulfate (SDS), 5% dextran sulfate and 0.1 mg/mL denatured salmon sperm DNA at 50 °C for sixteen hours and wash twice at room temperature with 2X SSPE, 1% SDS for 5 min, then wash with fresh solution for 10 min, then wash for 5 min at 50 °C in 0.5X SSPE, 1% SDS;

(e) hybridization in 50 mM Tris, pH 7.6, 6X SSC, 5X Denhardt's, 0.5% sodium dodecyl sulfate (SDS), 100 µg denatured calf thymus DNA at 50°C overnight and wash with 6X SSC, 0.5% SDS at room temperature for 15 min, then wash twice with 2X SSC, 0.5% SDS at 45°C for 30 min each and then wash twice with 0.2X SSC, 0.5% SDS at 60 °C for 30 min each; or

(f) hybridization in 50 mM Tris-HCl, pH 7.5, 1M NaCl, 1% sodium dodecyl sulfate (SDS), 5% dextran sulfate and 0.1 mg/mL denatured salmon sperm DNA at 50 °C for eighteen hours and wash twice at room temperature with 2X SSPE, 1% SDS for 5 min, followed by washing for 5 min at 50°C in 0.2X SSPE, 1% SDS.

21. An isolated nucleic acid fragment of Claim 20 wherein said fragment is isolated from a plant selected from the group consisting of soybean, oilseed *Brassica* species, *Arabidopsis thaliana* and corn.

22. A chimeric gene capable of causing altered levels of linolenic acid in a transformed plant cell, the gene comprising a nucleic acid fragment of Claim 20, the fragment operably linked to regulatory sequences.

23. Plants containing the chimeric gene of claim 22.

24. Seeds obtained from the plants of claim 23.

25. Oil obtained from the seeds of claim 24.

26. A method of producing seed oil containing altered levels of linolenic (18:3) acid comprising:

- (a) transforming a plant cell of an oil-producing species with a chimeric gene of Claim 22;
- (b) growing fertile plants from the transformed plant cells of step (a);
- (c) screening progeny seeds from the fertile plants of step (b) for the desired levels of linolenic (18:3) acid; and
- (d) processing the progeny seed of step (c) to obtain seed oil containing altered levels of linolenic (18:3) acid.

26. A method of claim 25 wherein said plant cell of an oil-producing species is selected from the group consisting of *Arabidopsis thaliana*, soybean, oilseed *Brassica napus*, sunflower, cotton, peanut, and corn.

27. The method of claim 25 wherein said plant cell of an oil-producing species is selected from the group consisting of *Arabidopsis thaliana*, soybean, oilseed *Brassica napus*, sunflower, cotton and corn. --

#### In the Specification

Please amend the specification by adding after the Title at line 3

--This is a continuation of application number 08/244,205 filed August 26, 1994, now allowed, which is a 371 of PCT/US92/10284, filed December 3, 1992, now abandoned, which is a Continuation-in-Part of application number 07/804,259 filed December 4, 1991, now abandoned.--

#### Remarks

This is a continuation under 37 CFR §1.53 (b).


Claims 21 - 27 concern nucleic acid fragments useful in inhibition of plant target genes by antisense or cosuppression technologies. Support for this can be found in the specification on page 53 starting at line 11 through line 30 on page 54. Support for the hybridization conditions can be found in the specification on pages 40, 41, 42, 77, 79, 80 and 82. Thus, no new matter is being added.

Enclosed herewith is an Information Disclosure Statement setting forth all references which have been cited by Applicants or the Examiner in connection with

the prosecution of application number 08/244,205 which is a 371 of  
PCT/US92/10284, filed December 3, 1992, now abandoned.

Please charge any fees which are required in connection with the filing of this  
Preliminary Amendment, Information Disclosure Statement to Deposit Account No.  
04-1928 (E. I. du Pont de Nemours and Company).

Respectfully submitted,



Lynne M. Christenbury  
Attorney for Applicants  
Registration No. 30,971  
Telephone: 302-992-5481

09216035-122300

TITLE

## FATTY ACID DESATURASE GENES FROM PLANTS

## FIELD OF THE INVENTION

The invention relates to the preparation and use of  
5 nucleic acid fragments encoding fatty acid desaturase  
enzymes to modify plant lipid composition.

## BACKGROUND OF THE INVENTION

Plant lipids have a variety of industrial and nutritional uses and are central to plant membrane function and climatic adaptation. These lipids represent a vast array of chemical structures, and these structures determine the physiological and industrial properties of the lipid. Many of these structures result either directly or indirectly from metabolic processes that alter the degree of unsaturation of the lipid. Different metabolic regimes in different plants produce these altered lipids, and either domestication of exotic plant species or modification of agronomically adapted species is usually required to economically produce large amounts of the desired lipid.

Plant lipids find their major use as edible oils in the form of triacylglycerols. The specific performance and health attributes of edible oils are determined largely by their fatty acid composition. Most vegetable oils derived from commercial plant varieties are composed primarily of palmitic (16:0), stearic (18:0), oleic (18:1), linoleic (18:2) and linolenic (18:3) acids. Palmitic and stearic acids are, respectively, 16- and 18-carbon-long, saturated fatty acids. Oleic, linoleic, and linolenic acids are 18-carbon-long, unsaturated fatty acids containing one, two, and three double bonds, respectively. Oleic acid is referred to as a mono-unsaturated fatty acid, while linoleic and linolenic acids are referred to as poly-unsaturated fatty acids. The relative amounts of saturated and

unsaturated fatty acids in commonly used, edible vegetable oils are summarized below (Table 1):

TABLE 1

Percentages of Saturated and Unsaturated Fatty Acids in the Oils of Selected Oil Crops

	<u>Saturated</u>	<u>Mono-unsaturated</u>	<u>Poly-unsaturated</u>
<u>Canola</u>	6%	58%	36%
<u>Soybean</u>	15%	24%	61%
<u>Corn</u>	13%	25%	62%
<u>Peanut</u>	18%	48%	34%
<u>Safflower</u>	9%	13%	78%
<u>Sunflower</u>	9%	41%	51%
<u>Cotton</u>	30%	19%	51%

Many recent research efforts have examined the role that saturated and unsaturated fatty acids play in reducing the risk of coronary heart disease. In the past, it was believed that mono-unsaturates, in contrast to saturates and poly-unsaturates, had no effect on serum cholesterol and coronary heart disease risk. Several recent human clinical studies suggest that diets high in mono-unsaturated fat and low in saturated fat may reduce the "bad" (low-density lipoprotein) cholesterol while maintaining the "good" (high-density lipoprotein) cholesterol (Mattson et al., Journal of Lipid Research (1985) 26:194-202).

A vegetable oil low in total saturates and high in mono-unsaturates would provide significant health benefits to consumers as well as economic benefits to oil processors. As an example, canola oil is considered a very healthy oil. However, in use, the high level of poly-unsaturated fatty acids in canola oil renders the oil unstable, easily oxidized, and susceptible to development of disagreeable odors and flavors (Gailliard, 1980, Vol. 4, pp. 85-116 In: Stumpf, P. K., Ed., The Biochemistry of Plants, Academic Press, New



York). The levels of poly-unsaturates may be reduced by hydrogenation, but the expense of this process and the concomitant production of nutritionally questionable trans isomers of the remaining unsaturated fatty acids  
 5 reduces the overall desirability of the hydrogenated oil (Mensink et al., New England J. Medicine (1990) N323: 439-445). Similar problems exist with soybean and corn oils.

For specialized uses, high levels of poly-  
 10 unsaturates can be desirable. Linoleate and linolenate are essential fatty acids in human diets, and an edible oil high in these fatty acids can be used for nutritional supplements, for example in baby foods. Linseed oil, derived from the Flax plant (*Linum*  
 15 *usitatissimum*), contains over 50% linolenic acid and has widespread use in domestic and industrial coatings since the double bonds of the fatty acids react rapidly with oxygen to polymerize into a soft and flexible film. Although the oil content of flax is comparable to canola  
 20 (around 40% dry weight of seed), high yields are only obtained in warm temperatures or subtropical climates. In the USA flax is highly susceptible to rust infection. It will be commercially useful if a crop such as soybean or canola could be genetically transformed by the  
 25 appropriate desaturase gene(s) to synthesize oils with a high linolenic acid content.

Mutation-breeding programs have met with some success in altering the levels of poly-unsaturated fatty acid levels found in the edible oils of agronomic  
 30 species. Examples of commercially grown varieties are high (85%) oleic sunflower and low (2%) linolenic flax (Knowles, (1980) pp. 35-38 In: Applewhite, T. H., Ed., World Conference on Biotechnology for the Fats and Oils Industry Proceedings, American Oil Chemists' Society).  
 35 Similar commercial progress with the other plants shown

00210035-122320

in Table 1 has been largely elusive due to the difficult nature of the procedure and the pleiotropic effects of the mutational regime on plant hardiness and yield potential.

- 5       The biosynthesis of the major plant lipids has been the focus of much research (Browse et al., Ann. Rev. Plant Physiol. Mol. Biol. (1991) 42:467-506). These studies show that, with the notable exception of the soluble stearyl-acyl carrier protein desaturase, the
- 10       controlling steps in the production of unsaturated fatty acids are largely catalyzed by membrane-associated fatty acid desaturases. Desaturation reactions occur in plastids and in the endoplasmic reticulum using a variety of substrates including galactolipids,
- 15       sulfolipids, and phospholipids. Genetic and physiological analyses of Arabidopsis thaliana nuclear mutants defective in various fatty acid desaturation reactions indicates that most of these reactions are catalyzed by enzymes encoded at single genetic loci in
- 20       the plant. The analyses show further that the different defects in fatty acid desaturation can have profound and different effects on the ultra-structural morphology, cold sensitivity, and photosynthetic capacity of the plants (Ohlrogge, et al., Biochim. Biophys. Acta (1991)
- 25       1082:1-26). However, biochemical characterization of the desaturase reactions has been meager. The instability of the enzymes and the intractability of their proper assay has largely limited researchers to investigations of enzyme activities in crude membrane
- 30       preparations. These investigations have, however, demonstrated the role of delta-12 desaturase and delta-15 desaturase activities in the production of linoleate and linolenate from 2-oleoyl-phosphatidylcholine and 2-linoleoyl-phosphatidylcholine,
- 35       respectively (Wang et al., Plant Physiol. Biochem.

00010025-12200

(1988) 26:777-792). Thus, modification of the activities of these enzymes represents an attractive target for altering the levels of lipid unsaturation by genetic engineering.

- 5 Genes from plants for stearoyl-acyl carrier protein desaturase, the only soluble fatty acid desaturase known, have been described (Thompson, et al., Proc. Natl. Acad. Sci. U.S.A. (1991) 88:2578-2582; Shanklin et al., Proc. Natl. Acad. Sci. USA (1991) 88:2510-2514).
- 10 Stearoyl-coenzyme-A desaturase genes from yeast, rat, and mice have also been described (Stukey, et al., J. Biol. Chem. (1990) 265:20144-20149; Thiede, et al., J. Biol. Chem. (1986) 261:13230-13235; Kaestner, et al., J. Biol. Chem. (1989) 264:14755-1476). No evidence exists
- 15 in the public art that describes the isolation of fatty acid desaturases other than stearoyl-ACP desaturases from higher plants or their corresponding genes. A fatty acid desaturase gene from the cyanobacterium, Synechocystis PCC 6803, has also been described (Wada, et al., Nature (1990) 347:200-203). This gene encodes a
- 20 fatty acid desaturase, designated des A, that catalyzes the conversion of oleic acid at the 1 position of galactolipids to linoleic acid. However, these genes have not proven useful for isolating plant fatty acid
- 25 desaturases other than stearoyl-ACP desaturase via sequence-dependent protocols, and the present art does not indicate how to obtain plant fatty acid desaturases other than stearoyl-ACP desaturases or how to obtain fatty acid desaturase-related enzymes. Thus, the
- 30 present art does not teach how to obtain glycerolipid desaturases from plants. Furthermore, there is no evidence that a method to control the nature and levels of unsaturated fatty acids in plants using nucleic acids encoding fatty acid desaturases other than stearoyl-ACP
- 35 desaturase is known in the art.

The biosynthesis of the minor plant lipids has been less well studied. While hundreds of different fatty acids have been found, many from the plant kingdom, only a tiny fraction of all plants have been surveyed for their lipid content (Gunstone, et al., Eds., (1986) The Lipids Handbook, Chapman and Hall Ltd., Cambridge). Accordingly, little is known about the biosynthesis of these unusual fatty acids and fatty acid derivatives. Interesting chemical features found in such fatty acids include, for example, allenic and conjugated double bonds, acetylenic bonds, trans double bonds, multiple double bonds, and single double bonds in a wide number of positions and configurations along the fatty acid chain. Similarly, many of the structural modifications found in unusual lipids (e.g., hydroxylation, epoxidation, cyclization, etc.) are probably produced via further metabolism following chemical activation of the fatty acid by desaturation or they involve a chemical reaction that is mechanistically similar to desaturation. For example, evidence for the mechanism of hydroxylation of fatty acids being part of a general mechanism of enzyme-catalyzed desaturation in eukaryotes has been obtained by substituting a sulfur atom in the place of carbon at the delta-9 position of stearic acid. When incubated with yeast cell extracts the thiostearate was converted to a 9-sulfoxide (Buist et al. (1987) Tetrahedron Letters 28:857-860). This sulfoxidation was specific for sulfur at the delta-9 position and did not occur in a yeast delta-9-desaturase deficient mutant (Buist & Marecak (1991) Tetrahedron Letters 32:891-894). The 9-sulfoxide is the sulfur analogue of 9-hydroxyoctadecastearate, the proposed intermediate of stearate desaturation. Thus fatty-acid desaturase cDNAs may serve as useful probes for cDNAs encoding fatty-acid hydroxylases and other cDNAs which encode enzymes with

- reaction mechanisms similar to fatty-acid desaturation. Many of these fatty acids and derivatives having such features within their structure could prove commercially useful if an agronomically viable species could be
- 5 induced to synthesize them by introduction of a gene encoding the appropriate desaturase.

#### SUMMARY OF THE INVENTION

- Applicants have discovered a means to control the nature and levels of unsaturated fatty acids in plants.
- 10 Nucleic acid fragments from glycerolipid desaturase cDNAs or genes are used to create chimeric genes. The chimeric genes may be used to transform various plants to modify the fatty acid composition of the plant or the oil produced by the plant. More specifically, one
- 15 embodiment of the invention is an isolated nucleic acid fragment comprising a nucleotide sequence encoding a plant delta-15 fatty acid desaturase or a fatty acid desaturase-related enzyme with an amino acid identity of 50%, 65%, 90% or greater to the polypeptide encoded by
- 20 SEQ ID NOS:1, 4, 6, 8, 10, 12, 14, or 16. The isolated fragment in these embodiments is isolated from a plant selected from the group consisting of soybean, oilseed Brassica species, *Arabidopsis thaliana* and corn.

- Another embodiment of this invention involves the
- 25 use of these nucleic acid fragments in sequence-dependent protocols. Examples include use of the fragments as hybridization probes to isolate other glycerolipid desaturase cDNAs or genes. A related embodiment involves using the disclosed sequences for
- 30 amplification of DNA fragments encoding other glycerolipid desaturases.

- Another aspect of this invention involves chimeric genes capable of causing altered levels of the linolenic acid in a transformed plant cell, the gene comprising
- 35 nucleic acid fragments encoding encoding a plant

delta-15 fatty acid desaturase or a fatty acid desaturase-related enzyme with an amino acid identity of 50%, 65%, 90% or greater to the polypeptide encoded by SEQ ID NOS:1, 4, 6, 8, 10, 12, 14, or 16 operably linked  
5 in suitable orientation to suitable regulatory sequences. Preferred are those chimeric genes which incorporate nucleic acid fragments encoding delta-15 fatty acid desaturase cDNAs or genes. Plants and oil from seeds of plants containing the chimeric genes  
10 described are also claimed.

Yet another embodiment of the invention involves a method of producing seed oil containing altered levels of linolenic (18:3) acid comprising: (a) transforming a plant cell with a chimeric gene described above; (b)  
15 growing fertile plants from the transformed plant cells of step (a); (c) screening progeny seeds from the fertile plants of step (b) for the desired levels of linolenic (18:3) acid, and (d) processing the progeny seed of step (c) to obtain seed oil containing altered  
20 levels of the unsaturated fatty acids. Preferred plant cells and oils are derived from soybean, rapeseed, sunflower, cotton, cocoa, peanut, safflower, coconut, flax, oil palm, and corn. Preferred methods of transforming such plant cells would include the use of  
25 Ti and Ri plasmids of *Agrobacterium*, electroporation, and high-velocity ballistic bombardment.

The invention also is embodied in a method of breeding plant species to obtain altered levels of poly-unsaturated fatty acids, specifically linolenic (18:3)  
30 acid in seed oil of oil-producing plants. This method involves (a) making a cross between two varieties of an oilseed plant differing in the linolenic acid trait; (b) making a Southern blot of restriction enzyme digested genomic DNA isolated from several progeny plants  
35 resulting from the cross of step (a); and (c)

hybridizing the Southern blot with the radiolabeled nucleic acid fragments encoding the claimed glycerolipid desaturases.

The invention is also embodied in a method of RFLP mapping that uses the isolated Arabidopsis thaliana delta-15 desaturase sequences described herein.

The invention is also embodied in plants capable of producing altered levels of glycerolipid desaturase by virtue of containing the chimeric genes described herein. Further, the invention is embodied by seed oil obtained from such plants.

The invention is also embodied in a method of RFLP mapping in a genomic RFLP marker comprising (a) making a cross between two varieties of plants; (b) making a Southern blot of restriction enzyme digested genomic DNA isolated from several progeny plants resulting from the cross of step (a); and (c) hybridizing the Southern blot with a radiolabelled nucleic acid fragments of the claimed fragments.

The invention is also embodied in a method to isolate nucleic acid fragments encoding fatty acid desaturases and fatty acid desaturase-related enzymes, comprising (a) comparing SEQ ID NOS:2, 5, 7, 9, 11, 13, 15 and 17 with other fatty acid desaturase polypeptide sequences; (b) identifying the conserved sequence(s) of 4 or more amino acids obtained in step a; (c) making region-specific nucleotide probe(s) or oligomer(s) based on the conserved sequences identified in step b; and d) using the nucleotide probe(s) or oligomers(s) of step c to isolate sequences encoding fatty acid desaturases and fatty-acid desaturase-related enzymes by sequence-dependent protocols. The product of the method of isolation method described is also part of the invention.

# BRIEF DESCRIPTION OF THE SEQUENCE DESCRIPTIONS

The invention can be more fully understood from the following detailed description and the Sequence Descriptions which form a part of this application. The Sequence Descriptions contain the one letter code for nucleotide sequence characters and the three letter code for amino acids in conformity with the IUPAC-IUB standard described in Nucleic Acids Research 13:3021-3030 (19085) and 37 C.F.R. 1.822 which are incorporated herein by reference.

SEQ ID NO:1 shows the complete 5' to 3' nucleotide sequence of 1350 base pairs of the Arabidopsis cDNA which encodes delta-15 desaturase in plasmid pCF3. Nucleotides 46 to 48 are the putative initiation codon of the open reading frame (nucleotides 46 to 1206). Nucleotides 1204 to 1206 are the termination codon. Nucleotides 1 to 45 and 1207 to 1350 are the 5' and 3' untranslated nucleotides, respectively. The 386 amino acid protein sequence in SEQ ID NO:1 is that deduced from the open reading frame.

SEQ ID NO:2 is the deduced peptide of the open-reading frame of SEQ ID NO:1.

SEQ ID NO:3 is a partial nucleotide sequence of the Arabidopsis genomic DNA insert in plasmid pF1 which shows the genomic sequence in the region of the Arabidopsis genome that encodes delta-15 desaturase. Nucleotides 68-255 are identical to nucleotides 1-188 of SEQ ID NO:1. Nucleotides 47 to 49 and 56 to 58 are termination codons in the same reading frame as the open reading frame in SEQ ID NO:1.

SEQ ID NO:4 shows the 5' to 3' nucleotide sequence of the insert in plasmid pACF2-2 of 1525 base pairs of the Arabidopsis thaliana cDNA that encodes a plastid delta-15 fatty acid desaturase. Nucleotides 10-12 and nucleotides 1348 to 1350 are, respectively, the putative



initiation codon and the termination codon of the open reading frame (nucleotides 10 to 1350). Nucleotides 1 to 9 and 1351 to 1525 are, respectively, the 5' and 3' untranslated nucleotides.

- 5        SEQ ID NO:5 is the deduced peptide of the open reading frame of SEQ ID NO:4.

- SEQ ID NO:6 shows the complete 5' to 3' nucleotide sequence of 1336 base pairs of the Brassica napus seed cDNA, found in plasmid pBNSF3-2, which encodes a  
10        microsomal delta-15 glycerolipid desaturase. Nucleotides 79 to 82 are the putative initiation codon of the open reading frame (nucleotides 79 to 1212). Nucleotides 1210 to 1212 are the termination codon. Nucleotides 1 to 78 and 1213 to 1336 are the 5' and 3'  
15        untranslated nucleotides respectively.

         SEQ ID NO:7 is the deduced peptide of the open reading frame of SEQ ID NO:6.

- SEQ ID NO:8 is the complete 5' to 3' nucleotide sequence of 1416 base pairs of the Brassica napus seed  
20        cDNA found in plasmid pBNSFd-2 which encodes a plastid delta-15 glycerolipid desaturase. Nucleotides 1 to 1215 correspond to a continuous open reading frame of 404 amino acids. Nucleotides 1213 to 1215 are the termination codon. Nucleotides 1215 to 1416 are the 3'  
25        untranslated nucleotides.

         SEQ ID NO:9 is the deduced peptide of the open reading frame of SEQ ID NO:8.

- SEQ ID NO:10 is the complete nucleotide sequence of the soybean (glycine max) microsomal delta-15 desaturase  
30        cDNA, found in plasmid pXF1, which the 2184 nucleotides of this sequence contain both the coding sequence and the 5' and 3' non-translated regions of the cDNA. Nucleotides 855 to 857 are the putative initiation codon of the open reading frame (nucleotides 855 to 2000).  
35        Nucleotides 1995 to 1997 are the termination codon.

Nucleotides 1 to 854 and 1998 to 2184 are the 5' and 3' untranslated nucleotides respectively. The 380 amino acid protein sequence in SEQ ID NO:7 is that deduced from the open reading frame.

- 5        SEQ ID NO:11 is the deduced peptide of the open reading frame in SEQ ID NO:10.

- SEQ ID NO:12 is the complete 5' to 3' nucleotide sequence of 1676 base pairs of the soybean (*Glycine max*) seed cDNA found in plasmid pSFD-118bwp which encodes a  
10        soybean plastid delta-15 desaturase. Nucleotides 169 to 1530 correspond to a continuous open reading frame of 453 amino acids. Nucleotides 169 to 171 are the putative initiation codon of the open reading frame. Nucleotides 1528 to 1530 are the termination codon.  
15        Nucleotides 1531 to 1676 are the 3' untranslated nucleotides. Nucleotides 169 to 382 encode the putative plastid transit peptide, based on comparison of the deduced peptide with the soybean microsomal delta-15 peptide.

- 20        SEQ ID NO:13 is the deduced peptide of the open reading frame in SEQ ID NO:12.

- SEQ ID NO:14 is the complete nucleotide sequence of a 396 bp polymerase chain reaction product derived from corn seed mRNA that is found in the insert of plasmid  
25        pPCR20. Nucleotides 1 to 31 and 364 to 396 correspond to the amplification primers described in SEQ ID NO:18 and SEQ ID NO:19, respectively. Nucleotides 31 to 363 encode an internal region of a corn seed delta-15 desaturase that is 61.9% identical to the region between  
30        amino acids 137 and 249 of the Brassica napus delta-15 desaturase peptide sequence shown in SEQ ID NO:7.

          SEQ ID NO:15 is the deduced amino acid sequence of SEQ ID NO:14.

- 35        SEQ ID NO:16 shows the partial composite 5' to 3' nucleotide sequence of 472 bp derived from the inserts

in plasmids pFadx-2 and pYacp7 for Arabidopsis thaliana cDNA that encodes a plastid delta-15 fatty acid desaturase. Nucleotides 2-4 and nucleotides 468 to 470 are, respectively, the first and the last codons in the open reading frame.

SEQ ID NO:17 is deduced partial peptide sequence of the open reading frame in SEQ ID NO:16.

SEQ ID NO:18 One hundred and twenty eight fold degenerate sense 31-mer PCR primer. Nucleotides 1 to 8 correspond to the Bam H1 restriction enzyme recognition sequence. Nucleotides 9 to 137 correspond to amino acid residues 130 to 137 of SEQ ID NO:6 with a deoxyinosine base at nucleotide 11.

SEQ ID NO:19 Two thousand and forty eight-fold degenerate antisense 35-mer PCR primer. Nucleotides 1 to 8 correspond to the Bam H1 restriction enzyme recognition sequence. Nucleotides 9 to 35 correspond to amino acid residues 249 to 256 of SEQ ID NO:6 with a deoxyinosine base at nucleotide 15.

SEQ ID NO:20 Sixteen-fold degenerate sense 36-mers made to amino acid residues 97-108 in SEQ ID NO:2.

SEQ ID NO:21 Sixteen-fold degenerate sense 36-mers made to amino acid residues 97-108 in SEQ ID NO:2.

SEQ ID NO:22 Seventy two-fold degenerate sense 18-mers made to amino acid residues 100-105 in SEQ ID NO:2.

SEQ ID NO:23 Seventy two-fold degenerate sense 18-mers made to amino acid residues 100-105 in SEQ ID NO:2.

SEQ ID NO:24 Seventy two-fold degenerate antisense 18-mers made to amino acid residues 299-304 in SEQ ID NO:2.

SEQ ID NO:25 Seventy two-fold degenerate antisense 18-mers made to amino acid residues 299-304 in SEQ ID NO:2.

SEQ ID NO:26 Seventy two-fold degenerate antisense  
18-mers made to amino acid residues 304-309 in SEQ ID  
NO:2.

5 SEQ ID NO:27 Seventy two-fold degenerate antisense  
18-mers made to amino acid residues 304-309 in SEQ ID  
NO:2.

SEQ ID NO:28 Sixteen-fold degenerate sense 36-mers  
made to amino acid residues 97-108 in SEQ ID NO:2.

10 SEQ ID NO:29 Sixteen-fold degenerate sense 36-mers  
made to amino acid residues 97-108 in SEQ ID NO:2.

SEQ ID NO:30 Sixty four-fold degenerate antisense  
38-mers made to amino acid residues 299-311 in SEQ ID  
NO:2.

15 SEQ ID NO:31 Sixty four-fold degenerate antisense  
38-mers made to amino acid residues 299-311 in SEQ ID  
NO:2.

SEQ ID NO:32 A 135-mer made as an antisense strand  
to amino acid residues 97-141 in SEQ ID NO:2.

#### DETAILED DESCRIPTION OF THE INVENTION

20 Applicants have isolated nucleic acid fragments  
that encode plant fatty acid desaturases and that are  
useful in modifying fatty acid composition in oil-  
producing species by transformation.

25 Thus, transfer of the nucleic acid fragments of the  
invention or a part thereof that encodes a functional  
enzyme, along with suitable regulatory sequences that  
direct the transcription of their mRNA, into a living  
cell will result in the production or over-production of  
plant fatty acid desaturases and will result in  
30 increased levels of unsaturated fatty acids in cellular  
lipids, including triacylglycerols.

Transfer of the nucleic acid fragments of the  
invention or a part thereof, along with suitable  
regulatory sequences that direct the transcription of  
35 their antisense RNA, into plants will result in the

inhibition of expression of the endogenous fatty acid desaturase that is substantially homologous with the transferred nucleic acid fragment and will result in decreased levels of unsaturated fatty acids in cellular lipids, including triacylglycerols.

Transfer of the nucleic acid fragments of the invention or a part thereof, along with suitable regulatory sequences that direct the transcription of their mRNA, into plants may result in inhibition by cosuppression of the expression of the endogenous fatty acid desaturase gene that is substantially homologous with the transferred nucleic acid fragment and may result in decreased levels of unsaturated fatty acids in cellular lipids, including triacylglycerols.

The nucleic acid fragments of the invention can also be used as restriction fragment length polymorphism (RFLP) markers in Arabidopsis genetic mapping and plant breeding programs.

The nucleic acid fragments of the invention or oligomers derived therefrom can also be used to isolate other related glycerolipid desaturase genes using DNA, RNA, or a library of cloned nucleotide sequences from the same or different species by well known sequence-dependent protocols, including, for example, methods of nucleic acid hybridization and amplification by the polymerase chain reaction.

#### Definitions

In the context of this disclosure, a number of terms shall be used. The term "fatty acid desaturase" used herein refers to an enzyme which catalyzes the breakage of a carbon-hydrogen bond and the introduction of a carbon-carbon double bond into a fatty acid molecule. The fatty acid may be free or esterified to another molecule including, but not limited to, acyl-carrier protein, coenzyme A, sterols and the glycerol

moiety of glycerolipids. The term "glycerolipid desaturases" used herein refers to a subset of the fatty acid desaturases that act on fatty acyl moieties esterified to a glycerol backbone. "Delta-12 desaturase" refers to a fatty acid desaturase that catalyzes the formation of a double bond between carbon positions 6 and 7 (numbered from the methyl end), (i.e., those that correspond to carbon positions 12 and 13 (numbered from the carbonyl carbon) of an 18 carbon-long fatty acyl chain or carbon positions 10 and 11 (numbered from the carbonyl carbon) of a 16 carbon-long fatty acyl chain). "Delta-15 desaturase" refers to a fatty acid desaturase that catalyzes the formation of a double bond between carbon positions 3 and 4 (numbered from the methyl end), (i.e., those that correspond to carbon positions 15 and 16 (numbered from the carbonyl carbon) of an 18 carbon-long fatty acyl chain and carbon positions 13 and 14 (numbered from the carbonyl carbon) of a 16 carbon-long fatty acyl chain). Examples of fatty acid desaturases include, but are not limited to, the microsomal delta-12 and delta-15 desaturases that act on phosphatidylcholine lipid substrates; the chloroplastic delta-12 and delta-15 desaturases that act on phosphatidyl glycerol and galactolipids; and other desaturases that act on such fatty acid substrates such as phospholipids, galactolipids, and sulfolipids. "Microsomal desaturase" refers to the cytoplasmic location of the enzyme, while "chloroplast desaturase" and "plastid desaturase" refer to the plastid location of the enzyme. These fatty acid desaturases may be found in a variety of organisms including, but not limited to, higher plants, diatoms, and various eukaryotic and prokaryotic microorganisms such as fungi and photosynthetic bacteria and algae. The term "homologous fatty acid desaturases" refers to fatty acid

desaturases that catalyze the same desaturation on the same lipid substrate. Thus, microsomal delta-15 desaturases, even from different plant species, are homologous fatty acid desaturases. The term

5 "heterologous fatty acid desaturases" refers to fatty acid desaturases that catalyze desaturations at different positions and/or on different lipid substrates. Thus, for example, microsomal delta-12 and delta-15 desaturases, which act on phosphatidylcholine

10 lipids, are heterologous fatty acid desaturases, even when from the same plant. Similarly, microsomal delta-15 desaturase, which acts on phosphatidylcholine lipids, and chloroplast delta-15 desaturase, which acts on galactolipids, are heterologous fatty acid

15 desaturases, even when from the same plant. It should be noted that these fatty acid desaturases have never been isolated and characterized as proteins. Accordingly the terms such as "delta-12 desaturase" and "delta-15 desaturase" are used as a convenience to

20 describe the proteins encoded by nucleic acid fragments that have been isolated based on the phenotypic effects caused by their disruption. The term "fatty acid desaturase-related enzyme" refers to enzymes whose catalytic product may not be a carbon-carbon double bond

25 but whose mechanism of action is similar to that of a fatty acid desaturase (that is, catalysis of the displacement of a carbon-hydrogen bond of a fatty acid chain to form a fatty-hydroxyacyl intermediate or end-product). This term is different from "related fatty

30 acid desaturases", which refers to structural similarities between fatty acid desaturases.

The term "nucleic acid" refers to a large molecule which can be single-stranded or double-stranded, composed of monomers (nucleotides) containing a sugar, a

35 phosphate and either a purine or pyrimidine. A "nucleic

acid fragment" is a fraction of a given nucleic acid molecule. In higher plants, deoxyribonucleic acid (DNA) is the genetic material while ribonucleic acid (RNA) is involved in the transfer of the information in DNA into proteins. A "genome" is the entire body of genetic material contained in each cell of an organism. The term "nucleotide sequence" refers to the sequence of DNA or RNA polymers, which can be single- or double-stranded, optionally containing synthetic, non-natural or altered nucleotide bases capable of incorporation into DNA or RNA polymers. The term "oligomer" refers to short nucleotide sequences, usually up to 150 bases long. "Region-specific nucleotide probes" refers to isolated nucleic acid fragments derived from a cDNA or gene using a knowledge of the amino acid regions conserved between different fatty-acid desaturases which may be used to isolate cDNAs or genes for other fatty-acid desaturases or fatty acid desaturase-related enzymes using sequence dependent protocols. As used herein, the term "homologous to" refers to the relatedness between the nucleotide sequence of two nucleic acid molecules or between the amino acid sequences of two protein molecules. Estimates of such homology are provided by either DNA-DNA or DNA-RNA hybridization under conditions of stringency as is well understood by those skilled in the art (Hames and Higgins, Eds. (1985) *Nucleic Acid Hybridisation*, IRL Press, Oxford, U.K.); or by the comparison of sequence similarity between two nucleic acids or proteins, such as by the method of Needleman et al. (*J. Mol. Biol.* (1970) 48:443-453). As used herein, "substantially homologous" refers to nucleotide sequences that have more than 90% overall identity at the nucleotide level with the coding region of the claimed sequence, such as genes and pseudo-genes corresponding to the coding



regions. The nucleic acid fragments described herein include molecules which comprise possible variations, both man-made and natural, such as but not limited to (a) those that involve base changes that do not cause a change in an encoded amino acid, or (b) which involve base changes that alter an amino acid but do not affect the functional properties of the protein encoded by the DNA sequence, (c) those derived from deletions, rearrangements, amplifications, random or controlled mutagenesis of the nucleic acid fragment, and (d) even occasional nucleotide sequencing errors.

"Gene" refers to a nucleic acid fragment that expresses a specific protein, including regulatory sequences preceding (5' non-coding) and following (3' non-coding) the coding region. "Fatty acid desaturase gene" refers to a nucleic acid fragment that expresses a protein with fatty acid desaturase activity. "Native" gene refers to an isolated gene with its own regulatory sequences as found in nature. "Chimeric gene" refers to a gene that comprises heterogeneous regulatory and coding sequences not found in nature. "Endogenous" gene refers to the native gene normally found in its natural location in the genome and is not isolated. A "foreign" gene refers to a gene not normally found in the host organism but that is instead introduced by gene transfer. "Pseudo-gene" refers to a genomic nucleotide sequence that does not encode a functional enzyme.

"Coding sequence" refers to a DNA sequence that codes for a specific protein and excludes the non-coding sequences. It may constitute an "uninterrupted coding sequence", i.e., lacking an intron or it may include one or more introns bounded by appropriate splice junctions. An "intron" is a nucleotide sequence that is transcribed in the primary transcript but that is removed through cleavage and re-ligation of the RNA within the cell to

create the mature mRNA that can be translated into a protein.

"Initiation codon" and "termination codon" refer to a unit of three adjacent nucleotides in a coding sequence that specifies initiation and chain termination respectively, of protein synthesis (mRNA translation). "Open reading frame" refers to the coding sequence uninterrupted by introns between initiation and termination codons that encodes an amino acid sequence.

"RNA transcript" refers to the product resulting from RNA polymerase-catalyzed transcription of a DNA sequence. When the RNA transcript is a perfect complementary copy of the DNA sequence, it is referred to as the primary transcript or it may be a RNA sequence derived from posttranscriptional processing of the primary transcript and is referred to as the mature RNA. "Messenger RNA (mRNA)" refers to the RNA that is without introns and that can be translated into protein by the cell. "cDNA" refers to a double-stranded DNA that is complementary to and derived from mRNA. "Sense" RNA refers to RNA transcript that includes the mRNA. "Antisense RNA" refers to a RNA transcript that is complementary to all or part of a target primary transcript or mRNA and that blocks the expression of a target gene by interfering with the processing, transport and/or translation of its primary transcript or mRNA. The complementarity of an antisense RNA may be with any part of the specific gene transcript, i.e., at the 5' non-coding sequence, 3' non-coding sequence, introns, or the coding sequence. In addition, as used herein, antisense RNA may contain regions of ribozyme sequences that increase the efficacy of antisense RNA to block gene expression. "Ribozyme" refers to a catalytic RNA and includes sequence-specific endoribonucleases.

As used herein, "suitable regulatory sequences" refer to nucleotide sequences in native or chimeric genes that are located upstream (5'), within, and/or downstream (3') to the nucleic acid fragments of the invention, which control the expression of the nucleic acid fragments of the invention. The term "expression", as used herein, refers to the transcription and stable accumulation of the sense (mRNA) or the antisense RNA derived from the nucleic acid fragment(s) of the invention that, in conjunction with the protein apparatus of the cell, results in altered levels of the fatty acid desaturase(s). Expression or overexpression of the gene involves transcription of the gene and translation of the mRNA into precursor or mature fatty acid desaturase proteins. "Antisense inhibition" refers to the production of antisense RNA transcripts capable of preventing the expression of the target protein. "Overexpression" refers to the production of a gene product in transgenic organisms that exceeds levels of production in normal or non-transformed organisms. "Cosuppression" refers to the expression of a foreign gene which has substantial homology to an endogenous gene resulting in the suppression of expression of both the foreign and the endogenous gene. "Altered levels" refers to the production of gene product(s) in transgenic organisms in amounts or proportions that differ from that of normal or non-transformed organisms.

"Promoter" refers to a DNA sequence in a gene, usually upstream (5') to its coding sequence, which controls the expression of the coding sequence by providing the recognition for RNA polymerase and other factors required for proper transcription. In artificial DNA constructs promoters can also be used to transcribe antisense RNA. Promoters may also contain DNA sequences that are involved in the binding of

protein factors which control the effectiveness of transcription initiation in response to physiological or developmental conditions. It may also contain enhancer elements. An "enhancer" is a DNA sequence which can stimulate promoter activity. It may be an innate element of the promoter or a heterologous element inserted to enhance the level and/or tissue-specificity of a promoter. "Constitutive promoters" refers to those that direct gene expression in all tissues and at all times. "Tissue-specific" or "development-specific" promoters as referred to herein are those that direct gene expression almost exclusively in specific tissues, such as leaves or seeds, or at specific development stages in a tissue, such as in early or late embryogenesis, respectively.

The "3' non-coding sequences" refers to the DNA sequence portion of a gene that contains a polyadenylation signal and any other regulatory signal capable of affecting mRNA processing or gene expression. The polyadenylation signal is usually characterized by affecting the addition of polyadenylic acid tracts to the 3' end of the mRNA precursor.

The term "Transit Peptide" refers to the N-terminal extension of a protein that serves as a signal for uptake and transport of that protein into an organelle such as a plastid or mitochondrion.

"Transformation" herein refers to the transfer of a foreign gene into the genome of a host organism and its genetically stable inheritance. "Restriction fragment length polymorphism" refers to different sized restriction fragment lengths due to altered nucleotide sequences in or around variant forms of genes. "Fertile" refers to plants that are able to propagate sexually.

"Oil-producing species" herein refers to plant species which produce and store triacylglycerol in specific organs, primarily in seeds. Such species include soybean (Glycine max), rapeseed and canola (including Brassica napus, B. campestris), sunflower (Helianthus annus), cotton (Gossypium hirsutum), corn (Zea mays), cocoa (Theobroma cacao), safflower (Carthamus tinctorius), oil palm (Elaeis guineensis), coconut palm (Cocos nucifera), flax (Linum usitatissimum), castor (Ricinus communis) and peanut (Arachis hypogaea). The group also includes non-agronomic species which are useful in developing appropriate expression vectors such as tobacco, rapid cycling Brassica species, and Arabidopsis thaliana, and wild species which may be a source of unique fatty acids.

"Sequence-dependent protocols" refer to techniques that rely on a nucleotide sequence for their utility. Examples of sequence-dependent protocols include, but are not limited to, the methods of nucleic acid and oligomer hybridization and methods of DNA and RNA amplification such as are exemplified in various uses of the polymerase chain reaction. "PCR product" refers to the DNA product obtained through polymerase chain reaction.

Various solutions used in the experimental manipulations are referred to by their common names such as "SSC", "SSPE", "Denhardt's solution", etc. The composition of these solutions may be found by reference to Appendix B of Sambrook, et al. (Molecular Cloning, A Laboratory Manual, 2nd ed. (1989), Cold Spring Harbor Laboratory Press).

00210035.122393

T-DNA Mutagenesis and Identification of an  
Arabidopsis Mutant Defective in Delta-15 Desaturation

In T-DNA mutagenesis (Feldmann, et al., Science  
(1989) 243:1351-1354), the integration of T-DNA in the  
5 genome can interrupt normal expression of the gene at or  
near the site of the integration. If the resultant  
mutant phenotype can be detected and shown genetically  
to be tightly linked to the T-DNA insertion, then the  
"tagged" locus and its wild type counterpart can be  
10 readily isolated by molecular cloning by one skilled in  
the art.

Arabidopsis thaliana seeds were transformed by  
Agrobacterium tumefaciens C58Clrif strain harboring the  
avirulent Ti-plasmid pGV3850::pAK1003 that has the T-DNA  
15 region between the left and right T-DNA borders replaced  
by the origin of replication region and ampicillin  
resistance gene of plasmid pBR322, a bacterial kanamycin  
resistance gene, and a plant kanamycin resistance gene  
(Feldmann, et al., Mol. Gen. Genetics (1987) 208:1-9).  
20 Plants from the treated seeds were self-fertilized and  
the resultant progeny seeds, germinated in the presence  
of kanamycin, were self-fertilized to give rise to a  
population, designated T3, that was segregating for  
T-DNA insertions. T3 seeds from approximately 6000 T2  
25 plants were analyzed for fatty acid composition. One  
line, designated 3707, showed a reduced level of  
linolenic acid (18:3). One more round of self-  
fertilization of mutant line 3707 produced T4 progeny  
seeds. The ratio of 18:2/18:3 in seeds of the  
30 homogyzous mutant in T4 population was ca. 14; this  
ratio is ca 1.8 and ca. 23, respectively, in wild-type  
Arabidopsis and Arabidopsis fad 3 mutant [Lemieux et al.  
(1990) Theor. App. Gen. 80:234-240 ] obtained via  
chemical mutagenesis. These seeds were planted and 263  
35 individual plants were analyzed for the presence of

nopaline in leaf extracts. T5 seeds from these plants were further analyzed for fatty acid composition and the ability to germinate in the presence of kanamycin. The mutant fatty acid phenotype was found to segregate in a 1:2:1 ratio, as was germinability on kanamycin. Nopaline was found in all plants with an altered fatty acid phenotype, but not in wild type segregants. These results provided evidence that the locus controlling delta-15 desaturation was interrupted by T-DNA in mutant line 3707.

Isolation of *Arabidopsis* Genomic DNA  
Containing the Gene Controlling Delta-15 Desaturation

In order to isolate the gene controlling delta-15 desaturation from wild-type *Arabidopsis*, a T-DNA-plant DNA "junction" fragment containing a T-DNA border integrated into the host plant DNA was isolated from *Arabidopsis* mutant 3707. For this, genomic DNA from the mutant plant was isolated and completely digested by either Bam HI or Sal I restriction enzymes. In each case, one of the resultant fragments was expected to contain the origin of replication and ampicillin-resistance gene of pBR322 as well as the left T-DNA-plant DNA junction fragment. Such fragments were rescued as plasmids by ligating the digested genomic DNA fragments at a dilute concentration to facilitate self-ligation and then using the ligated fragments to transform *E. coli* cells. Ampicillin-resistant *E. coli* transformants were isolated and screened by colony hybridization to fragments containing either the left or the right T-DNA border. Of the 192 colonies obtained from the plasmid rescue of Sal I digested genomic DNA, 31 hybridized with the left T-DNA border fragment, 4 hybridized to the right T-DNA border fragment, and none hybridized to both. Of the 85 colonies obtained from the plasmid rescue of Bam HI digested genomic DNA, 63

hybridized to the left border and none to the right border. Restriction analysis of seven rescued plasmids that were obtained from the Bam HI digestion and that hybridized to the left T-DNA border showed that they

5 were indistinguishable and contained 1.4 kb of putative, flanking plant DNA. Restriction analysis of another rescued plasmid, pS1, that was obtained from the Sal I digestion and hybridized only to the left T-DNA border, showed that it contained 2.9 kb of putative, flanking

10 plant DNA. This flanking DNA had a Bam HI site and a Hind III site 1.4 kb and 2.2 kb, respectively, away from the left T-DNA border, suggesting that the 1.4 kb putative plant DNA in Bam HI rescued plasmids was contained within the 2.9 kb putative plant DNA in the

15 Sal I rescued plasmids. Southern blot analysis of wild type and mutant 3707 Arabidopsis genomic DNA using the radiolabeled 1.4 kb DNA fragment as the hybridization probe confirmed that this fragment contained plant DNA and that the T-DNA integration site was in a 2.8 kb

20 Bam HI, a 5.2 kb Hind III, a 3.5 kb Sal I, a 5.5 kb Eco RI, and an approximately 9 kb Cla I fragment of wild type Arabidopsis DNA. Nucleotide sequencing of plasmid pS1 with a primer made to a left T-DNA border sequence revealed that pS1 was colinear with the sequence of the

25 left T-DNA border (Yadav et al., Proc. Natl. Acad. Sci. USA (1982) 79:6322-6326) up to nucleotide position 65, which is in the T-DNA border repeats. Approximately 800 bp of additional sequence in pS1 beyond the T-DNA-plant DNA junction, that is, in the plant DNA adjoining the

30 left T-DNA border, showed no significant homology to the T-DNA of pGV3850::pAK1003 and no significant open reading frame.

The nucleic acid fragment from wild-type Arabidopsis corresponding to the plant DNA flanking

35 T-DNA in the line 3707 was isolated by screening a



lambda phage Arabidopsis thaliana genomic library with the 1.4 kb plant DNA isolated from the rescued plasmids as a hybridization probe. Seven positively-hybridizing genomic clones were isolated that fell in one of five classes based on partial restriction mapping. While their average insert size was approximately 15 kb, taken together they spanned a total of approximately 40 kb of genomic DNA. A combination of restriction and Southern analyses revealed that the five clones overlapped the site of integration of the left border of the T-DNA and that there was no detectable rearrangement of plant DNA in the rescued plasmids as compared to that in the wild type genomic plant DNA. One of these lambda phage clones, designated 1111, was representative of the recovered clones and contained an approximately 20 kb genomic DNA insert which was more or less symmetrically arranged around the site of insertion of the left border of the T-DNA. This clone was deposited on November 27, 1991 with the American Type Culture Collection of Rockville, Maryland under the provisions of the Budapest Treaty and bears accession number ATCC 75167.

#### Isolation of Arabidopsis Delta-15

##### Desaturase cDNA

A 5.2 kb Hind III fragment containing wild-type genomic DNA, which hybridized to the 1.4 kb flanking plant DNA recovered from line 3707 and which was interrupted near its middle by the T-DNA insertion in line 3707, was isolated from lambda phage clone 41A1 and cloned into the Hind III site of the pBluescript SK vector (Stratagene) by standard cloning procedures described in Sambrook et al., Molecular Cloning, A Laboratory Manual, 2nd ed. (1989, Cold Spring Harbor Laboratory Press). The resultant plasmid was designated pF1. The isolated 5.2 kb Hind III fragment was also used as a radiolabeled hybridization probe to screen a

cdNA library made to poly A<sup>+</sup> mRNA from 3-day-old etiolated *Arabidopsis thaliana* (ecotype Columbia) seedling hypocotyls in a lambda ZAP II vector (Stratagene). Of the several positively-hybridizing

5 plaques, four strongly-hybridizing ones were subjected to plaque purification. Sequences of the pBluescript (Stratagene) vector, including the cdNA inserts, from each of the purified phage stocks were excised in the presence of a helper phage. The resultant phagemids

10 were used to infect *E. coli* cells which yielded double-stranded plasmids, pCF1, pCF2, pCF3, and pCF4. All four were shown to contain at least one approximately 1.3 to 1.4 kb Not I insert fragment (Not I/Eco RI adaptors were used in the preparation of the cdNA library) which

15 hybridized to the same region of wild-type plant genomic DNA present in the isolated phage clones. This region, which was near the site of integration of the left T-DNA border in line 3707, was on the side of the T-DNA insertion opposite to that of the plant DNA flanking the

20 left T-DNA border isolated previously via plasmid rescue. Partial sequence determination of the different cDNAs revealed common identity. Since multiple versions of only one type of cdNA were obtained from a cdNA library made from etiolated tissue which is expected to

25 express delta-15 desaturation, and since these cDNAs hybridized to the genomic DNA that corresponds to the site of T-DNA integration in line 3707 which had a high linoleic acid/low linolenic acid phenotype, Applicants were lead to conclude that the T-DNA in line 3707

30 interrupted the normal expression of the gene encoding delta-15 desaturase. The complete nucleotide sequence of one cdNA, designated pCF3, was determined and is shown as SEQ ID NO:1. It reveals an open reading frame that encodes a 386 amino acid polypeptide. One of the

35 sequencing primers made to the pCF3 insert was also used

to obtain 255 bp of sequence from pF1 that is shown as  
 SEQ ID NO:3. Nucleotides 68 to 255 of the genomic DNA  
 in pF1 (SEQ ID NO:3) are identical to nucleotides 1 to  
 188 of the cDNA (SEQ ID NO:1), which shows that they are  
 5 colinear and that the cDNA is encoded for by the gene in  
 the isolated genomic DNA. Nucleotides 113 to 115 in SEQ  
 ID NO:3 are the initiation codon of the largest open  
 reading frame corresponding to nucleotides 46-48 in SEQ  
 ID NO:1. This is evident from the presence of in-frame  
 10 termination codons at nucleotides 47 to 49 and  
 nucleotides 56 to 58 and the absence of observable  
 intron splice junctions in SEQ ID NO:3. The  
 identification of the 386 amino acid polypeptide as a  
 desaturase was confirmed by comparing its amino acid  
 15 sequence with all the protein sequences found in Release  
 19.0 of the SWISSPROTEIN database using the FASTA  
 algorithm of Pearson and Lipman (Proc. Natl. Acad. Sci.  
 USA (1988) 85:2444-2448) and the BLAST program (Altschul  
 et al., J. Mol. Biol. (1990) 215:403-410). The most  
 20 homologous protein found in both searches was the desA  
 fatty acid desaturase from the cyanobacterium  
Synechocystis PCC6803 (Wada, et al., Nature (1990)  
 347:200-203; Genbank ID:CSDESA; GenBank Accession  
 No:X53508). The 386 amino acid peptide in SEQ ID NO:1  
 25 was also compared to the 351 amino acid sequence of desA  
 by the method of Needleman et al. (J. Mol. Biol. (1970)  
 48:443-453). Over their entire length, these proteins  
 were 26% identical, the comparison imposing four major  
 gaps in the desA protein sequence. While this overall  
 30 homology is poor, homology in shorter stretches was  
 better. For instance, in a stretch of 78 amino acids  
 the Arabidopsis delta-15 desaturase (amino acids 78 to  
 155 in SEQ ID NO:1) and the desA protein (amino acids 67  
 to 144) showed 40% identity and 66% similarity.

Homology in yet shorter stretches was even greater as shown in Table 2.

TABLE 2

Peptide Length	AA positions in SEQ ID NO:1	AA positions in <i>desA</i>	Percent Identity
12	97-108	86-97	83
7	115-121	104-110	71
9	133-141	22-130	56
11	299-309	282-292	64

These high percent identities in short stretches of amino acids between the cyanobacterial desaturase polypeptide and SEQ ID NO:2 suggests significant relatedness between the two.

To analyse the developmental expression of the gene encoding mRNA corresponding to SEQ ID NO:1, the cDNA insert in plasmid pCF3 was used as a radiolabeled hybridization probe on mRNA samples from leaf, root, germinating seedling, and developing siliques from both wild type and mutant 3707 *Arabidopsis* plants, essentially as described in Maniatis et al., Molecular Cloning, A Laboratory Manual (1982) Cold Spring Harbor Laboratory Press. The results indicated that while the mRNA corresponding to SEQ ID NO:1 is detected in all tissues from the mutant plant, its levels are lower than in wild-type tissues. This is consistent with the observation that the fatty acid mutation in line 3707 is leaky relative to the known *Arabidopsis fad 3* mutant obtained via chemical mutagenesis. These results confirmed that the T-DNA in line 3707 had interrupted the normal expression of a fatty acid desaturase gene. Based on the fatty acid phenotype of homozygous mutant line 3707, Applicants concluded that the cDNA insert in pCF3 encoded the delta-15 desaturase. Further, Applicants concluded that it was the microsomal delta-15 desaturase, and not the chloroplastic delta-15

- desaturase, since: a) the mutant phenotype was expressed strongly in the seed but expressed poorly, if at all, in the leaf of line 3707, and b) the delta-15 desaturase polypeptide, by comparison to the desA polypeptide, did not have an N-terminal extension of a transit peptide expected for a nuclear-encoded chloroplast desaturase.

- The identity of SEQ ID NO:2 as the Arabidopsis microsomal delta-15 desaturase was confirmed by its biological overexpression in plant tissues. For this, the 1.4 kB Not I fragment of plasmid pCF3 containing the delta-15 desaturase cDNA was placed in the sense orientation behind either the CaMV 35S promotor, to provide constitutive expression, or behind the promotor for the gene encoding soybean  $\alpha'$  subunit of the  $\beta$ -conglycinin (7S) seed storage protein, to provide embryo-specific expression. The chimeric genes 35S promoter/sense SEQ ID NO:1/3' nopaline synthase and  $\beta$ -conglycinin/sense SEQ ID NO:1/3' phaseolin were then transformed into plant cells by Agrobacterium tumefaciens's binary Ti plasmid vector system [Hoekema et al. (1983) Nature 303:179-180; Bevan (1984) Nucl. Acids Res. 12:8711-8720].

- To confirm the identity of SEQ ID NO:1 and to test the biological effect of its overexpression in a heterologous plant species, the chimeric genes 35S promoter/sense SEQ ID NO:1/3' nopaline synthase was transformed into a binary vector, which was then transferred into Agrobacterium tumefaciens strain R1000, carrying the Ri plasmid pRiA4b from Agrobacterium rhizogenes [Moore et al. (1979) Plasmid 2:617-626]. Carrot (Daucus carota L.) cells were transformed by co-cultivation of carrot root disks with strain R1000 carrying the chimeric gene by the method of Petit et al. (1986) [Mol. Gen. Genet. 202:388-393]. Fatty acid

analyses of transgenic carrot "hairy" roots show that overexpression of Arabidopsis microsomal delta-15 desaturase can result in over 10-fold increase in 18:3 at the expense of 18:2.

- 5 To complement the delta-15 desaturation mutation in the T-DNA mutant line 3707 and to test the biological effect of overexpression of SEQ ID NO:1 in seed, the embryo-specific promoter/SEQ ID NO:1/3' phaseolin chimeric gene was transformed into a binary vector, 10 which was then transformed into the avirulent *Agrobacterium* strain LBA4404/pAL4404 [Hoekema et al. (1983) *Nature* 303:179-180]. Roots of line 3707 were transformed by the engineered *Agrobacterium*, transformed plants were selected and grown to give rise to seeds. 15 Fatty acid analysis of the seeds from two plants showed that the one out of six seeds in each plant showed the mutant fatty acid phenotype, while the remaining seeds show more than 10-fold increase in 18:3 to ca. 55%. While the sample size is small, this segregation 20 suggests Mendelian inheritance of the fatty acid phenotype. While most of the increase occurs at the expense of 18:2, some of it also occurs at the expense of 18:1. Thus, overexpression of this gene in oils crops, especially canola, which is a close relative of 25 Arabidopsis, is also expected to result in the high levels of 18:3 that are found in specialty oil of linseed.

- Comparisons of the sequence of the 386 amino acid polypeptide by the method of Needleman et al. (*J. Mol. Biol.* (1970) 48:443-453) with those for the microsomal 30 stearyl-CoA (delta-9) desaturases from rat, mouse and yeast revealed 21%, 19%, and 17% identities, respectively. While the membrane-associated Arabidopsis delta-15 desaturase protein showed significant but 35 limited homology to the desA protein, it showed no

significant homology to the soluble stearoyl-ACP (delta-9) desaturases from higher plants, including one from Arabidopsis.

Comparison of partial nucleotide sequences of  
5 plasmids pF1 and pS1 showed that the left T-DNA  
border:plant DNA junction is ca. 700 bp from the  
initiation codon in SEQ ID NO:1. To determine the  
position of the other T-DNA:plant DNA junction with  
respect to the pF1 sequence, the T-DNA:plant DNA  
10 junction fragment was isolated. Genomic DNA from mutant  
line 3707, isolated as described previously, was  
partially digested by restriction enzyme Mbo I to give  
an average fragment size of ca. 15 kB. The fragment  
ends were partially-filled with dGTP and gATP by Klenow  
15 and cloned into Xho I half-sites of LambdaGEM®-11  
(Promega Corporation) following the manufacturer's  
protocol. The phage library was titered and used  
essentially as described in Ausubel et al. [Current  
Protocols in Molecular Biology (1989) John Wiley &  
20 Sons]. The genomic phage library was screened with  
radiolabeled PCR product, ca. 0.6 kB, derived from 5'  
end of the gene in pF1. This product spans from 3 bp to  
the right of where the left-T-DNA border inserted to 15  
bp to the left of nucleotide position 1 in SEQ ID NO:1.  
25 Southern blot analysis of DNA from one of the purified,  
positively-hybridizing phages following Eco RI  
restriction digestion and electrophoresis showed that a  
4 kB Eco RI fragment hybridized to the 0.6 kB PCR  
product. The Eco RI fragment was subcloned and subject  
30 to sequence analyses. Comparison of the sequences  
derived from this fragment, pF1 and pS1 showed that the  
insertion of T-DNA resulted in a 56 bp deletion at the  
site of insertion and that the T-DNA interrupted the  
Arabidopsis gene 711 bp 5' to the initiation codon in SEQ  
35 ID NO:1. Thus, the T-DNA inserts 5' to the open reading

frame, consistent with the leaky expression of the gene encoding SEQ ID NO:1 and the leaky fatty acid phenotype in mutant 3707. While the left T-DNA:plant DNA junction is precise, that is without any sequence rearrangement in either the left T-DNA border or the flanking plant DNA, the other T-DNA:plant DNA junction is complex and not fully characterized.

Plasmid pCF3 was deposited on December 3, 1991 with the American Type Culture Collection of Rockville, Maryland under the provisions of the Budapest Treaty and bears accession number ATCC 68875.

Using Arabidopsis Delta-15 Desaturase cDNA as a  
Hybridization Probe to Isolate cDNAs Encoding  
Related Desaturases from Arabidopsis

The 1.4 kb Not I insert fragment isolated from plasmid pCF3 was purified, radiolabeled, and used to screen approximately 80,000 clones from the cDNA library made to poly A<sup>+</sup> mRNA from 3-day-old etiolated Arabidopsis thaliana as described above, except that lower stringency hybridizations (1 M NaCl, 50 mM Tris-HCl, pH 7.5, 1% SDS, 5% dextran sulfate, 0.1 mg/mL denatured salmon sperm DNA and 50°C) and washes (sequentially with 2X SSPE, 0.1% SDS at room temperature for 5 min and then again with fresh solution for 10 min, and finally with 0.5X SSPE, 0.1% SDS at 50°C for 5 min.) were used. Approximately 17 strongly-hybridizing and 17 weakly-hybridizing plaques were identified in the primary screen. Four of the weakly-hybridizing plaques were picked and subjected to one or two further rounds of screening with the radiolabeled probe as above until they were pure. To ensure that these were not delta-15 desaturase clones, they were further analyzed to determine whether they hybridized to an 18 bp oligomer specific to the 3' non-coding region of delta-15 desaturase cDNA (pCF3). After autoradiography of the



filters, one of the clones was found not to hybridize to this probe. This clone was picked, and a plasmid clone containing the cDNA insert was obtained as described above. Restriction analysis of this plasmid, designated

5 pCM2, showed that it had an approximately 1.3 kb cDNA insert which lacked a 0.7 kb Nco I - Bgl II fragment characteristic of the Arabidopsis delta-15 desaturase cDNA of pCF3. (This fragment corresponds to the DNA located between the Nco I site at nucleotides 474 to 479

10 and the Bgl II site at nucleotides 1164 to 1169 in SEQ ID NO:1). Partial nucleotide sequences of single strands from the 5' region and 3' region of pCM2 revealed that the cDNA insert was incomplete and that it encoded a polypeptide that is similar to, but distinct

15 from, that encoded by the cDNA in pCF3. In order to isolate a full-length version of the cDNA in plasmid pCM2, the 1.3 kb Not I fragment from plasmid pCM2 containing the cDNA insert was isolated and used as a radiolabeled hybridization probe to rescreen the same

20 Arabidopsis cDNA library as above. Three strongly hybridizing plaques were purified and the plasmids excised as described previously. The three resultant plasmids were digested by Not I restriction enzyme and shown to contain cDNA inserts ranging in size between 1

25 kB and 1.5 kB. Complete nucleotide sequence determination of the cDNA insert in one of these plasmids, designated pACF2-2, is shown in SEQ ID NO:4. SEQ ID NO:4 shows the 5' to 3' nucleotide sequence of base pairs of the Arabidopsis thaliana cDNA which

30 encodes a fatty acid desaturase. Nucleotides 10-12 and nucleotides 1358 to 1350 are, respectively, the putative initiation codon and the termination codon of the open reading frame (nucleotides 10 to 1350). The open reading frame was confirmed by comparison of its deduced

35 amino acid sequences with that of the related delta-15

fatty acid desaturase from soybean in this application. Nucleotides 1 to 9 and 1351 to 1525 are, respectively, the 5' and 3' untranslated nucleotides. The 446 amino acid protein sequence in SEQ ID NO:5 is that deduced from the open reading frame in SEQ ID NO:4 and has an estimated molecular weight of 51 kD. Alignment of SEQ ID NOS:2 and 5 shows an overall homology of approximately 80% and that the former has an approximately 55 amino acid long N-terminal extension, which is deduced to be a transit peptide found in nuclear-encoded plastid proteins.

To analyse the developmental expression of the gene corresponding to SEQ ID NO:4, this sequence was used as a radiolabeled hybridization probe on mRNA samples from leaf, root, germinating seedling, and developing siliques from both wild type and mutant line 3707 Arabidopsis plants, essentially as described in Maniatis et al. [Molecular Cloning, A Laboratory Manual (1982) Cold Spring Harbor Laboratory Press]. The results indicated that, in contrast to the constitutive expression of the gene encoding SEQ ID NO:1, the mRNA corresponding to SEQ ID NO:4 is abundant in green tissues, rare in roots and leaves, and is about three-fold more abundant in leaf than that of SEQ ID NO:1. The cDNA in plasmid pCM2 was also shown to hybridize polymorphically to genomic DNA from Arabidopsis thaliana (ecotype Wassileskija and marker line W100 ecotype Landesberg background) digested with Eco RI. It was used as a RFLP marker to map the genetic locus for the gene encoding this fatty acid desaturase in Arabidopsis. A single genetic locus was positioned corresponding to this desaturase cDNA. Its location was thus determined to be on chromosome 3 between the lambda AT228 and cosmid c3838 RFLP markers, "north" of the glabrous locus (Chang et al., Proc. Natl. Acad. Sci. USA (1988))

85:6856-6860; Nam et al., Plant Cell (1989) 1:699-705). This approximates the region to which Arabidopsis fatty acid desaturase fad\_2, fad\_D, and fad\_B mutations map [Somerville et al., (1992) in press]. Unsuccessful  
 5 efforts to clone the microsomal delta-12 fatty acid desaturase using cDNA inserts of pCF3 and pACF2-2 alongwith the above data led Applicants to conclude that the cDNA in pACF2-2 encodes a plastid delta-15 fatty acid desaturase that corresponds to the fad\_D locus.  
 10 This conclusion will be confirmed by biological expression of the cDNA in pACF2-2.

Plasmid pCM2 was deposited on November 27, 1991 with the American Type Culture Collection of Rockville, Maryland under the provisions of the Budapest Treaty and  
 15 bears accession number ATCC 68852.

The 1.4 kb, 1.3 kb, and 1.5 kb Not I cDNA insert fragments isolated from plasmids pCF3, pCM2 and pACF2-2 were purified, radiolabeled, and used several times to screen at low stringency as described above two  
 20 different cDNA libraries: one was made to poly A<sup>+</sup> mRNA from 3-day-old etiolated Arabidopsis thaliana ("etiolated" library) as described above and one made to polyA<sup>+</sup> mRNA from the above-ground parts of Arabidopsis thaliana plants, which varied in size from those that  
 25 had just opened their primary leaves to plants which had bolted and were flowering [Elledge et al. (1991) Proc. Natl. Acad Sci. USA 88:1731-1735]. The cDNA inserts in the library were made into an Xho I site flanked by Eco RI sites in lambda Yes vector [Elledge et al. (1991)  
 30 Proc. Natl. Acad Sci. USA 88:1731-1735] ("leaf" library). Several plaques from both libraries that hybridized weakly and in duplicate lifts to both SEQ ID NOS:1 and 4 were subjected to plaque purification. Phagemids were excised from the pure phages from  
 35 "etiolated" library as described above. Plasmids were

excised from the purified phages of the "leaf" library by site-specific recombination using the cre-lox recombination system in *E. coli* strain BNN132 [Elledge et al. (1991) Proc. Natl. Acad. Sci. USA 88:1731-1735].

- 5 In all cases, nucleotide sequencing of the cloned DNA revealed clones either identical to SEQ ID NOS:1 or 4 or unrecognizable sequences.

- In another set of experiments ca. 400,000 phages in the "leaf" library was screened with SEQ ID NOS:1 and 4  
 10 at low stringency (26 C, 1 M Na<sup>+</sup>, 50% formamide) and high stringency (42 C, 1 M Na<sup>+</sup>, 50% formamide). Of the several positive signals on the primary plaque lifts, 11 showed high stringency hybridization to SEQ ID NO:1, 35 showed high stringency hybridization to SEQ ID NO:4, and  
 15 39 hybridized to both at low stringency only. Twenty seven plaques of the low stringency signals came through a secondary low-stringency screen, 17 of which were used to make DNA from excised plasmids. Of the 7 plasmid DNA were sequenced, 8 were unrecognizable sequences, 5 were  
 20 identical to SEQ ID NO:1, 2 were identical to SEQ ID NO:2, and 2 were identical to one another and related but distinct to SEQ ID NOS:1 and 4. The novel desaturase sequence, designated pFad-x2, was also isolated from the "leaf" library independently by using  
 25 as a hybridization probe a 0.6 kB PCR product derived by polymerase chain reaction on poly A<sup>+</sup> RNA made from both canola seed as well as Arabidopsis leaves, as described elsewhere in this application, using degenerate oligomers made to conserved sequences between plant  
 30 delta-15 desaturases and the cyanobacterial *des A* desaturase. The PCR-derived plasmid, designated pYacp7, was sequenced partially from both ends. Comparison of the sequences of pFad-x2 and pYacp7 revealed that the two independently cloned cDNAs contained an identical  
 35 sequence that was related to the other delta-15

desaturases and that both were incomplete cDNAs. A partial composite sequence derived from both plasmids, pFadx-2 and pYacp7, is shown in SEQ ID NO:16 as a 5' to 3' nucleotide sequence of 472 bp. Nucleotides 2-4 and nucleotides 468 to 470 are, respectively, the first and the last codons in the open reading frame. This open reading frame is shown in SEQ ID NO:17. Comparison of SEQ ID NO:17 to the other delta-15 desaturase polypeptides disclosed in this application by the method of Needleman et al. [J. Mol. Biol. (1970) 48:443-453] using gap weight and gap length weight values of 3.0 and 0.1, respectively. The overall identities are between 65% and 68% between SEQ ID NO:17 and the microsomal delta-15 desaturases from Arabidopsis, canola and soybean and the overall identities are between 77% and 87% between SEQ ID NO:17 and the plastid delta-15 desaturases from Arabidopsis, canola and soybean. In addition SEQ ID NO:17 has an N-terminal peptide extension compared to the microsomal delta-15 desaturases that shows homology of the transit peptide sequence in Arabidopsis plastid delta-15 desaturase. On the basis of these comparisons it is deduced that SEQ ID NO:16 encodes a plastid delta-15 desaturase. There is genetic data in Arabidopsis suggesting the presence of two loci for plastid delta-15 desaturase. The full-length version of SEQ ID NO:16 can be readily isolated by one skilled in the art. The biological effect of introducing SEQ ID NO:16 or its full-length version into plants will be used to confirm its identity.

Plasmid pYacp7 was deposited on 20 November 1992 with the American Type Culture Collection of Rockville, Maryland under the provisions of the Budapest Treaty and bears accession number ATCC 69129.

Using Arabidopsis Delta-15 Desaturase cDNAs  
as Hybridization Probes to Isolate  
Delta-15 Desaturase cDNAs from Other Plant Species

For the purpose of cloning the Brassica napus seed  
5 cDNAs encoding delta-15 fatty acid desaturases, the cDNA  
inserts from pCF3 and pCM2 were isolated by polymerase  
chain reaction from the respective plasmids,  
radiolabeled, and used as hybridization probes to screen  
a lambda phage cDNA library made with poly A<sup>+</sup> mRNA from  
10 developing Brassica napus seeds 20-21 days after  
pollination. This cDNA library was screened several  
times at low stringency, using the Arabidopsis cDNA  
probes mentioned above. One of the Brassica napus  
cDNAs obtained in the initial screens was used as probe  
15 in a subsequent high stringency screen.

Arabidopsis pCM2 insert was radiolabeled and used  
as probe to screen approximately 300,000 plaques under  
low stringency hybridization conditions. The filter  
hybridizations were performed in 50 mM Tris pH 7.6, 6X  
20 SSC, 5X Denhardt's, 0.5% SDS, 100 ug denatured calf  
thymus DNA at 50°C overnight, and the posthybridization  
washes were carried out in 6X SSC, 0.5% SDS at room  
temperature for 15 min, then repeated with 2X SSC, 0.5%  
SDS at 45°C for 30 min, and then repeated twice with  
25 0.2X SSC, 0.5% SDS at 50°C for 30 min. Five strongly-  
hybridizing phages were obtained. These were plaque  
purified and used to excise the phagemids as described  
in the manual of the pBluescriptII Phagemid Kit from  
Stratagene (Stratagene 1991 catalogue, item 212205).  
30 One of these, designated pBNSF3-2, contained a 1.3 kb  
insert. pBNSF3-f2 was sequenced completely on both  
strands and the nucleotide sequence is shown in SEQ ID  
NO:6. Plasmid pBNSF3-2 was deposited on 27 November  
1991 with the American Type Culture Collection of

An additional low stringency screen using pCM2 probe provided eight strongly hybridizing phages. One of these, designated pBNSFd 8, contained a 0.4kb insert. pBNSFd-8 was sequenced completely on one strand, this nucleotide sequence showed significant divergence from the sequence SEQ ID NO:6 in the homologous region, which suggested that it corresponded to a novel Brassica napus seed desaturase different from that shown in SEQ ID NO:6. pBNSFd-8 insert was radiolabelled and used as hybridization probe in a high stringency screen of the Brassica napus seed cDNA library. The hybridization conditions were identical to those of the low stringency screen described above except for the temperature of the final two 30 min posthybridization washes in 0.2x SSC, 0.5% SDS was increased to 60°C. This screen resulted in three strongly hybridizing phages that were purified and excised. One of the excised plasmids pBNSFd-3 contained a 1.4kb insert that was sequenced completely on both strands. SEQ ID NO:8 shows the complete nucleotide sequence of pBNSFd-2.

A cDNA library was made to poly A<sup>+</sup> mRNA isolated from developing soybean seeds, and screened essentially as described above, except that filters were cross-hybridized in 25 mL of hybridization buffer consisting of 50mM Tris-HCl, pH 7.5, 1 M NaCl, 1% SDS, 0.5% dextran sulfate and 0.1 mg/mL denatured salmon sperm DNA (Sigma Chemical Co.) at 50°C for 2 h. Radiolabeled probes prepared from pCF3 as described above was added, and allowed to hybridize for 18 h at 50°C. The probes were washed twice at room temperature with 2X SSPE, 1%

SDS for five min followed by washing for 5 min at 50°C in 0.2X SSPE, 1% SDS. Autoradiography of the filters indicated that there was one strongly hybridizing plaque, and approximately five weakly hybridizing  
5 plaques. The more strongly hybridizing plaque was subjected to a second round of screening as before, except that the final wash was for 5 min at 60°C in 0.2X SSPE, 1% SDS. Numerous, strongly hybridizing plaques were observed, and one, well-isolated from other phage,  
10 was picked for further analysis.

Sequences of the pBluescript vector from the purified phage, including the cDNA insert, were excised in the presence of a helper phage and the resultant phagemid was used to infect *E. coli* XL-1 Blue cells.  
15 DNA from the plasmid, designated pXF1, was made by the alkaline lysis miniprep procedure described in Sambrook et al. (Molecular Cloning, A Laboratory Manual, 2nd ed. (1989) Cold Spring Harbor Laboratory Press). The alkali-denatured double-stranded DNA from pXF1 was  
20 completely sequenced on both strands. The insert of pXF1 contained a stretch of 1783 nucleotides which contained an unknown open-reading frame and also contained a poly-A stretch of 16 nucleotides 3' to the open reading frame, from nucleotides 1767 to 1783,  
25 followed by an Eco RI restriction site. The 2184 bases that followed this Eco RI site contained a 1145 bp open reading frame which encoded a polypeptide of about 68% identity to, and colinear with, the *Arabidopsis* delta-15 desaturase polypeptide listed in SEQ ID No:2. The  
30 putative start methionine of the 1145 bp open-reading frame corresponded to the start methionine of the *Arabidopsis* microsomal delta-15 peptide and there were no amino acids corresponding to a plastid transit peptide 5' to this methionine. When the insert in pXF1  
35 was digested with Eco RI four fragments were observed,



fragments of approximately 370 bp and 1400 bp fragments, derived from the first 1783 bp of the insert in pXF1, and fragments of approximately 600 bp and 1600 bp derived from the the other 2184 nucleotides of the

5 insert in pXF1. Only the 600 bp and 1600 bp fragments hybridized with probe derived from pCF3 on Southern blots. It was deduced that pXF1 contained two different cDNA inserts separated by an Eco RI site and the second of these inserts was a 2184 bp cDNA encoding a soybean

10 microsomal delta-15 desaturase. The complete nucleotide sequence of the 2184 bp soybean microsomal delta-15 cDNA contained in plasmid pXF1 is listed in SEQ ID No:10. Plasmid pXF1 was deposited on December 3, 1991 with the American Type Culture Collection of Rockville, Maryland

15 under the provisions of the Budapest Treaty and bears accession number ATCC 68874.

Using Soybean Microsomal Delta-15 Desaturase cDNA as a Hybridization Probe to Isolate

cDNAs Encoding Related Desaturases from Soybean

20 A 1.0 kb fragment of DNA corresponding to part of the coding region of the soybean microsomal delta-15 desaturase cDNA contained in plasmid pXF1, was excised with the restriction enzyme Hha I and gel purified. The fragment was labeled with  $^{32}\text{P}$  as described above and

25 used to probe a soybean cDNA library as described above. Autoradiography of the filters indicated that there were eight hybridizing plaques and these were subjected to a second round of screening. Sequences of the pBluescript vector from all eight of the purified phages, including

30 the cDNA inserts, were excised in the presence of a helper phage and the resultant phagemids were used to infect *E. coli* XL-1 Blue cells. DNA from the plasmids was made by the alkaline lysis miniprep procedure described in Sambrook et al. (Molecular Cloning, A

35 Laboratory Manual, 2nd ed. (1989) Cold Spring Harbor

Laboratory Press). Restriction analysis showed they contained inserts ranging from 1.0 kb to 3.0 kb in size. One of these inserts, designated pSFD-118bwp, contained an insert of about 1700 bp. The alkali-denatured double-stranded DNA from pSFD-118bwp was completely sequenced on both strands, shown in SEQ ID NO:12. The insert of pSFD-118bwp contained a stretch of 1675 nucleotides which contained an open-reading frame encoding a polypeptide, shown in SEQ ID NO:13, of about 80% identity with, and colinear with, the Arabidopsis plastid delta-15 desaturase polypeptide listed in SEQ ID No:5. The open-reading frame also encoded amino acids corresponding to a plastid transit peptide at the 5' end of the open-reading frame. The transit peptide was colinear with, and shared some homology to, the transit peptide described for the Arabidopsis plastid delta-15 glycerolipid desaturase. The complete nucleotide sequence of the 1675 bp soybean plastid delta-15 glycerolipid desaturase cDNA is listed in SEQ ID No:12.

Comparison of the different delta-15 desaturase sequences disclosed in the application by the method of Needleman et al. (J. Mol. Biol. (1970) 48:443-453) using gap weight and gap length weight values of 3.0 and 0.1, respectively, reveals the relatedness between them as shown in Table 3.

TABLE 3  
Percent Identities Between Different Delta-15  
Fatty Acid Desaturases at the Amino Acid Level

	<u>aD</u>	<u>c3</u>	<u>cD</u>	<u>s3</u>	<u>sD</u>
a3	66	93	66	68	67
aD	-	67	90	67	69
c3	-	-	68	68	68
cD	-	-	-	68	74

- a3, aD, c3, cD, s3 and sD refer, respectively, to  
 SEQ ID NO:2 (Arabidopsis microsomal delta-15  
 desaturase), SEQ ID NO:5 (Arabidopsis plastid delta-15  
 desaturase), SEQ ID NO:7 (canola microsomal delta-15  
 desaturase), SEQ ID NO:9 (canola plastid delta-15  
 desaturase), SEQ ID NO:11 (soybean microsomal delta-15  
 desaturase), and SEQ ID NO:13 (soybean plastid delta-15  
 desaturase). Based on these comparisons, the delta-15  
 desaturases, of both microsomal and plastid types, have  
 overall identities of 65% or more at the amino acid  
 levels, even when from different plant species.

Isolation of Nucleotide Sequences Encoding  
Homologous and Heterologous Glycerolipid Desaturases

- Fragments of the instant invention may be used to  
 isolate cDNAs and genes of homologous and heterologous  
 glycerolipid desaturases from the same species as the  
 fragment of the invention or from different species.  
 Isolation of homologous genes using sequence-dependent  
 protocols is well-known in the art. Southern blot  
 analysis revealed that the Arabidopsis microsomal  
 delta-15 desaturase cDNA (SEQ ID NO:1) hybridized to  
 genomic DNA fragments of corn and soybean. In addition,  
 Applicants have demonstrated that it can be used to  
 isolate cDNAs encoding seed microsomal delta-15  
 desaturases from Brassica napus (SEQ ID NO:6) and  
 soybean (SEQ ID NO:10). Thus, one can isolate cDNAs and

genes for homologous glycerolipid desaturases from the same or different higher plant species, especially from the oil-producing species.

More importantly, one can use the fragments of the invention to isolate cDNAs and genes for heterologous glycerolipid desaturases, including those found in plastids. Thus, *Arabidopsis* microsomal delta-15 desaturase cDNA (SEQ ID NO:1) was successfully used as a hybridization probe to isolate cDNAs encoding the related plastid delta-15 desaturases from *Arabidopsis* (SEQ ID NO:4) and *Brassica napus* (SEQ ID NO: 8), and the soybean microsomal delta-15 soybean (SEQ ID NO:10) was successfully used to isolate soybean cDNA encoding plastid delta-15 desaturase (SEQ ID NO:12).

In a particular embodiment of the present invention, regions of the nucleic acid fragments of the invention that are conserved between different desaturases may be used by one skilled in the art to design a mixture of degenerate oligomers for use in sequence-dependent protocols aimed at isolating nucleic acid fragments encoding other homologous or heterologous glycerolipid desaturase cDNA's or genes. For example, by comparing all desaturase polypeptides one can identify stretches of amino acids that are conserved between them, and then use the conserved amino acid sequence to design oligomers, both short degenerate or long ones, or "guessmers" as known by one skilled in the art (see Sambrook et al., (Molecular Cloning, A Laboratory Manual, 2nd ed. (1989), Cold Spring Harbor Laboratory Press). Such oligomers and "guessmers" may be used as hybridization probes as known to one skilled in the art.

For example, comparison of cyanobacterial desA and plant delta-15 desaturases revealed a particularly well conserved stretch of amino acids (amino acids 97-108 in

SEQ ID NO:1). SEQ ID NOS:20 and 21 represent two sets of 36-mers each 16-fold degenerate made to this region. End-labeled oligomers represented in SEQ ID NOS:20 and 21 were mixed and used as hybridization probes to screen

5 Arabidopsis cDNA libraries. Most of the positively-hybridizing plaques also hybridized to cDNAs encoding Arabidopsis microsomal and plastid delta-15 desaturases (SEQ ID NOS:1 and 4). However, the use of SEQ ID NOS:20 and 21 did not give consistent and reproducible results.

10 A 135 base-long oligomer (SEQ ID NO:32) was also made as an antisense strand to a longer stretch of the same conserved region, amino acids 97 to 141 in SEQ ID NO:1 (FVLGHDCGHSFSDIPLLNSVVGHILHSFILVPYHGWRISHRTHH). At positions of ambiguity, the design used either

15 deoxyinosines or most frequently used codons based on the codon usage in Arabidopsis genes. When used as a hybridization probe, the 135-mer hybridized to all plaques that also hybridized to cDNAs encoding Arabidopsis microsomal and plastid delta-15 desaturases

20 (SEQ ID NOS:1 and 4). In addition, it also hybridized to plaques that did not hybridize to SEQ ID NOS:1 and 4). The latter were purified and excised as described previously. Nucleotide sequencing of the cDNA inserts in the resultant plasmids revealed DNA sequences that

25 did not show any relatedness to any desaturase.

For another example, in the polymerase chain reaction (Innis, et al., Eds, (1990) PCR Protocols: A Guide to Methods and Applications, Academic Press, San Diego), two short pieces of the present fragment of the

30 invention can be used to amplify a longer glycerolipid desaturase DNA fragment from DNA or RNA. The polymerase chain reaction may also be performed on a library of cloned nucleotide sequences with one primer based on the fragment of the invention and the other on either the

35 poly A<sup>+</sup> tail or a vector sequence. These oligomers may



SEQ ID NO	Length	Fold Degeneracy	AA positions in SEQ ID NO:2	AA Sequence
20	36	16	97-108 (S)	FVLGHDCGHGSF
21	36	16	97-108 (S)	FVLGHDCGHGSF
28	36	16	97-108 (S)	FVLGHDCGHGSF
29	36	16	97-108 (S)	FVLGHDCGHGSF
22	18	72	100-105 (S)	GHDCGH
23	18	72	100-105 (S)	GHDCGH
24	18	72	299-304 (AS)	HDIGTH
25	18	72	299-304 (AS)	HDIGTH
26	23	416	304-309 (AS)	HVIHHL
27	23	416	304-309 (AS)	HVIHHL
30	38	64	299-311 (AS)	HDIGTHVIHHLFP
31	38	64	299-311 (AS)	HDIGTHVIHHLFP

- In one experiment, PCRs were performed using SEQ ID NOS:22 and 23 as sense primers and either SEQ ID NOS:24 and 25 or SEQ ID NOS:26 and 27 as antisense primers on poly A+ RNA purified from both Arabidopsis leaf and canola developing seeds. All PCRs resulted in PCR products of the correct size (ca. 630 bp). The PCR products from Arabidopsis and canola were purified and used as radiolabeled hybridization probes to screen the Lambda Yes Arabidopsis cDNA library, as described above.
- 5 This led to the isolation of a pure phage, which was excised to give plasmid pYacp7. The cDNA insert in pYacp7 was partially sequenced. It's sequence showed that it encoded an incomplete desaturase polypeptide that was identical to another cDNA (in plasmid pFadx-2)
- 10 isolated by low-stringency hybridization as described previously. The composite sequence derived from the partial sequences from the cDNA inserts in pFadx-2 and pYacp7 is shown in SEQ ID NO:16 and the polypeptide encoded by it in SEQ ID NO:17. As discussed previously,
- 15 SEQ ID NO:17 is a putative plastid delta-15 desaturase. This is further supported by Southern blot analysis
- 20

using radiolabeled cDNA inserts from either pCF3, pACF2-2, or pYacp7 on Arabidopsis genomic DNA digested with one of several enzymes. It shows that the different inserts hybridize to different restriction fragments and that only the inserts from pACF2-2 and pYacp7 show some cross-hybridization.

In another PCR experiment, PCR was performed using ca. 80 pmoles each of SEQ ID NOS:28 and 29 as sense primers and ca. 94 pmoles each of SEQ ID NOS:30 and 31 as antisense primers on poly A+ RNA purified from Arabidopsis mutant line 3707. This was performed using GeneAmp® RNA PCR Kit (Perkin Elmer Cetus) following manufacturer's protocol and using the following program:

a) 1 cycle of 2 min at 95°C, b) 35 cycles of 1 min at 95°C (denaturation), 1 min at 50°C (annealing) and 1 min at 65°C (extension), and c) 1 cycle of 7 min at 65°C. The resulting PCR product, of the correct size (ca. 630 bp), was purified, radiolabeled, and used as a hybridization probe on a Southern blot of Arabidopsis genomic DNA as described above. While it hybridized to restriction fragments that also hybridized to SEQ ID NOS:1 (Arabidopsis microsomal delta-15 desaturase), 4 (Arabidopsis plastid delta-15 desaturase), and 16 (Arabidopsis plastid delta-15 desaturase), it also hybridized to novel fragments that did not hybridize to previously cloned desaturase cDNAs. However, even after several attempts, the radiolabeled PCR product did not hybridize to any novel cDNA clone when used as a probe on different Arabidopsis cDNA libraries: in all cases it hybridized only to plaques that also hybridized to the known desaturase cDNAs. Furthermore, the PCR product was subcloned into a plasmid vector and after screening about a 100 of these, none gave rise to a clone with a novel desaturase sequence.



The isolation of other glycerolipid desaturases will become easier as more examples of glycerolipid desaturases are isolated using the fragments of the invention. Knowing the conserved amino acid sequences from diverse desaturases will also allow one to identify more and better consensus sequences. Such sequences can be used to make hybridization probes or amplification primers which will further aid in the isolation of different glycerolipid desaturases, including those from non-plant sources such as fungi, algae, and even cyanobacteria, as well as other membrane-associated desaturases from other organisms.

The function of the diverse nucleotide fragments encoding glycerolipid desaturases that can be isolated using the present invention can be identified by transforming plants with the isolated desaturase sequences, linked in sense or antisense orientation to suitable regulatory sequences required for plant expression, and observing the fatty acid phenotype of the resulting transgenic plants. Preferred target plants for the transformation are the same as the source of the isolated nucleotide fragments when the goal is to obtain inhibition of the corresponding endogenous gene by antisense inhibition or cosuppression. Preferred target plants for use in expression or overexpression of the isolated nucleic acid fragments are plants with known mutations in desaturation reactions, such as the Arabidopsis desaturase mutants, mutant flax deficient in delta-15 desaturation, or mutant sunflower deficient in delta-12 desaturation. Alternatively, the function of the isolated nucleic acid fragments can be determined similarly via transformation of other organisms, such as yeast or cyanobacteria, with chimeric genes containing the nucleic acid fragment and suitable regulatory

sequences followed by analysis of fatty acid composition and/or enzyme activity.

Overexpression of the Glycerolipid  
Desaturase Enzymes in Transgenic Species

5       The nucleic acid fragment(s) of the instant  
invention encoding functional glycerolipid  
desaturase(s), with suitable regulatory sequences, can  
be used to overexpress the enzyme(s) in transgenic  
organisms. Such recombinant DNA constructs may include  
10 either the native glycerolipid desaturase gene or a  
chimeric glycerolipid desaturase gene isolated from the  
same or a different species as the host organism. For  
overexpression of glycerolipid desaturase(s), it is  
preferable that the introduced gene be from a different  
15 species to reduce the likelihood of cosuppression. For  
example, overexpression of delta-15 desaturase in  
soybean, rapeseed, or other oil-producing species to  
produce altered levels of polyunsaturated fatty acids  
may be achieved by expressing RNA from the entire cDNA  
20 found in pCF3. Similarly, the isolated nucleic acid  
fragments encoding glycerolipid desaturases from  
Arabidopsis, rapeseed, and soybean can also be used by  
one skilled in the art to obtain substantially  
homologous full-length cDNAs, if not already obtained,  
25 as well as the corresponding genes as fragments of the  
invention. These, in turn, may be used to overexpress  
the corresponding desaturases in plants. One skilled in  
the art can also isolate the coding sequence(s) from the  
fragment(s) of the invention by using and/or creating  
30 sites for restriction endonucleases, as described in  
Sambrook et al., (Molecular Cloning, A Laboratory  
Manual, 2nd ed. (1989), Cold Spring Harbor Laboratory  
Press). For example, the fragment in SEQ ID NO:1 in  
plasmid pCF3 is flanked by Not I sites and can be  
35 isolated as a Not I fragment that can be introduced in

the sense orientation relative to suitable plant regulatory sequences. Alternatively, sites for Nco I (5'-CCATGG-3') or Sph I (5'-GCATGC-3') that allow precise removal of coding sequences starting with the initiating codon "ATG" may be engineered into the fragment(s) of the invention. For example, for utilizing the coding sequence of delta-15 desaturase from pCF3, an Sph I site can be engineered by substituting nucleotides at positions 44, 45, and 49 of SEQ ID NO:1 with G, C, and C, respectively.

#### Inhibition of Plant Target Genes by Use of Antisense RNA

Antisense RNA has been used to inhibit plant target genes in a tissue-specific manner (see van der Krol et al., *Biotechniques* (1988) 6:958-976). Antisense inhibition has been shown using the entire cDNA sequence (Sheehy et al., *Proc. Natl. Acad. Sci. USA* (1988) 85:8805-8809) as well as a partial cDNA sequence (Cannon et al., *Plant Molec. Biol.* (1990) 15:39-47). There is also evidence that the 3' non-coding sequences (Ch'ng et al., *Proc. Natl. Acad. Sci. USA* (1989) 86:10006-10010) and fragments of 5' coding sequence, containing as few as 41 base-pairs of a 1.87 kb cDNA (Cannon et al., *Plant Molec. Biol.* (1990) 15:39-47), can play important roles in antisense inhibition.

The use of antisense inhibition of the glycerolipid desaturases may require isolation of the transcribed sequence for one or more target glycerolipid desaturase genes that are expressed in the target tissue of the target plant. The genes that are most highly expressed are the best targets for antisense inhibition. These genes may be identified by determining their levels of transcription by techniques, such as quantitative analysis of mRNA levels or nuclear run-off transcription, known to one skilled in the art.

For example, antisense inhibition of delta-15 desaturase in Brassica napus resulting in altered levels of polyunsaturated fatty acids may be achieved by expressing antisense RNA from the entire or partial cDNA  
 5 found in pBNSF3-2.

#### Inhibition of Plant Target Genes by Cosuppression

The phenomenon of cosuppression has also been used to inhibit plant target genes in a tissue-specific  
 10 manner. Cosuppression of an endogenous gene using the entire cDNA sequence (Napoli et al., The Plant Cell (1990) 2:279-289; van der Krol et al., The Plant Cell (1990) 2:291-299) as well as a partial cDNA sequence (730 bp of a 1770 bp cDNA) (Smith et al., Mol. Gen.  
 15 Genetics (1990) 224:477-481) are known.

The nucleic acid fragments of the instant invention encoding glycerolipid desaturases, or parts thereof, with suitable regulatory sequences, can be used to reduce the level of glycerolipid desaturases, thereby  
 20 altering fatty acid composition, in transgenic plants which contain an endogenous gene substantially homologous to the introduced nucleic acid fragment. The experimental procedures necessary for this are similar to those described above for the overexpression of the  
 25 glycerolipid desaturase nucleic acid fragments. For example, cosuppression of delta-15 desaturase in Brassica napus resulting in altered levels of polyunsaturated fatty acids may be achieved by expressing in the sense orientation the entire or  
 30 partial seed delta-15 desaturase cDNA found in pBNSF3-2.

#### Selection of Hosts, Promoters and Enhancers

A preferred class of heterologous hosts for the expression of the nucleic acid fragments of the invention are eukaryotic hosts, particularly the cells  
 35 of higher plants. Particularly preferred among the

higher plants are the oil-producing species, such as soybean (Glycine max), rapeseed (including Brassica napus, B. campestris), sunflower (Helianthus annuus), cotton (Gossypium hirsutum), corn (Zea mays), cocoa  
 5 (Theobroma cacao), safflower (Carthamus tinctorius), oil palm (Elaeis guineensis), coconut palm (Cocos nucifera), flax (Linum usitatissimum), and peanut (Arachis hypogaea).

Expression in plants will use regulatory sequences  
 10 functional in such plants. The expression of foreign genes in plants is well-established (De Blaere et al., Meth. Enzymol. (1987) 153:277-291). The source of the promoter chosen to drive the expression of the fragments  
 15 of the invention is not critical provided it has sufficient transcriptional activity to accomplish the invention by increasing or decreasing, respectively, the level of translatable mRNA for the glycerolipid desaturases in the desired host tissue. Preferred  
 20 promoters include (a) strong constitutive plant transcripts in cauliflower mosaic virus (Odell et al., Nature (1985) 313:810-812; Hull et al., Virology (1987) 86:482-493), and (b) tissue- or developmentally-specific promoters. Examples of tissue-specific promoters are  
 25 the light-inducible promoter of the small subunit of ribulose 1,5-bis-phosphate carboxylase (if expression is desired in photosynthetic tissues), the maize zein protein promoter (Matzke et al., EMBO J. (1984) 3:1525-1532), and the chlorophyll a/B binding protein  
 30 promoter (Lampa et al., Nature (1986) 316:750-752).

Particularly preferred promoters are those that allow seed-specific expression. This may be especially useful since seeds are the primary source of vegetable oils and also since seed-specific expression will avoid  
 35 any potential deleterious effect in non-seed tissues.

Examples of seed-specific promoters include, but are not limited to, the promoters of seed storage proteins, which can represent up to 90% of total seed protein in many plants. The seed storage proteins are strictly regulated, being expressed almost exclusively in seeds in a highly tissue-specific and stage-specific manner (Higgins et al., *Ann. Rev. Plant Physiol.* (1984) 35:191-221; Goldberg et al., *Cell* (1989) 56:149-160). Moreover, different seed storage proteins may be expressed at different stages of seed development.

Expression of seed-specific genes has been studied in great detail (See reviews by Goldberg et al., *Cell* (1989) 56:149-160 and Higgins et al., *Ann. Rev. Plant Physiol.* (1984) 35:191-221). There are currently numerous examples of seed-specific expression of seed storage protein genes in transgenic dicotyledonous plants. These include genes from dicotyledonous plants for bean b-phaseolin (Sengupta-Gopalan et al., *Proc. Natl. Acad. Sci. USA* (1985) 82:3320-3324; Hoffman et al., *Plant Mol. Biol.* (1988) 11:717-729), bean lectin (Voelker et al., *EMBO J.* (1987) 6:3571-3577), soybean lectin (Okamuro et al., *Proc. Natl. Acad. Sci. USA* (1986) 83:8240-8244), soybean Kunitz trypsin inhibitor (Perez-Grau et al., *Plant Cell* (1989) 1:095-1109), soybean b-conglycinin (Beachy et al., *EMBO J.* (1985) 4:3047-3053; pea vicilin (Higgins et al., *Plant Mol. Biol.* (1988) 11:683-695), pea convicilin (Newbigin et al., *Planta* (1990) 180:461-470), pea legumin (Shirsat et al., *Mol. Gen. Genetics* (1989) 215:326-331); rapeseed napin (Radke et al., *Theor. Appl. Genet.* (1988) 75:685-694) as well as genes from monocotyledonous plants such as for maize 15 kD zein (Hoffman et al., *EMBO J.* (1987) 6:3213-3221), maize 18 kD oleosin (Lee et al., *Proc. Natl. Acad. Sci. USA* (1991) 888:6181-6185), barley b-hordein (Marris et al., *Plant Mol. Biol.* (1988)

- 10:359-366) and wheat glutenin (Colot et al., EMBO J. (1987) 6:3559-3564). Moreover, promoters of seed-specific genes operably linked to heterologous coding sequences in chimeric gene constructs also maintain
- 5 their temporal and spatial expression pattern in transgenic plants. Such examples include use of Arabidopsis thaliana 2S seed storage protein gene promoter to express enkephalin peptides in Arabidopsis and B. napus seeds (Vandekerckhove et al.,
- 10 Bio/Technology (1989) 7:929-932), bean lectin and bean b-phaseolin promoters to express luciferase (Riggs et al., Plant Sci. (1989) 63:47-57), and wheat glutenin promoters to express chloramphenicol acetyl transferase (Colot et al., EMBO J. (1987) 6:3559-3564).
- 15 Of particular use in the expression of the nucleic acid fragment of the invention will be the heterologous promoters from several soybean seed storage protein genes such as those for the Kunitz trypsin inhibitor (Jofuku et al., Plant Cell (1989) 1:1079-1093; glycinin
- 20 (Nielson et al., Plant Cell (1989) 1:313-328), and b-conglycinin (Harada et al., Plant Cell (1989) 1:415-425). Promoters of genes for a- and b-subunits of soybean  $\beta$ -conglycinin storage protein will be particularly useful in expressing the mRNA or the
- 25 antisense RNA in the cotyledons at mid- to late-stages of seed development (Beachy et al., EMBO J. (1985) 4:3047-3053) in transgenic plants. This is because there is very little position effect on their expression in transgenic seeds, and the two promoters show
- 30 different temporal regulation. The promoter for the a-subunit gene is expressed a few days before that for the b-subunit gene. This is important for transforming rapeseed where oil biosynthesis begins about a week before seed storage protein synthesis (Murphy et al., J. Plant Physiol. (1989) 135:63-69).
- 35

Also of particular use will be promoters of genes expressed during early embryogenesis and oil biosynthesis. The native regulatory sequences, including the native promoters, of the glycerolipid desaturase genes expressing the nucleic acid fragments of the invention can be used following their isolation by those skilled in the art. Heterologous promoters from other genes involved in seed oil biosynthesis, such as those for *B. napus* isocitrate lyase and malate synthase (Comai et al., Plant Cell (1989) 1:293-300), delta-9 desaturase from safflower (Thompson et al. Proc. Natl. Acad. Sci. USA (1991) 88:2578-2582) and castor (Shanklin et al., Proc. Natl. Acad. Sci. USA (1991) 88:2510-2514), acyl carrier protein (ACP) from *Arabidopsis* (Post-Beittenmiller et al., Nucl. Acids Res. (1989) 17:1777), *B. napus* (Safford et al., Eur. J. Biochem. (1988) 174:287-295), and *B. campestris* (Rose et al., Nucl. Acids Res. (1987) 15:7197), b-ketoacyl-ACP synthetase from barley (Siggaard-Andersen et al., Proc. Natl. Acad. Sci. USA (1991) 88:4114-4118), and oleosin from *Zea mays* (Lee et al., Proc. Natl. Acad. Sci. USA (1991) 88:6181-6185), soybean (Genbank Accession No: X60773) and *B. napus* (Lee et al., Plant Physiol. (1991) 96:1395-1397) will be of use. If the sequence of the corresponding genes is not disclosed or their promoter region is not identified, one skilled in the art can use the published sequence to isolate the corresponding gene and a fragment thereof containing the promoter. The partial protein sequences for the relatively-abundant enoyl-ACP reductase and acetyl-CoA carboxylase are also published (Slabas et al.; Biochim. Biophys. Acta (1987) 877:271-280; Cottingham et al., Biochim. Biophys. Acta (1988) 954:201-207) and one skilled in the art can use these sequences to isolate the corresponding seed genes with their promoters. Similarly, the fragments of the

00210005-122290



present invention encoding glycerolipid desaturases can be used to obtain promoter regions of the corresponding genes for use in expressing chimeric genes.

Attaining the proper level of expression of the  
5 nucleic acid fragments of the invention may require the use of different chimeric genes utilizing different promoters. Such chimeric genes can be transferred into host plants either together in a single expression vector or sequentially using more than one vector.

10 It is envisioned that the introduction of enhancers or enhancer-like elements into the promoter regions of either the native or chimeric nucleic acid fragments of the invention will result in increased expression to accomplish the invention. This would include viral  
15 enhancers such as that found in the 35S promoter (Odell et al., Plant Mol. Biol. (1988) 10:263-272), enhancers from the opine genes (Fromm et al., Plant Cell (1989) 1:977-984), or enhancers from any other source that result in increased transcription when placed into a  
20 promoter operably linked to the nucleic acid fragment of the invention.

Of particular importance is the DNA sequence element isolated from the gene for the a-subunit of b-conglycinin that can confer 40-fold seed-specific  
25 enhancement to a constitutive promoter (Chen et al., Dev. Genet. (1989) 10:112-122). One skilled in the art can readily isolate this element and insert it within the promoter region of any gene in order to obtain seed-specific enhanced expression with the promoter in  
30 transgenic plants. Insertion of such an element in any seed-specific gene that is expressed at different times than the b-conglycinin gene will result in expression in transgenic plants for a longer period during seed development.

The invention can also be accomplished by a variety of other methods to obtain the desired end. In one form, the invention is based on modifying plants to produce increased levels of glycerolipid desaturases by virtue of introducing more than one copy of the foreign gene containing the nucleic acid fragments of the invention. In some cases, the desired level of polyunsaturated fatty acids may require introduction of foreign genes for more than one kind of glycerolipid desaturase.

Any 3' non-coding region capable of providing a polyadenylation signal and other regulatory sequences that may be required for the proper expression of the nucleic acid fragments of the invention can be used to accomplish the invention. This would include 3' ends of the native glycerolipid desaturase(s), viral genes such as from the 35S or the 19S cauliflower mosaic virus transcripts, from the opine synthesis genes, ribulose 1,5-bisphosphate carboxylase, or chlorophyll a/b binding protein. There are numerous examples in the art that teach the usefulness of different 3' non-coding regions.

#### Transformation Methods

Various methods of transforming cells of higher plants according to the present invention are available to those skilled in the art (see EPO Pub. 0 295 959 A2 and 0 318 341 A1). Such methods include those based on transformation vectors utilizing the Ti and Ri plasmids of Agrobacterium spp. It is particularly preferred to use the binary type of these vectors. Ti-derived vectors transform a wide variety of higher plants, including monocotyledonous and dicotyledonous plants (Sukhapinda et al., Plant Mol. Biol. (1987) 8:209-216; Potrykus, Mol. Gen. Genet. (1985) 199:183). Other transformation methods are available to those skilled in the art, such as direct uptake of foreign DNA constructs

- (see EPO Pub. 0 295 959 A2), techniques of electroporation (Fromm et al., Nature (1986) (London) 319:791) or high-velocity ballistic bombardment with metal particles coated with the nucleic acid constructs
- 5 (Kline et al., Nature (1987) (London) 327:70). Once transformed, the cells can be regenerated by those skilled in the art.

- Of particular relevance are the recently described methods to transform foreign genes into commercially
- 10 important crops, such as rapeseed (De Block et al., Plant Physiol. (1989) 91:694-701), sunflower (Everett et al., Bio/Technology (1987) 5:1201), and soybean (Christou et al., Proc. Natl. Acad. Sci USA (1989) 86:7500-7504).

15 Application to RFLP Technology

- The use of restriction fragment length polymorphism (RFLP) markers in plant breeding has been well-documented in the art (Tanksley et al., Bio/Technology (1989) 7:257-264). The nucleic acid fragments of the
- 20 invention can be used as RFLP markers for traits linked to expression of glycerolipid desaturases. These traits will include altered levels of unsaturated fatty acids. The nucleic acid fragment of the invention can also be used to isolate the glycerolipid desaturase gene from
- 25 variant (including mutant) plants with altered levels of unsaturated fatty acids. Sequencing of these genes will reveal nucleotide differences from the normal gene that cause the variation. Short oligonucleotides designed around these differences may be used as hybridization
- 30 probes to follow the variation in polyunsaturates. Oligonucleotides based on differences that are linked to the variation may be used as molecular markers in breeding these variant oil traits.

EXAMPLES

The present invention is further defined in the following Examples, in which all parts and percentages are by weight and degrees are Celsius, unless otherwise stated. It should be understood that these Examples, while indicating preferred embodiments of the invention, are given by way of illustration only. From the above discussion and these Examples, one skilled in the art can ascertain the essential characteristics of this invention, and without departing from the spirit and scope thereof, can make various changes and modifications of the invention to adapt it to various usages and conditions. All publications, including patents and non-patent literature, referred to in this specification are expressly incorporated by reference herein.

EXAMPLE 1

ISOLATION OF GENOMIC DNA FLANKING THE T-DNA SITE OF  
INSERTION IN ARABIDOPSIS THALIANA MUTANT LINE 3707  
20 Identification of an Arabidopsis thaliana T-DNA Mutant  
with Low Linolenic Acid Content

A population of Arabidopsis thaliana (geographic race Wassilewskija) transformants containing the T-DNA of Agrobacterium tumefaciens was generated by seed  
25 transformation as described by Feldmann et al., (Mol. Gen. Genetics (1987) 208:1-9). In this population the transformants contain DNA sequences encoding the pBR322 bacterial vector, nopaline synthase, neomycin phosphotransferase (NPTII, confers kanamycin  
30 resistance), and b-lactamase (confers ampicillin resistance) within the T-DNA border sequences. The integration of the T-DNA into different areas of the chromosomes of individual transformants may cause a disruption of plant gene function at or near the site of  
35 insertion, and phenotypes associated with this loss of

gene function can be analyzed by screening the population for the phenotype.

- T3 seed was generated from the wild type seed treated with Agrobacterium tumefaciens by two rounds of self-fertilization as described by Feldmann et al., (Science (1989) 243:1351-1354). These progeny were segregating for the T-DNA insertion, and thus for any mutation resulting from the insertion. Approximately 100 seeds of each of 6000 lines were combined and the fatty acid content of each of the 6000 pooled samples was determined by gas chromatography of the fatty acyl methyl esters essentially as described by Browse et al., (Anal. Biochem. (1986) 152:141-145) except that 2.5% H<sub>2</sub>SO<sub>4</sub> in methanol was used as the methylation reagent and samples were heated for 1.5 h at 80°C to effect the methanolysis of the seed triglycerides. A line designated "3707" produced seeds that gave an altered fatty acid profile compared to that of the total population. T3 plants were grown from individual T3 seeds produced by line 3707 and self-fertilized to produce T4 seeds on individual plants that were either homozygous wild type, homozygous mutant, or heterozygous for the mutation. The percent fatty acid compositions of a representative subsample of the entire population, of the pooled 3707 T3 seeds, and of a homozygous T4 mutant segregant are shown in Table 4.

TABLE 4

<u>Fatty Acid Methyl Ester</u>	<u>T3 Pools from lines 3501-4000 average and (std. deviation)</u>	<u>3707 T3 Pool</u>	<u>3707 Homozygous T4 Segregant</u>
palmitic	7.4 (0.37)	7.0	6.4
stearic	3.0 (0.22)	2.9	3.0
oleic	17.0 (1.5)	17.7	15.9
linoleic	29.3 (0.78)	35.0	42.4
linolenic	16.1 (1.1)	10.2	3.1
eicosenoic	20.2 (0.73)	20.5	23.6

- The phenotype of the segregating T3 pool of line 3707 (high linoleic acid, low linolenic acid) was intermediate between that of the population subsample and the homozygous T4 mutant seeds suggesting that line 3707 harbored a mutation at a locus which controls the conversion of linoleic to linolenic acid in the seed. Still, it was not apparent whether the mutant phenotype in line 3707 was the result of a T-DNA insertion.
- Therefore, Applicants checked a segregating T4 population to determine whether the mutant fatty acid phenotype cosegregated with the nopaline synthase activity and kanamycin resistance encoded by the T-DNA insert. A total of 263 T4 plants were grown and assayed for the presence of nopaline in leaf extracts (Errampalli et al., The Plant Cell (1991) 3:149-157). In addition, T5 seeds were collected from each of the T4 plants and samples of 10-50 seeds were taken to determine the seed fatty acid composition and to determine their ability to germinate in the presence of kanamycin (Feldmann, et al., (1989) Science 243:1351-1354). The 263 plants fell into 3 classes as in Table 5.

00210035-102306

TABLE 5

<u>Number of Individuals</u>	<u>Phenotype</u>
63	T4 plants: little or no nopaline present; T5 seeds: wild type fatty acid composition, all kanamycin sensitive
134	T4 plants: nopaline present; T5 seeds: heterozygous fatty acid composition similar to 3707 T3 pool, segregating for kanamycin resistance
64	T4 plants: nopaline present; T5 seeds homozygous mutant fatty acid composition, all kanamycin resistant

The cosegregation of the fatty acid phenotype with the phenotypes conferred by T-DNA sequences in an approximately 1:2:1 pattern provided strong evidence

- 5 that the mutation in line 3707 was the result of a T-DNA insertion. Further experiments were then conducted with the intent of using probes containing T-DNA sequences to clone the T-DNA insert and flanking genomic DNA from line 3707.

10 Preparation of Genomic DNA from Homozygous 3707 Plants

- Seeds from a homozygous line derived from Arabidopsis thaliana (geographic race Wassilewskija (WS)) line 3707 were surface sterilized for 5 min at room temperature in a solution of 5.25% sodium hypochlorite (w/v)/0.15% Tween 20 (v/v), then washed 15 several times in sterile distilled water, with a final rinse in 50% ethanol. Immediately following the ethanol wash, the seeds were transferred to sterile filter paper to dry. One to three seeds were then transferred to 20 250-mL flasks containing 50 mL of sterile Gamborgs B5 media (Gibco, 500-1153EA), pH 6.0. Cultures were incubated at 22°C, 70  $\mu\text{E}\cdot\text{m}^{-2}\cdot\text{sec}^{-1}$  of continuous light for approximately three weeks, after which time the root tissue was harvested, made into 10 g aliquots (wet 25 weight), lyophilized, and stored at -20°C.

- Using a variation of the procedure of Shure et al., (Cell (1983) 35:225-233) genomic DNA was isolated from the root tissue. Two aliquots of lyophilized tissue were ground to a fine powder using a mortar and pestle.
- 5 The ground tissue was added to a flask containing 85 mL of lysis buffer (7 M urea, 0.35 M NaCl, 0.05 M Tris-HCl, pH 8.0, 0.02 M EDTA, 1% Sarkosyl, 5% phenol) and mixed gently with a glass rod to obtain a homogeneous suspension. To this suspension an equal volume of
- 10 phenol:chloroform:isoamyl alcohol (25:24:1) (equilibrated with 10 mM Tris, pH 8, 1 mM EDTA) was added. After the addition of 8.5 mL of 10% SDS the mixture was swirled on a rotating platform for 15 min at room temperature. After centrifugation at 2000xg for 15
- 15 min, the upper aqueous phase was removed to a new tube and extracted two more times, as above, but without the addition of SDS. To the final aqueous phase was added 1/20th the volume of 3 M potassium acetate, pH 5.5 and two times the volume of ice cold 100% ethanol.
- 20 Precipitation of the DNA was facilitated by incubation at -20°C for one hour followed by centrifugation at 12,000xg for 10 min. The resulting pellet was resuspended in 3 mL of 10 mM Tris, pH 8, 1 mM EDTA to which was added 0.95 g of cesium chloride (CsCl) and
- 25 21.4 µL of 10 mg/mL ethidium bromide (EtBr) per mL of solution. The DNA was then purified by centrifugation to equilibrium in a CsCl/EtBr density gradient for 16 h at 15°C, 265,000xg. After removal from the gradient, the DNA was extracted with isopropanol saturated with TE
- 30 buffer (10 mM Tris, pH 8; 1 mM EDTA) and CsCl to remove EtBr and then dialyzed overnight at 4°C against 10 mM Tris, pH 8, 1 mM EDTA to remove CsCl. The DNA was removed from dialysis and the concentration was determined using the Hoechst fluorometric assay in which
- 35 an aliquot of DNA is added to 3 mL of  $1.5 \times 10^{-6}$  M bis-



benzimidazole (Hoechst 33258, Sigma) in 1X SSC (0.15 M NaCl, 0.015 M sodium citrate), pH 7.0, incubated at room temperature for 5 min, and read on a fluorometer at excitation 360, emission 450, against a known set of DNA standards.

#### Plasmid Rescue and Analysis

Five micrograms of genomic DNA from the homozygous 3707 mutant, prepared as described above, was digested with 20 units of either Bam HI or Sal I restriction enzyme (Bethesda Research Laboratory) in a 50  $\mu$ L reaction volume according to the manufacturer's specifications. After digestion the DNA was extracted with buffer-saturated phenol (Bethesda Research Laboratory) followed by precipitation in ethanol. The resulting pellet was resuspended in a final volume of 10  $\mu$ L of 10 mM Tris, pH 8, and the concentration of the DNA was determined using the Hoechst fluorometric assay as above.

To facilitate circularization, as opposed to end-to-end joining, a dilute ligation reaction was set up containing 250 ng of Bam HI or Sal I digested genomic DNA, 3 Weiss units of T4 DNA ligase (Promega), 50  $\mu$ L of 10X ligase buffer (30 mM Tris-HCl, pH 7.8, 100 mM MgCl<sub>2</sub>, 100 mM DTT, 5 mM ATP) and 5  $\mu$ L of 100 mM ATP in a 500  $\mu$ L reaction volume. The reaction was incubated for 16 h at 16°C, heated for 10 min at 70°C, and extracted once with buffer saturated phenol (Bethesda Research Laboratory). The DNA was then precipitated with the addition of two volumes of 100% ethanol and 1/10th volume of 7.5 M ammonium acetate. The resulting pellet was resuspended in a final volume of 10  $\mu$ L of 10 mM Tris, pH 8, and the concentration of the DNA was determined using the Hoechst fluorometric assay as above.

Competent DH10B cells (Bethesda Research Laboratory) were transfected with 50 ng of ligated DNA

at a concentration of 10 ng of DNA per 100  $\mu$ L of cells according to the manufacturer's specifications. Transformants from Sal I or Bam HI digests were selected on LB plates (10 g Bacto-tryptone, 5 g Bacto-yeast extract, 5 g NaCl, 15 g agar per liter, pH 7.4) containing 100  $\mu$ g/mL ampicillin or 25  $\mu$ g/mL kanamycin sulfate, respectively. Ampicillin-resistant ( $Amp^r$ ; ampicillin sensitivity,  $Amp^s$ ) Sal I transformants were screened for the presence of the kanamycin resistance ( $Kan^r$ ; kanamycin sensitivity,  $Kan^s$ ) gene by picking primary transformants and stabbing them first to LB plates containing 100  $\mu$ g/mL ampicillin then to LB plates containing 25  $\mu$ g/mL kanamycin. After overnight incubation at 37°C the plates were scored for  $Amp^r/Kan^s$  colonies. Kanamycin-resistant Bam HI transformants were screened for the presence of the ampicillin resistance gene by picking primary transformants and stabbing them first to LB plates containing 25  $\mu$ g/mL kanamycin and then to LB plates containing 100  $\mu$ g/mL ampicillin. After overnight incubation at 37°C the plates were scored for  $Kan^r/Amp^r$  colonies.

Cultures were made of 192  $Amp^r/Kan^s$  Sal I transformants and 85  $Kan^r/Amp^r$  Bam HI transformants directly into deep-well microtiter plates containing 200  $\mu$ L of LB broth (10 g Bacto-tryptone, 5 g Bacto-yeast extract, 5 g NaCl per liter) with 100  $\mu$ g/mL ampicillin. Using the Schleicher and Schuell Minifold I apparatus and Nytran membranes, dot blots were set up, in duplicate, using the following conditions: 50  $\mu$ L of culture was diluted into 150  $\mu$ L of 5X SSC, the culture was lysed and the DNA denatured by the addition of 150  $\mu$ L of 0.5 M NaOH, 1.5 M NaCl solution for 3 min at room temperature, the filter was removed from the apparatus and neutralized in 0.5 M Tris, pH 8, 1.5 M NaCl, the DNA was then UV cross-linked to the filters

using the Stratagene Stratalinker, and the filters were heated for 2 h at 80°C and stored at room temperature.

- To determine whether T-DNA was contained within any of the rescued plasmids, the dot blots were probed with portions of the right and left borders of T-DNA. The right border probe consisted of a 2.2 kb Hind III-Dra I fragment of DNA obtained from plasmid H23pKC7 (composed of the 3.2 kb Hind III 23 fragment from Ti plasmid pTiC58 (Lemmers et al., J. Mol. Biol. (1989) 144;353-376) cloned into plasmid vector pKC7 (Maniatis et al., Molecular Cloning, A Laboratory Manual (1982) Cold Spring Harbor Laboratory Press)), and the left border probe consisted of a 2.9 kb Hind III-Eco RI fragment obtained from plasmid H10pKC7 (composed of the 6.5 kb Hind III 10 fragment from Ti plasmid pTiC58 (Lemmers et al., J. Mol. Biol. (1989) 144;353-376) cloned into plasmid vector pKC7 (Maniatis et al., Molecular Cloning, A Laboratory Manual (1982) Cold Spring Harbor Laboratory Press)) using standard digestion, electrophoresis, and electroelution conditions as described in Sambrook et al., (Molecular Cloning, A Laboratory Manual, 2nd ed (1989) Cold Spring Harbor Laboratory Press). Final DNA purification was obtained by passage of the eluted DNA over an Elutip-D column (Schleicher and Schuell) using the manufacturer's specifications. Concentration of the DNA was determined using the Hoechst fluorometric assay as above. Approximately 100 ng of each probe was labeled with a[<sup>32</sup>P]dCTP using a Random Priming Kit from Bethesda Research Laboratories under conditions recommended by the manufacturer. Labeled probe was separated from unincorporated a[<sup>32</sup>P]dCTP by passing the reaction through a Sephadex G-25 spun column under standard conditions as described in Sambrook et al., (Molecular

Cloning, A Laboratory Manual, 2nd ed. (1989) Cold Spring Harbor Laboratory Press).

The filters were pre-hybridized in 150 mL of buffer consisting of 6X SSC, 10X Denhardt's solution, 1% SDS, and 100 µg/mL denatured calf thymus DNA for 16 h at 42°C. The denatured, purified, labeled probe was added to the pre-hybridized filters following transfer of the filters to 50 mL of hybridization buffer consisting of 6X SSC, 1% SDS, 10% dextran sulfate, and 50 µg/mL denatured calf thymus DNA. Following incubation of the filters in the presence of the probe for 16 h at 65°C, the filters were washed twice in 150 mL of 6X SSC, 0.5% SDS, twice in 1X SSC, 1% SDS and once in 0.1X SSC, 1% SDS, all at 65°C. The washed filters were subjected to autoradiography on Kodak XAR-2 film at 80°C overnight.

Of the 85 Bam HI candidates, 63 hybridized with the left border probe and none hybridized with the right border probe. Of the 192 Sal I candidates, 31 hybridized with the left border probe, 4 hybridized with the right border probe, and none hybridized with both probes. Twelve of the Bam HI candidates, 7 positive and 5 negative for the presence of the left border of T-DNA, were further analyzed by restriction digests.

DNA from the Bam HI candidates was made by the alkaline lysis miniprep procedure of Birnboim et al., (Nuc. Acid Res. (1979) 7:1513-1523), as described in Sambrook et al., (Molecular Cloning, A Laboratory Manual, 2nd ed. (1989), Cold Spring Harbor Laboratory Press). The plasmid DNA was digested with Eco RI restriction enzyme (Bethesda Research Laboratories) in accordance with the manufacturer's specifications and electrophoresed through a 0.8% agarose gel in 1X TBE buffer (0.089 M Tris-borate, 0.089 M boric acid, 0.002 M EDTA). All of the Bam HI candidates which hybridized with the left border probe of T-DNA had the same Eco RI

restriction pattern, which indicated the presence of 14.2 kb of T-DNA and 1.4 kb of putative plant genomic DNA in these clones.

- DNA from Sal I candidates was isolated,
- 5 restriction-analyzed using Eco RI, Bam HI and Sal I enzymes, and electrophoresed through a 0.8% agarose gel, as above. All of the Sal I candidates which hybridized with the left border probe of T-DNA included 2.9 kb of putative plant DNA. Contained within this 2.9 kb
- 10 fragment was a 1.4 kb Bam HI-Eco RI fragment as seen with the Bam HI rescued plasmids, suggesting that the 1.4 kb fragment was a subset of the 2.9 kb fragment and that it was adjacent to the left border of the T-DNA at its site of insertion into the plant genome. Sequence
- 15 analysis of one Sal I candidate (pS1) using a primer homologous to the left border sequence of T-DNA, revealed that the sequence of pS1 was colinear with the sequence of the T-DNA left border (Yadav et al., Proc. Natl. Acad. Sci. USA (1982) 79:6322-6326) up to
- 20 nucleotide 65, followed by non-T-DNA (putative plant) sequences.

#### Southern Analysis with Putative Plant

##### DNA from Rescued Plasmids

- DNA from the seven Bam HI candidates which
- 25 hybridized with the left border of the T-DNA was pooled and a portion was digested with Eco RI and Bam HI restriction endonucleases and electrophoretically separated on a 0.8% agarose gel in 1X TBE buffer. After excising a 1.4 kb Eco RI-Bam HI fragment from the
- 30 agarose gel, the 1.4 kb fragment was purified by use of a Gene Clean Kit from Bio 101. Fifty nanograms of the resulting DNA fragment was labeled with a [ $^{32}$ P]dCTP using a Random Priming Kit (Bethesda Research Laboratory) under conditions recommended by the manufacturer.

- Three micrograms of total genomic DNA from homozygous wild-type Arabidopsis and homozygous 3707 mutant Arabidopsis plants was digested to completion with one of the following restriction enzymes: Sal I, Hind III, Eco RI, Cla I, and Bam HI under conditions suggested by the manufacturer. The digested DNA was subjected to electrophoresis and Southern transfer to Hybond-N membranes (Amersham) as described in Sambrook et al. (Molecular Cloning, A Laboratory Approach, 2nd. ed. (1989) Cold Spring Harbor Laboratory Press). Southern transfer, the membranes were exposed to UV light using the Stratalinker (Stratagene) as per the manufacturer's instructions, air dried, and heated at 68°C for 2 h.
- The filters were prehybridized in 1 M NaCl, 50 mM Tris-Cl, pH 7.5, 1% sodium dodecyl sulfate, 5% dextran sulfate, 100 µg/mL of denatured salmon sperm DNA at 65°C overnight. Fifty nanograms of the radiolabeled 1.4 kb Eco RI-Bam HI plant DNA fragment prepared above was added to the prehybridization solution containing the Southern blot and further incubated at 65°C overnight. The filter was washed for 10 min twice in 200 mL 2X SSPE, 0.1% sodium dodecyl sulfate at 65°C and for 10 min in 200 mL 0.5% SSPE, 0.1% sodium dodecyl sulfate at 65°C. Hybridizing fragments were detected by autoradiography. The analysis confirmed that the probe fragment contained plant DNA and that the T-DNA integration site was in a 2.8 kb Bam HI, a 5.2 kb Hind III, a 3.5 kb Sal I, a 5.5 kb Eco RI, and an approximately 9 kb Cla I fragment of wild type Arabidopsis DNA.

Isolation of Lambda Clones Containing the Wild Type  
Arabidopsis Delta-15 Desaturase Gene

- The 1.4 kb Eco RI-Bam HI fragment (see above) was used as a probe to screen a lGem-11 library made from

genomic DNA isolated from wildtype *Arabidopsis thaliana* plants, geographic race WS. To construct the library, genomic DNA was partially digested with Sau3A enzyme, and size-fractionated over a salt gradient as described in Sambrook et al. (Molecular Cloning, A Laboratory Approach, 2nd ed. (1989) Cold Spring Harbor Laboratory Press). The size-fractionated DNA was then cloned into Bam HI-digested lGem-11 phage DNA (Promega) following the protocol outlined by the manufacturer. About 25,000 plaque-forming units of phage each were plated on five 150 mm petri plates containing a lawn of KW251 cells on NZY agar media (5 g NaCl, 2 g  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ , 5 g yeast extract, 10 g NZ Amine (casein hydrolysate from ICN Pharmaceuticals), 15 g agar per liter; pH 7.5). The plaques were adsorbed onto nylon membranes (Colony/Plaque Screen, New England Nuclear), in duplicate, and prepared according to the manufacturer's instructions with the addition of a 2 h incubation at 80°C after air drying the filters. The filters were prehybridized at 65°C in hybridization buffer (1% BSA, 0.5 M  $\text{NaPi}$ , pH 7.2, ( $\text{NaH}_2\text{PO}_4$  and  $\text{Na}_2\text{HPO}_4$ ), 10 mM EDTA, and 7% SDS) for 4 h, after which time they were transferred to fresh buffer containing the denatured radiolabeled probe (see above) and incubated overnight at 65°C. The filters were rinsed twice with 0.1X SSC, 1% SDS at 65°C for 30 min each and subjected to autoradiography on Kodak XA-R film at 80°C overnight. Seven positively-hybridizing plaques were subjected to plaque purification as described in Sambrook et al., (Molecular Cloning, A Laboratory Manual, 2nd ed. (1989), Cold Spring Harbor Laboratory Press).

Small scale (5 mL) liquid lysates from each of the 7 clones were prepared and titered on KW251 bacteria as described in Sambrook et al. (Molecular Cloning, A Laboratory Manual, 2nd ed (1989), Cold Spring Harbor

Laboratory Press). Phage DNA was isolated using a variation of the method of Chisholm (Biotechniques (1989) 7:21-23) in which the initial lysate was made according to Sambrook et al. (Molecular Cloning, A Laboratory Manual, 2nd ed (1989), Cold Spring Harbor Laboratory Press) the concentration of DNase I and RNase I (Sigma) was reduced by half, and the PEG precipitation step was increased to 16 h. Based on restriction analysis using Hind III, Sal I and Xho I enzymes, the original 7 positive phage fell into 5 different classes. While the average insert size was approximately 15 kb, taken together the clones spanned a 40 kb region of genomic DNA. Through restriction mapping using 4 different enzymes (Hind III, Bam HI, Kpn I, and Sal I) singly, and in pair-wise combinations, accompanied by Southern analysis with the 1.4 kb Eco RI-Bam HI probe (as above) and other probes obtained from the 1 clones themselves, a partial map was obtained in which all 5 clones (11111, 141A1, 14211, 14311 and 14411) were found to share an approximately 3 kb region of homology near the site of T-DNA insertion. Via restriction and Southern analysis, Applicants ascertained that a 5.2 kb Hind III fragment present in clones 1111, 41A1, and 4411 also spanned the site of the T-DNA insertion. This fragment was excised from lambda clone 41A1, inserted into the Hind III site of the pBluescript vector (Stratagene), and the resulting plasmid, designated pF1, was prepared and isolated using standard protocols. This Hind III fragment was subsequently used to probe an Arabidopsis cDNA library (see below).



EXAMPLE 2

CLONING OF ARABIDOPSIS THALIANA DELTA-15  
DESATURASE cDNA USING GENOMIC DNA FLANKING  
THE T-DNA SITE OF INSERTION IN ARABIDOPSIS THALIANA  
5 MUTANT LINE 3707 AS A HYBRIDIZATION PROBE

The 5.2 kb Hind III fragment from plasmid pF1 was purified by electrophoresis in agarose after digestion of the plasmid with Hind III and radiolabeled with <sup>32</sup>P as described above. For the preparation of an

10 Arabidopsis cDNA library, polyadenylated mRNA was prepared from 3 day-old, etiolated Arabidopsis (ecotype Columbia) seedling hypocotyls using standard protocols (Sambrook, et al., Molecular Cloning: A Laboratory Manual, 2nd Ed. (1989) Cold Spring Harbor Laboratory

15 Press). Five micrograms of this mRNA were used as template with an oligo d(T) primer, and Moloney Murine Leukemia Virus reverse transcriptase (Pharmacia) was used to catalyze first strand cDNA synthesis. Second-strand cDNA was made according to Gubler et al., (Gene

20 (1983) 25:263-272) except that DNA ligase was omitted. After the second strand synthesis, the ends of the cDNA were made blunt by reaction with the Klenow fragment of DNA polymerase and ligated to Eco RI/Not I adaptors (Pharmacia). The cDNA's were purified by spun-column

25 chromatography using Sephacryl S-300 and size-fractionated on a 1% low melting point agarose gel. Size-selected cDNAs (1-3 kb) were removed from the gel using agarase (New England Biolabs) and purified by phenol:chloroform extraction and ethanol precipitation.

30 One hundred nanograms of the cDNA was co-precipitated with 1 µg of 1 ZAP II (Stratagene) Eco RI-digested, dephosphorylated arms. The DNAs were ligated in a volume of 4 µL overnight, and the ligation mix was packaged in vitro using the Gigapack II Gold packaging

35 extract (Stratagene).

Approximately 80,000 phage were screened for positively hybridizing plaques using the radiolabeled 5.2 kb Hind III fragment as a probe essentially as described above and in Sambrook et al., (Molecular Cloning: A Laboratory Manual, 2nd ed. (1989) Cold Spring Harbor Laboratory Press). Replica filters of the phage plaques were soaked in 1 M NaCl, 50 mM Tris-HCl, pH 7.5, 1% SDS, 5% dextran sulfate, 0.1 mg/mL denatured salmon sperm DNA during the pre-hybridization step (8 hr at 65°C) and then probe was added and the hybridization proceeded over 16 hr at the same temperature. Filters were washed sequentially with 2X SSPE, 0.1% SDS at room temperature for 5 min and then again with fresh solution for 10 min, and finally with 0.5X SSPE, 0.1% SDS at 65°C for 5 min. Approximately 20 positively hybridizing plaques were identified in the primary screen. Four of these were picked and subjected to two further rounds of screening and purification. From the tertiary screen, four pure phage plaques were isolated. Plasmid clones containing the cDNA inserts were obtained through the use of a helper phage according to the in vivo excision protocol provided by Stratagene. Double-stranded DNA was prepared using the alkaline lysis method as previously described, and the resulting plasmids were size-analyzed by electrophoresis in agarose gels. The largest one of these, designated pCF3, contained an approximately 1.4 kb insert which was sequenced using Sequenase T7 DNA polymerase (US Biochemical Corp.) and the manufacturer's instructions, beginning with primers homologous to vector sequences that flank the cDNA insert and continuing serially with primers designed from the newly acquired sequences as the sequencing experiment progressed. The sequence of this insert is shown in SEQ ID NO:1.

00240035.122000

EXAMPLE 3CLONING OF AN ARABIDOPSIS CDNA ENCODING A PLASTID  
DELTA-15 FATTY ACID DESATURASE

- A related fatty acid desaturase was cloned in a similar fashion, except that the probe used was not derived from a PCR reaction on pCF3, but rather was the actual 1.4 kb Not I fragment isolated from pCF3 which was purified and radiolabeled as described above.
- Approximately 80,000 phage from the Arabidopsis etiolated hypocotyl cDNA library described above were plated out and screened essentially as before, except as indicated below. The filters were soaked in 1 M NaCl, 50 mM Tris-HCl, pH 7.5, 1% SDS, 5% dextran sulfate, 0.1 mg/mL denatured salmon sperm DNA during the pre-hybridization step (8 hr at 50°C). Then probe was added and the hybridization proceeded over 16 hr at the same temperature. Filters were washed sequentially with 2X SSPE, 0.1% SDS at room temperature for 5 min and then again with fresh solution for 10 min, and finally with 0.5X SSPE, 0.1% SDS at 50°C for 5 min. Approximately 17 strongly hybridizing and 17 weakly hybridizing plaques were identified in the primary screen. Four of the weakly hybridizing plaques were picked and subjected to one to two further rounds of screening with the radiolabeled probe as above until they were pure. To ensure that these were not delta-15 desaturase clones, they were further analyzed to determine whether they hybridized to a delta-15 desaturase 3' end-specific probe. The probe used was an 18 bp oligonucleotide which is complementary in sequence (i.e., antisense) to nucleotides 1229 - 1246 of SEQ ID NO:1. The probe was radiolabeled with gamma-<sup>32</sup>P ATP using T4 polynucleotide kinase and hybridized to filters containing DNA from the isolated clones in 6X SSC, 5X Denhardt's, 0.1 mg/mL denatured salmon sperm DNA, 1 mM EDTA, 1% SDS at 44°C

09219935.123308

overnight. The filters were washed twice in 6X SSC, 0.1% SDS for 5 min at room temperature, then in 6X SSC, 0.1% SDS at 44°C for 3-5 min. After autoradiography of the filters, one of the clones failed to show

5 hybridization to this probe. This clone was picked, and a plasmid clone containing the cDNA insert was obtained through the use of a helper phage according to the in vivo excision protocol provided by Stratagene. Double-stranded DNA was prepared using the alkaline lysis

10 method as previously described, and the resulting plasmid was size-analyzed by electrophoresis in agarose gels following either Not I digestion or digestion with both Nco I and Bgl II. The results were consistent with the presence in this plasmid, designated pCM2, of an

15 approximately 1.3 kb cDNA insert which lacked a 0.7 kb Nco I - Bgl II fragment characteristic of the Arabidopsis delta-15 desaturase cDNA of pCF3. (This fragment corresponds to the DNA located between the Nco I site at nucleotides 474-479 and the Bgl II site at

20 nucleotides 1164-1169 in SEQ ID NO:1). The complete nucleotide sequence of pCM2 is shown in SEQ ID NO:4.

#### EXAMPLE 4

#### CLONING OF PLANT FATTY ACID DESATURASE cDNAs FROM OTHER SPECIES BY HYBRIDIZATION TECHNIQUES

25 An approximately 1.4 kb fragment containing the Arabidopsis delta-15 desaturase coding sequence of SEQ ID NO:1 was obtained from plasmid pCF3 through the use of the polymerase chain reaction (PCR). Primers (M13(-20) and T7-17mer primers, 1991 Stratagene

30 Catalogue numbers 300303 and 300302, respectively) flanking the pCF3 insert were used in the PCR which was carried out essentially as described in the instructions provided by the vendor in the Perkin-Elmer/Cetus PCR kit. This fragment was digested with Not I to remove

35 vector sequences, purified by agarose gel electro-

phoresis, and radiolabeled with  $^{32}\text{P}$  as previously described.

#### EXAMPLE 5

#### CLONING OF BRASSICA NAPUS SEED cDNAs ENCODING 5 DELTA-15 FATTY ACID DESATURASES

- A cDNA library from developing Brassica napus seeds was constructed using the polyadenylated mRNA fraction contained in a polysomal RNA preparation from developing Brassica napus seeds. Polysomal RNA was isolated following the procedure of Kamalay et al., (Cell (1980) 19:935-946) from seeds 20-21 days after pollination. The polyadenylated mRNA fraction was obtained by affinity chromatography on oligo-dT cellulose (Aviv et al., Proc. Natl. Acad. Sci. USA (1972) 69:1408-1411).
- 15 Four micrograms of polyadenylated mRNA were reverse transcribed and used to construct a cDNA library in lambda phage (Uni-ZAP<sup>TM</sup> XR vector) using the protocol described in the ZAP-cDNA<sup>TM</sup> Synthesis Kit (1991 Stratagene Catalog, Item # 200400).
- 20 For the purpose of cloning the Brassica napus seed cDNAs encoding delta-15 fatty acid desaturases, the Brassica napus seed cDNA library was screened several times using the inserts from the Arabidopsis cDNAs pCF3 and pCM2 as radiolabelled hybridization probes. One of
- 25 the Brassica napus cDNAs obtained in these screens was used as hybridization probe in a subsequent screen.
- For each screening experiment approximately 300,000 phages were screened under low stringency hybridization conditions. The filter hybridizations were carried out
- 30 in 50 mM Tris pH 7.6, 6X SSC, 5X Denhardt's, 0.5% SDS, 100 ug denatured calf thymus DNA at 50°C overnight and the p[ost hybridization washes were performed in 6X SSC, 0.5% SDS at room temperature for 15 min, then repeated with 2X SSC, 0.5% SDS at 45°C for 30 min, and then

repeated twice with 0.2X SSC, 0.5% SDS at 50°C for 30 min.

Using the Arabidopsis cDNA insert of pCM2 as a probe in a low stringency screen five strongly hybridizing phages were identified. These phages were purified and excised according to the protocols described in the ZAP-cDNA™ Synthesis Kit and pBluescript II Phagemid Kit (1991 Stratagene Catalog, Item # 200400 and 212205). One of these, designated pBNSF3-f2, contained a 1.3 kb insert. pBNSF3-f2 insert was sequenced completely on both strands. pBNSF3-f2 nucleotide sequence is shown in SEQ ID NO:6. A comparison of this sequence with that of the Arabidopsis thaliana delta-15 desaturase clone (SEQ ID NO:1) confirmed that pBNSF3-f2 is a Brassica napus cDNA that encodes a seed microsomal delta-15 desaturase.

An additional low stringency screen of the Brassica napus seed cDNA library using the cDNA insert in pCM2 as a probe identified eight strongly-hybridizing phages. These phages were plaque purified and used to excise the phagemids as described above. One of these, designated pBNSFd-8, contained a 0.3kb insert. pBNSFd-8 was sequenced completely on one strand, this sequence had significant divergence from the sequence of pBNSF3-f2. The cDNA insert in pBNSFd-8 was used as a hybridization probe in a high stringency screen of the Brassica napus seed cDNA library. The filter hybridizations were carried out in 50 mM Tris pH 7.6, 6X SSC, 5X Denhardt's, 0.5% SDS, 100 ug denatured calf thymus DNA overnight at 50°C and post hybridization washes were in 6X SSC, 0.5% SDS at room temperature for 15 min, then with 2X SSC, 0.5% SDS at 45°C for 30 min, and then twice with 0.2X SSC, 0.5% SDS at 60°C for 30 min. The high stringency screen resulted in three strongly hybridizing phages that were purified and excised as above. One of the

5 delta-15 desaturase clone (SEQ ID NO:4) confirmed that  
pBNSFd-3 is a Brassica napus cDNA that encodes a seed  
plastid delta-15 desaturase.

A cDNA library was made as follows: Soybean embryos (ca. 50 mg fresh weight each) were removed from the pods and frozen in liquid nitrogen. The frozen embryos were ground to a fine powder in the presence of liquid nitrogen and then extracted by Polytron homogenization and fractionated to enrich for total RNA by the method of Chirgwin et al. (Biochemistry (1979) 18:5294-5299). The nucleic acid fraction was enriched for poly A<sup>+</sup>RNA by passing total RNA through an oligo-dT cellulose column and eluting the poly A<sup>+</sup>RNA with salt as described by Goodman et al. (Meth. Enzymol. (1979) 68:75-90). cDNA was synthesized from the purified poly A<sup>+</sup>RNA using cDNA Synthesis System (Bethesda Research Laboratory) and the manufacturer's instructions. The resultant double-stranded DNA was methylated by Eco RI DNA methylase (Promega) prior to filling-in its ends with T4 DNA polymerase (Bethesda Research Laboratory) and blunt-end ligation to phosphorylated Eco RI linkers using T4 DNA ligase (Pharmacia). The double-stranded DNA was digested with Eco RI enzyme, separated from excess linkers by passage through a gel filtration column (Sephacrose CL-4B), and ligated to lambda ZAP vector (Stratagene) according to manufacturer's instructions. Ligated DNA was packaged into phage using the Gigapack packaging extract (Stratagene) according to manufacturer's instructions. The resultant cDNA library

10 A cDNA library was made as follows: Soybean embryos (ca. 50 mg fresh weight each) were removed from the pods and frozen in liquid nitrogen. The frozen embryos were ground to a fine powder in the presence of liquid nitrogen and then extracted by Polytron homogenization and fractionated to enrich for total RNA by the method of Chirgwin et al. (Biochemistry (1979) 18:5294-5299). The nucleic acid fraction was enriched for poly A<sup>+</sup>RNA by passing total RNA through an oligo-dT cellulose column and eluting the poly A<sup>+</sup>RNA with salt as described by Goodman et al. (Meth. Enzymol. (1979) 68:75-90). cDNA was synthesized from the purified poly A<sup>+</sup>RNA using cDNA Synthesis System (Bethesda Research Laboratory) and the manufacturer's instructions. The resultant double-stranded DNA was methylated by Eco RI DNA methylase (Promega) prior to filling-in its ends with T4 DNA polymerase (Bethesda Research Laboratory) and blunt-end ligation to phosphorylated Eco RI linkers using T4 DNA ligase (Pharmacia). The double-stranded DNA was digested with Eco RI enzyme, separated from excess linkers by passage through a gel filtration column (Sephacrose CL-4B), and ligated to lambda ZAP vector (Stratagene) according to manufacturer's instructions. Ligated DNA was packaged into phage using the Gigapack packaging extract (Stratagene) according to manufacturer's instructions. The resultant cDNA library

was amplified as per Stratagene's instructions and stored at -80°C.

- Following the instructions in the Lambda ZAP Cloning Kit Manual (Stratagene), the cDNA phage library
- 5 was used to infect *E. coli* BB4 cells and approximately 80,000 plaque forming units were plated onto 150 mm diameter petri plates. Duplicate lifts of the plates were made onto nitrocellulose filters (Schleicher & Schuell). The filters were prehybridized in 25 mL of
- 10 hybridization buffer consisting of 50mM Tris-HCl, pH 7.5, 1 M NaCl, 1% SDS, 5% dextran sulfate and 0.1 mg/mL denatured salmon sperm DNA (Sigma Chemical Co.) at 50°C for 2 h. Radiolabeled probe prepared from pCF3 as described above was added, and allowed to hybridize for
- 15 18 h at 50°C. The probes were washed twice at room temperature with 2X SSPE, 1% SDS for five minutes followed by washing for 5 min at 50°C in 0.2X SSPE, 1% SDS. Autoradiography of the filters indicated that there was one strongly hybridizing plaque, and
- 20 approximately five weakly hybridizing plaques. The more strongly hybridizing plaque was subjected to a second round of screening as before, excepting that the final wash was for 5 min at 60°C in 0.2X SSPE, 1% SDS. Numerous, strongly hybridizing plaques were observed,
- 25 and one, well-isolated from other phage, was picked for further analysis.

- Following the Lambda ZAP Cloning Kit Instruction Manual (Stratagene), sequences of the pBluescript
- 30 vector, including the cDNA inserts, from the purified phage was excised in the presence of a helper phage and the resultant phagemid was used to infect *E. coli* XL-1 Blue cells. DNA from the plasmid, designated pXF1, was made by the alkaline lysis miniprep procedure described in Sambrook et al. (Molecular Cloning, A Laboratory
- 35 Manual, 2nd ed. (1989) Cold Spring Harbor Laboratory



Press). The alkali-denatured double-stranded DNA from pXF1 was completely sequenced on both strands. The insert of pXF1 contained a stretch of 1783 nucleotides which contained an unknown open-reading frame and also contained a poly-A stretch of 16 nucleotides 3' to the open reading frame, from nucleotides 1767 to 1783, followed by an Eco RI restriction site. The 2184 bases that followed this Eco RI site contained a 1145 bp open reading frame which encoded a polypeptide of about 68% identity to, and colinear with, the Arabidopsis delta-15 desaturase polypeptide listed in SEQ ID No:2. The putative start methionine of the 1145 bp open-reading frame corresponded to the start methionine of the Arabidopsis microsomal delta-15 peptide and there were no amino acids corresponding to a plastid transit peptide 5' to this methionine. When the insert in pXF1 was digested with Eco RI four fragments were observed, fragments of approximately 370 bp and 1400 bp fragments, derived from the first 1783 bp of the insert in pXF1, and fragments of approximately 600 bp and 1600 bp derived from the the other 2184 nucleotides of the insert in pXF1. Only the 600 bp and 1600 bp fragments hybridized with probe derived from pCF3 on Southern blots. It was deduced that pXF1 contained two different cDNA inserts separated by an Eco RI site and the second of these inserts was a 2184 bp cDNA encoding a soybean microsomal delta-15 desaturase. The complete nucleotide sequence of the 2184 bp soybean microsomal delta-15 cDNA contained in plasmid pXF1 is listed in SEQ ID No:10.

Cloning of a Soybean Seed cDNA Encoding a Plastid  
Delta-15 Glycerolipid Desaturase Using  
Soybean Microsomal Delta-15 Desaturase cDNA  
as an Hybridization Probe

A 1.0 kb fragment of the coding region of the soybean microsomal delta-15 desaturase cDNA contained in

09210035.122300

plasmid pXF1 was excised by digestion with the restriction enzyme Hha I. This 1.0 Kb fragment was purified by agarose gel electrophoresis and radiolabeled with  $^{32}\text{P}$  as previously described. The radiolabeled

5 fragment was used to screen 100,000 plaque-forming units of the the soybean cDNA library as described above. Autoradiography of the filters indicated that there were eight hybridizing plaques and these were subjected to a second round of screening. Sequences of the pBluescript

10 vector from all eight of the purified phages, including the cDNA inserts, were excised in the presence of a helper phage and the resultant phagemids were used to infect *E. coli* XL-1 Blue cells. DNA from the plasmids was made by the alkaline lysis miniprep procedure

15 described in Sambrook et al. (Molecular Cloning, A Laboratory Manual, 2nd ed. (1989) Cold Spring Harbor Laboratory Press). Restriction analysis showed they contained inserts ranging from 1.0 kb to 3.0 kb in size. One of these inserts, designated pSFD-118bwp, contained

20 an insert of about 1700 bp. The alkali-denatured double-stranded DNA from pSFD-118bwp was completely sequenced on both strands. The insert of pSFD-118bwp contained a stretch of 1675 nucleotides which contained an open-reading frame encoding a polypeptide of about

25 80% identity with, and colinear with, the Arabidopsis plastid delta-15 desaturase polypeptide listed in SEQ ID No:5. The open-reading frame also encoded amino acids corresponding to a plastid transit peptide at the 5' end of the open-reading frame. The transit peptide was

30 colinear with, and shared some homology to, the transit peptide described for the *Arabidopsis* plastid delta-15 glycerolipid desaturase. Based on the homology to *Arabidopsis* plastid delta-15 glycerolipid desaturase and because of the presence of a plastid transit peptide,

35 the cDNA contained in plasmid pSFD-118bwp was deduced to

be a soybean plastid delta-15 glycerolipid desaturase. The complete nucleotide sequence of the 1675 bp soybean plastid delta-15 glycerolipid desaturase cDNA is listed in SEQ ID NO:12.

5

#### EXAMPLE 6

#### CLONING OF cDNA SEQUENCES ENCODING FATTY ACID DESATURASES BY POLYMERASE CHAIN REACTION

Analysis of the deduced protein sequences of the different higher plant glycerolipid desaturases described in this invention reveals to those skilled in the art regions of the amino acid sequences that have been conserved among higher plants and between higher plants and cyanobacterial *des A*. These short stretches of amino acids can be used to design oligomers as primers for polymerase chain reactions. Two amino acid sequences that are highly conserved between the *des A* and plant delta-15 desaturases polypeptides are amino acid sequences 97-108 and 299-311 (SEQ ID NO:2). Polymerase chain reactions (PCRs) were performed using GeneAmp® RNA PCR Kit (Perkin Elmer Cetus) following manufacturer's protocols. In one PCR experiment, SEQ ID NOS:22 and 23 were used as sense primers and either SEQ ID NOS:24 and 25 or SEQ ID NOS:26 and 27 as antisense primers on poly A+ RNA purified from both *Arabidopsis* leaf and canola developing seeds. For this, ca. 100 ng of polyA+ RNA was isolated as described previously and reverse-transcribed using the kit using random hexamers. Then the cDNA was used in PCR using 64 pmoles each of SEQ ID NOS:22 and 23 as sense primers and either a mixture of 64 pmoles of SEQ ID NO:24 and 78 pmoles of SEQ ID NO:25 or a mixture 35 pmoles of SEQ ID NO:26 and 50 pmoles of SEQ ID NO:27 by the following program: a) 1 cycle of 2 min at 95°C and 15 C at 50°C, b) 30 cycles of 3 min at 65°C (extension), 1 min 20 sec at 95°C (denaturation), 2 min at 50°C (annealing), and c) 1

35

00210025-122300

cycle of 7 min at 65°C. PCR products were analyzed by gel electrophoresis. All PCRs resulted in PCR products of the correct size (ca. 630 bp). The PCR products from Arabidopsis and canola were purified and used as

5 radiolabeled hybridization probes to screen the Lambda Yes Arabidopsis cDNA library at low stringency, as described above. This led to the isolation of a pure phage, which was excised to give plasmid pYacp7. The cDNA insert in pYacp7 was partially sequenced. Its

10 sequence showed that it encoded an incomplete desaturase polypeptide that was identical to another cDNA (in plasmid pFadx-2) isolated by low-stringency hybridization as described previously. The composite sequence derived from the partial sequences from the

15 cDNA inserts in pFadx-2 and pYacp7 is shown in SEQ ID NO:16 and the polypeptide encoded by it in SEQ ID NO:17. As discussed previously, SEQ ID NO:17 is a putative plastid delta-15 desaturase. A full-length version of pYacp7 can be readily isolated using it has a

20 hybridization probe.

Two additional conserved regions correspond to aminoacid residues 130 to 137 and 249 and 256 of SEQ ID NO:7 (Brassica napus glycerolipid desaturase delta-15). Degenerate oligomers were designed to these regions with

25 additional nucleotides containing a restriction site for Bam H1 were added to the 5' ends of each oligonucleotide to facilitate subcloning of the PCR products. The nucleotide sequences of these oligonucleotides named F2-3 and F2-3c are shown in SEQ ID NO:18 and SEQ ID

30 NO:19 respectively.

Mixtures of degenerate oligonucleotides F2-3 and F2-3c were used to amplify, isolate and clone glycerolipid desaturase sequences represented in corn seed mRNA population, essentially as described in the GeneAmp RNA

35 PCR Kit purchased from Perkin Elmer Cetus and in Innis,

et al., Eds, (1990) PCR Protocols: A Guide to Methods and Applications, Academic Press, San Diego.

- Corn seed RNA was obtained from developing corn seeds 15-20 days after pollination by the method of Chirgwin et al., (1979) Biochemistry 18:5294. Corn seed polyadenylated mRNA was isolated by affinity chromatography on oligo-dT cellulose (Aviv et al., Proc. Natl. Acad. Sci. USA (1972) 69:1408-1411). 20-50ng of A+mRNA were used in reverse transcription reactions with oligo-dT and random hexamers primers using the reaction buffer and conditions recommended by Perkin Elmer Cetus. The resulting cDNA was then used as template for the amplification of corn seed glycerolipid sequences using the set of degenerate primers in SEQ ID NO: 18 and 19.
- Reaction conditions were as described by Perkin Elmer Cetus, the amplification protocol consisted of a sequence of 95°C/1 min, 55°C/1 min, 72°C/2 min for 30-50 cycles. The resulting polymerase reaction products were phenol-chloroform extracted, digested with Bam HI and separated from unincorporated primers by gel filtration chromatopgraphy on Linker 6 spin columns (Pharmacia Inc.). The resulting PCR products were cloned into pBluescript SK at the Bam HI site, and transformed into *E. coli* DH5 competent cells. Restriction analysis of plasmid DNA from the transformed colonies obtained revealed a colony, PCR-20, that contained an insert of about 0.5 kB in size at the pBluescript SK BamHI site. The PCR-20 insert was completely sequenced on both strands. The nucleotide sequence of PCR20 insert is shown in SEQ ID NO:14 and the translated amino acid sequence is shown in SEQ ID NO:15. This aminoacid sequence shows an overall identity of 61.9% to the aminoacid sequence of Brassica napus microsomal delta-15 deaturase shown in SEQ ID NO:7. This result identifies the PCR20 insert as a polymerase reaction product of a

corn seed delta-15 desaturase cDNA. PCR20 insert may be used as a probe to readily isolate full length corn seed delta-15 desaturase cDNAs or as such to antisense or cosuppress corn seed glycerolipid delta-15 desaturase gene expression in transgenic corn plants by cloning it in the appropriate corn gene expression vector.

#### EXAMPLE 7

USE OF THE ARABIDOPSIS THALIANA DELTA-15 DESATURASE GENOMIC CLONES AS A RESTRICTION FRAGMENT LENGTH POLYMORPHISM (RFLP) MARKERS TO MAP THE DELTA-15 DESATURASE LOCI IN ARABIDOPSIS

DNA flanking the T-DNA insertion site in mutant line 3707 was used to map the genetic locus encoding the delta-15 desaturase of Arabidopsis thaliana seeds. An approximately 12 kB genomic DNA fragment containing the Arabidopsis delta-15 desaturase coding sequence was removed from the lambda-4211 clone by digestion with restriction endonuclease Xho I, separated from the Lambda arms by agarose gel electrophoresis, and purified using standard procedures. The isolated DNA was labeled with <sup>32</sup>P using a random priming kit from Pharmacia under conditions recommended by the manufacturer. The radioactive DNA was used to probe a Southern blot containing genomic DNA from Arabidopsis thaliana (ecotype Wassileskija and marker line W100 ecotype Landesberg background) digested with one of several restriction endonucleases. Following hybridization and washes under standard conditions (Sambrook et al., Molecular Cloning: A Laboratory Manual, 2nd ed. (1989) Cold Spring Harbor Laboratory Press), autoradiograms were obtained. Different patterns of hybridization (polymorphisms) were identified in digests using restriction endonucleases Bgl II, Cla I, Hind III, Nsi I, and Xba I. The same radiolabeled DNA fragment was used to map the polymorphism essentially as described by

Helentjaris et al., (Theor. Appl. Genet. (1986) 72:761-769). The radiolabeled DNA fragment was applied as described above to Southern blots of Xba I digested genomic DNA isolated from 117 recombinant inbred progeny (derived from single-seed descent lines to the F<sub>6</sub> generation) resulting from a cross between Arabidopsis thaliana marker line W100 and ecotype Wassileskija (Burr et al., Genetics (1988) 118:519-526). The bands on the autoradiograms were interpreted as resulting from inheritance of either paternal (ecotype Wassileskija) or maternal (marker line W100) DNA or both (a heterozygote). The resulting segregation data were subjected to genetic analysis using the computer program Mapmaker (Lander et al., Genomics (1987) 1:174-181). In conjunction with previously obtained segregation data for 63 anonymous RFLP markers and 9 morphological markers in Arabidopsis thaliana (Chang et al., Proc. Natl. Acad. Sci. USA (1988) 85:6856-6860; Nam et al., Plant Cell (1989) 1:699-705), a single genetic locus was positioned corresponding to the genomic DNA containing the delta-15 desaturase coding sequence. The location of the delta-15 desaturase gene was thus determined to be on chromosome 2 between the lambda AT283 and cosmid c6842 RFLP markers, near the py and erecta morphological markers.

The cDNA in plasmid pCM2 was also shown to hybridize polymorphically to genomic DNA from Arabidopsis thaliana (ecotype Wassileskija and marker line W100 ecotype Landesberg background) digested with Eco RI. It was used as a RFLP marker to map the genetic locus for the gene encoding this fatty acid desaturase in Arabidopsis as described above. A single genetic locus was positioned corresponding to this desaturase cDNA. Its location was thus determined to be on chromosome 3 between the lambda AT228 and cosmid c3838

RFLP markers, "north" of the glabrous locus (Chang et al., Proc. Natl. Acad. Sci. USA (1988) 85:6856-6860; Nam et al., Plant Cell (1989) 1:699-705).

#### EXAMPLE 8

#### 5      USE OF SOYBEAN SEED MICROSOMAL DELTA-15 GLYCEROLIPID DESATURASE cDNA SEQUENCE IN PLASMID AS A RESTRICTION FRAGMENT LENGTH POLYMORPHISM (RFLP) MARKER

A 600 bp fragment of the cDNA insert from plasmid pXF1, which contains about 300 bp of the coding sequence and 300 bp of the 3' untranslated sequence, was excised  
10 by digestion with restriction enzyme Eco RI in standard conditions as described in Sambrook et al. (Molecular Cloning, A Laboratory Manual, 2nd ed. (1989) Cold Spring Harbor Laboratory Press), purified by agarose gel  
15 electrophoresis and labeled with <sup>32</sup>P using a Random Priming Kit from Bethesda Research Laboratories under conditions recommended by the manufacturer. The resulting radioactive probe was used to probe a Southern blot (Sambrook et al., Molecular Cloning, A Laboratory  
20 Manual, 2nd ed. (1989) Cold Spring Harbor Laboratory Press) containing genomic DNA from soybean [Glycine max (cultivar Bonus) and Glycine soja (PI81762)], digested with one of several restriction enzymes. After hybridization and washes under standard conditions  
25 (Sambrook et al. Molecular Cloning, A Laboratory Manual, 2nd ed. (1989), Cold Spring Harbor Laboratory Press), autoradiograms were obtained and different patterns of hybridization (polymorphisms) were identified in digests performed with restriction enzymes Bam HI, Eco RV and  
30 Eco RI. The same probe was then used to map the polymorphic pXF1 locus on the soybean genome, essentially as described by Helentjaris et al. (Theor. Appl. Genet. (1986) 72:761-769). Plasmid pXF1/600 bp probe was applied, as described above, to Southern blots  
35 of EcoRI, PstI, EcoRV, BamHI, or Hin DIII digested



genomic DNAs isolated from 68 F2 progeny plants resulting from a G. max Bonus x G. soja PI81762 cross. The bands on the autoradiograms were interpreted as resulting from the inheritance of either paternal (Bonus) or maternal (PI81762) pattern, or both (a heterozygote). The resulting data were subjected to genetic analysis using the computer program Mapmaker (Lander et al., Genomics (1987) 1:174-181). In conjunction with previously obtained data for 436 anonymous RFLP markers in soybean (Tingey et al., J. Cell. Biochem., Supplement 14E (1990) p. 291, abstract R153], Applicants were able to position a single genetic locus corresponding to the pXF1/600 bp probe on the soybean genetic map. This confirms that the gene for microsomal delta-15 desaturase is located on chromosome 19 in the soybean genome. This information will be useful in soybean breeding targeted towards developing lines with altered polyunsaturate levels.

#### EXAMPLE 9

##### OVEREXPRESSION OF MICROSOMAL DELTA-15 FATTY ACID DESATURASE IN PLANTS

Detailed procedures for DNA manipulation, such as use of restriction endonucleases and other DNA modifying enzymes, agarose gel electrophoresis, isolation of DNA from agarose gels, transformation of E. coli cells with plasmid DNA, and isolation and sequencing of plasmid DNA are described in Sambrook et al. (1989) Molecular cloning, A Laboratory Manual, 2nd ed. Cold Spring Harbor Laboratory Press and Ausubel et al. (1989) Current Protocols in Molecular Biology John Wiley & Sons. All restriction enzymes and modifying enzymes were obtained from Bethesda Research Laboratory, unless otherwise noted.

To test the biological effect of overexpression of the microsomal delta-15 desaturase SEQ ID NO:1, i.e., the cDNA encoding Arabidopsis thaliana microsomal delta-15 desaturase, was placed in the sense orientation

5 behind either the CaMV 35S promotor, to provide constitutive expression, or behind the promotor for the gene encoding soybean  $\alpha'$  subunit of the  $\beta$ -conglycinin (7S) seed storage protein, to provide embryo-specific expression. To create the chimeric gene constructs,

10 specific expression cassettes were made to facilitate easy manipulation of the desired clones. The chimeric genes were then transformed into plant cells by Agrobacterium tumefaciens's binary Ti plasmid vector system [Hoekema et al., (1983) Nature 303:179-180; Bevan

15 (1984) Nucl. Acids Res. 12:8711-8720].

Overexpression of Arabidopsis Delta-15 Fatty Acid  
Desaturase in Transgenic Carrot Hairy Roots

To confirm the identity of SEQ ID NO:1 (Arabidopsis microsomal delta-15 fatty acid desaturase) and to test

20 the biological effect of its overexpression in a heterologous plant species, the constitutive chimeric gene 35S:SEQ ID NO:1 was introduced into carrot tissue by Agrobacterium. The cassette for constitutive gene expression in plasmid, pAW28, originated from pK35K

25 which, in turn, is derived from pKNK. Plasmid pKNK is a pBR322-based vector containing a chimeric gene for plant kanamycin resistance: nopaline synthase (NOS) promoter/neomycin phosphotransferase (NPT) II coding region/3' NOS chimeric gene. Plasmid pKNK has been

30 deposited on 7 January 1987 with the American Type Culture Collection of Rockville, Maryland, USA under the provisions of the Budapest Treaty and bears the deposit accession number 67284. A map of this plasmid is shown in Lin, et al., Plant Physiol. (1987) 84:856-861. The

35 NOS promoter region is a 296 bp Sau 3A-Pst I fragment

corresponding to nucleotides -263 to +33, with respect to the transcription start site, of the NOS gene described by Depicker et al. (1982) J. Appl. Genet. 1:561-574. The Pst I site at the 3' end was created at the translation initiation codon of the NOS gene. The NptII coding region is a 998 bp Hind III-Bam HI fragment obtained from transposon Tn5 (Beck et al., Gene (1982) 19:327-336) by the creation of Hind III and Bam HI sites at nucleotides 1540 and 2518, respectively. The 3' NOS is a 702 bp Bam HI-Cla I fragment from nucleotides 848 to 1550 of the 3' end of the NOS gene (Depicker et al., J. Appl. Genet. (1982) 1:561-574) including its polyadenylation region. pKNK was converted to pK35K by replacing its Eco RI-Hind III fragment containing the NOS promoter with a Eco RI-Hind III fragment containing the CaMV 35S promoter. The Eco RI-Hind III 35S promoter fragment is the same as that contained in pUC35K that has been deposited on 7 January 1987 with the American Type Culture Collection under the provisions of the Budapest Treaty and bears the deposit accession number 67285. The 35S promoter fragment was prepared as follows, and as described in Odell et al., Nature (1985) 313:810-813, except that the 3' end of the fragment includes CaMV sequences to +21 with respect to the transcription start site. A 1.15 KB Bgl II segment of the CaMV genome containing the region between -941 and +208 relative to the 35S transcription start site was cloned in the Bam HI site of the plasmid pUC13. This plasmid was linearized at the Sal I site in the polylinker located 3' to the CaMV fragment and the 3' end of the fragment was shortened by digestion with nuclease Bal31. Following the addition of Hind III linkers, the plasmid DNA was recircularized. From nucleotide sequence analysis of the isolated clones, a 3' deletion fragment was selected with the Hind III

linker positioned at +21. The 35S promoter fragment was isolated as an Eco RI-Hind III fragment, the Eco RI site coming from the polylinker of pUC13.

The NPTII coding region in plasmid pK35K was removed from plasmid pK35K by digestion with Hind III and Bam HI restriction enzymes. Following digestion, the ends of the DNA molecules were filled-in using Klenow enzyme. Not I linkers (New England Biolabs) were then ligated on the ends and the plasmid was recircularized to yield plasmid pK35Nt. A 1.7 kB fragment containing the 35S promoter region - Not I site - 3' untranslated region from nopaline synthase was liberated from pK35Nt using restriction endonucleases Eco RI and Cla I. Following restriction digestion the ends of the DNA molecules were filled-in using Klenow enzyme after which Xho I linkers (New England Biolabs) were added. The 1.7 kB fragment, now containing Xho I sites at either end, was gel isolated and cloned into the plasmid vector pURA3 (Clonetech) at its unique Xho I site. The vector pURA3 was chosen due to the absence of a Not I restriction site, the presence of a single Xho I restriction site and because the relatively large size of the vector (pURA3) would make the isolation of the gene expression cassettes relatively easy from the final construct.

The 1.4 kB Not I fragment in plasmid pCF3 containing Arabidopsis microsomal delta-15 desaturase (SEQ ID NO:1) was isolated and ligated to pAW28 (the constitutive expression cassette) previously linearized with Not I restriction enzyme and treated with calf intestinal alkaline phosphatase (Boehringer Mannheim) to result in plasmids pAW29 and pAW30 that had SEQ ID NO:1 cloned in a sense orientation and antisense orientation, respectively, with respect to the promoter. The orientation of the cDNA relative to the promoters was

established by digestion with appropriate restriction endonucleases or by sequencing across the promotor-cDNA junctions.

The chimeric genes 35S promotor/sense SEQ ID  
 5 NO:1/3'NOS and 35S promotor/antisense SEQ ID NO:1/3'NOS  
 were isolated as a 3 kB Xho I fragment from plasmids  
 pAW29 and pAW30, respectively, and cloned into the  
 binary vector pZS194b at its unique Sal I site to result  
 in plasmids pAW31 and pAW32, respectively. The  
 10 orientation of the plant selectable marker gene in pAW31  
 and pAW32 is the same as that of the 35S promoter as  
 ascertained by digestion with appropriate restriction  
 endonucleases. Binary vector pZS194b contains the  
 pBR322 origin of replication, the replication and  
 15 stability regions of the Pseudomonas aeruginosa plasmid  
 pVS1 [Itoh, et al., (1984) Plasmid 11:206-220] required  
 for replication and maintenance of the plasmid in  
Agrobacterium, the bacterial NPT II gene (kanamycin  
 resistance) from Tn5 [Berg et al., (1975) Proc. Nat'l.  
 20 Acad. Sci. U.S.A. 72:3628-3632] as a selectable marker  
 for transformed bacteria, left and right borders of the  
 T-DNA of the Ti plasmid [Bevan et al., (1984) Nucl.  
 Acids Res. 12:8711-8720], and, between the left and  
 right T-DNA borders are the chimeric NOS:NPT II gene for  
 25 plant kanamycin resistance, described above, as a  
 selectable marker for transformed plant cells and the E.  
coli lacZ a-complementing segment [Vieria and Messing  
 (1982) Gene 19:259-267] with unique restriction  
 endonuclease sites for Kpn I and Sal I.

30 The binary vectors pAW31 and pAW32 were transformed  
 by the freeze/thaw method [Holsters et al. (1978) Mol.  
 Gen. Genet. 163:181-187] into Agrobacterium tumefaciens  
 strain R1000, carrying the Ri plasmid pRiA4b from  
Agrobacterium rhizogenes [Moore et al., (1979) Plasmid

2:617-626] to result in transformants R1000/pAW31 and R1000/pAW32, respectively.

Carrot (Daucus carota L.) cells were transformed by co-cultivation of carrot root disks with strain R1000, R1000/pAW31, or R1000/pAW32 by the method of Petit et al., (1986) Mol. Gen. Genet. 202:388-393]. To prepare explants for inoculation, carrots purchased from the local supermarket were first scrubbed gently with water and dish detergent, then rinsed thoroughly with tap and distilled water. They were surface sterilized in a stirred solution of 50% Clorox and distilled water for 30 min and rinsed thoroughly with sterile distilled water. The carrots were peeled using an autoclaved vegetable peeler and then sliced with a scalpel blade into disks of approximately 5-10 mm thickness. The disks were placed in petri dishes, onto a medium consisting of distilled deionized water solidified with 0.7% agar, in an inverted orientation so that the cut surface nearest to the root apex of the carrot was exposed for inoculation.

Cultures of Agrobacterium strains R1000, R1000/pAW31, and R1000/pAW32 were initiated from freshly grown plates in LB broth plus the appropriate antibiotic selective agents (50 mg/L chloramphenicol for the R1000 or 50 mg/L each of chloramphenicol and kanamycin for R1000/pAW31 and R1000/pAW32) and grown at 28°C to an optical density of around 1.0 at 600 nm. Bacterial cells were pelleted by centrifugation, rinsed and resuspended in LB broth without antibiotics. Freshly cut carrot disks were inoculated by applying 100 µL of the bacterial suspension to the cut surface of each disk. As a control, some disks were inoculated with sterile LB broth only, to indicate the extent of root formation in the absence of Agrobacterium.

Inoculated root disks were incubated at 25°C in the dark in petri dishes sealed with Parafilm. After two weeks of co-cultivation of carrot disks with Agrobacterium, the carrot disks were transferred to fresh agar-solidified water medium containing 500 mg/L carbenicillin for the counterselection of Agrobacterium. At this time, hairy root formation was noted on some root disks. Transfer of the explants to fresh counterselection medium was done at four weeks.

10 Excision of individual roots from the explants was begun at six weeks. Ten days later, additional roots were taken from the explants as needed.

Approximately 5-10 mm long hairy roots were excised and individually subcultured on MS minimal organics medium with 30 g/L sucrose (Gibco, Grand Island, N. Y., Cat. No. 510-1118EA) and 500 mg/L carbenicillin. Approximately equal numbers of roots were subcultured in liquid medium and in a medium solidified with 0.6% agarose. Cultures on solid medium were grown in 60 x 100 mm petri dishes, liquid cultures were in 6-well culture dishes. When excising roots, an effort was made to select single roots from distinct callus-like outgrowths on the wounded surface. These sites of excision were marked on the lid of the petri dish to minimize repeat sampling of tissue originating from the same transformation event.

Two to three weeks after excision from the explants, individual hairy root cultures that were not visibly contaminated with Agrobacterium were transferred to fresh MS medium supplemented with 500 mg/L carbenicillin. The root mass of each culture was cut into segments including one or more branch roots, and these segments were transferred as a group to a plate or well of fresh medium. Approximately 20 mg fresh weight of tissue of root cultures which grew to adequate size

- within the next two to three weeks were sampled for fatty acid composition by gas chromatography of the fatty acyl methyl esters essentially as described by Browse et al., (Anal. Biochem. (1986) 152:141-145)
- 5 except that 2.5%  $H_2SO_4$  in methanol was used as the methylation reagent and samples were heated for 1.5 h at 80°C to effect the methanolysis of the seed triglycerides. The results are shown in Table 6. A second sample of tissue consisting of an actively
- 10 growing root tip of approximately 1 cm was excised and placed on MS medium supplemented with 500 mg/L carbenicillin and 25-50 mg/L kanamycin to test for kanamycin resistance select for hairy roots co-transformed with the binary vector [Simpson et al.
- 15 (1986) Plant Mol. Biol. 6:403-415].



**TABLE 6**  
 Percent 18:3 and 18:2/18:3 Ratio in  
Roots of Transgenic Carrots

<u>Root Sample</u>	<u>Transformation Vector Used</u>	<u>%18:3</u>	<u>%18:2/18:3</u>
1	R1000/pAW31	62	0.09
2	R1000/pAW31	8	7.30
3	R1000/pAW31	10	5.69
4	R1000/pAW31	62	0.06
5	R1000/pAW31	10	5.07
6	R1000/pAW31	4	14.2
7	R1000/pAW31	61	0.18
8	R1000/pAW31	4	15.1
9	R1000/pAW31	61	0.07
10	R1000/pAW31	63	0.09
11	R1000/pAW31	15	3.04
12	R1000/pAW31	64	0.14
13	R1000/pAW31	5	9.94
14	R1000/pAW31	9	6.72
15	R1000/pAW31	8	7.08
16	R1000/pAW31	8	6.31
17	R1000/pAW31	23	1.86
18	R1000/pAW31	8	7.33
19	R1000/pAW31	10	5.99
20	R1000/pAW31	7	8.83
21	R1000/pAW32	9	6.80

<u>Root Sample</u>	<u>Transformation Vector Used</u>	<u>\$18:3</u>	<u>\$18:2/18:3</u>
22	R1000/pAW32	4	11.8
23	R1000/pAW32	3	18.8
24	R1000/pAW32	10	6.21
25	R1000/pAW32	7	8.57
26	R1000/pAW32	3	16.4
27	R1000/pAW32	6	8.29
28	R1000/pAW32	5	9.19
29	R1000/pAW32	5	8.47
30	R1000/pAW32	8	7.17
31	R1000/pAW32	4	11.9
32	R1000/pAW32	8	7.20
33	R1000/pAW32	5	10.4
34	R1000/pAW32	8	7.29
35	R1000/pAW32	3	17.2
36	R1000/pAW32	8	7.27
37	R1000/pAW32	9	6.01
38	R1000/pAW32	9	6.62
40	R1000/pAW32	9	6.02
41	R1000	8	7.23
42	R1000	8	7.83
43	R1000	10	6.20
44	R1000	9	5.97
45	R1000	9	6.73
46	R1000	9	6.27
47	R1000	8	7.27
48	R1000	7	8.30
49	R1000	9	7.11

The ability of R10Q0 transformed "hairy" roots to grow in the absence of exogenous phytohormones can be attributed to the Ri plasmid, pRiA4b. When R1000/pAW31 or R1000/pAW32 strains are used to transform, only a fraction (about half) of the "hairy" roots will also be

transformed with the experimental binary vector, pAW31 or pAW32. Thus, as expected, not all hairy roots resulting from transformation with R1000/pAW31 show the high 18:3 phenotype. The absence of any significant fatty acid phenotype in "hairy roots" transformed with R1000/pAW31 is expected, since carrot and Arabidopsis delta-15 desaturase sequences are not expected to be sufficiently related. These results show that overexpression of Arabidopsis microsomal delta-15 desaturase can result in over 10-fold increase in 18:3 at the expense of 18:2 in heterologous plant tissue.

Overexpression of Arabidopsis Delta-15 Fatty Acid Desaturase in Seeds and Complementation of the Mutation in Delta-15 Desaturation in Mutant 3707

To complement the delta-15 desaturation mutation in the T-DNA mutant 3707 and to test the biological effect of overexpression of SEQ ID NO:1 (Arabidopsis microsomal delta-15 fatty acid desaturase) in seed, the embryo-specific promoter:SEQ ID NO:1 chimeric gene was transformed into the mutant plant. This embryo-specific expression cassette in pAW42 was produced, in part, using a modified version of vector pCW109. Vector pCW109 itself was made by inserting into the Hind III site of the cloning vector pUC18 (Bethesda Research Laboratory) a 555 bp 5' non-coding region (containing the promoter region) of the  $\beta$ -conglycinin gene followed by the multiple cloning sequence containing the restriction endonuclease sites for Nco I, Sma I, Kpn I and Xba I, then 1174 bp of the common bean phaseolin 3' untranslated region into the Hind III site [Slightom et al., Proc. Nat'l Acad. Sci. U.S.A. (1983) 80:1897-1901]. The  $\beta$ -conglycinin promoter region used is an allele of the published  $\beta$ -conglycinin gene (Doyle et al., J. Biol. Chem. (1986) 261:9228-9238) due to differences at 27

00210505-122399

nucleotide positions. Further sequence description may be found in Slightom (W091/13993).

The modifications to vector pCW109 were as follows: The potential translation start site was destroyed by digestion with Nco I and Xba I restriction enzymes followed by treatment with mung bean nuclease (New England Biolabs) to create linear, blunt ended DNA molecules. After ligation of Not I linkers (New England Biolabs) and digestion with Not I restriction enzyme (New England Biolabs) the plasmid was recircularized. Confirmation of the desired change was obtained by dideoxy sequencing. The resulting plasmid was designated pAW35. The 1.8 kB Hind III fragment from pAW35 containing the modified  $\beta$ -conclycinin promotor/3' phaseolin region was subcloned into the Hind III site in plasmid vector pBluescript SK<sup>+</sup> (Stratagene) creating plasmid pAW36. Plasmid pAW36 was linerized at its unique Eco RI site and ligated to Eco RI/Xho I adaptors (Stratagene). Following digestion with Xho I, the 1.7 kB Xho I fragment containing the  $\beta$ -conclycinin promotor/Not I site/3'-phaseolin untranslated region was cloned into the Xho I site in pURA3 vector (Clonetech). The resultant plasmid, pAW42, contains the seed specific expression cassette bordered by Xho I sites to facilitate cloning into modified T-DNA binary vectors and a unique Not I site to facilitate cloning of target cDNA sequences. Vector pURA3 was choosen due to the absence of a Not I restriction site, the presence of a single Xho I restriction site, and the relatively large size of the vector (pURA3) would make the isolation of the gene expression cassettes relatively easy from the final construct.

The 1.4 kB Not I fragment in plasmid pCF3 containing Arabidopsis microsomal delta-15 desaturase (SEQ ID NO:1) was isolated and ligated to plasmid pAW42

- (the seed-specific expression cassette) that had previously been linearized with Not I restriction enzyme and treated with calf intestinal alkaline phosphatase (Boehringer Mannheim) to result in plasmids pAW45 that
- 5 had SEQ ID NO:1 cloned in a sense orientation with respect to the promoter. The orientation of the cDNA relative to the promoters was established by digestion with appropriate restriction endonucleases or by sequencing across the promoter-cDNA junctions.
- 10 The chimeric  $\beta$ -conclycinin promoter/sense SEQ ID NO:1/phaseolin 3' was isolated as a 3.2 kB Xho I fragment from plasmid pAW45 and subcloned into the binary vector pAW25 at its unique Sal I site. In the resulting vector, pAW50, the orientation of the plant
- 15 selectable marker is the same as that of the  $\beta$ -conclycinin promoter as ascertained by digestion with appropriate restriction endonucleases. Plasmid pAW25, is derived from plasmids pZS94K and pML2. Plasmid pZS94K contains the pBR322 origin of replication, the
- 20 replication and stability regions of the Pseudomonas aeruginosa plasmid pVS1 [Itoh, et al., (1984) Plasmid 11:206-220] required for replication and maintenance of the plasmid in Agrobacterium, the bacterial NPT II gene (kanamycin resistance) from Tn5 [Berg et al., (1975)
- 25 Proc. Nat'l. Acad. Sci. U.S.A. 72:3628-3632] as a selectable marker for transformed bacteria, a T-DNA left border fragment of the octopine Ti plasmid pTiA6 and right border fragment derived from TiAch5 describe by van den Elzen et al. (Plant Mol. Biol. (1985)
- 30 5:149-154). Between these borders are the E. coli lacZ a-complementing segment [Vieria and Messing (1982) Gene 19:259-267] with restriction endonuclease sites Sal I and Asp 718 derived from pUC18. A 4.5 kB Asp 718-Sal I DNA fragment containing the chimeric herbicide
- 35 sulfonylurea (SU)-resistant acetolactate (ALS) gene was

8/21/93 5:12:38

obtained from plasmid pML2 and cloned into the Asp 718-Sal I sites of plasmid pZS94K. This chimeric ALS gene contained the CaMV 35S promoter/Cab22L Bgl II-Nco I fragment that is described by Harpster et al., [Mol. Gen. Genet. (1988) 212:182-190] and the Arabidopsis ALS coding and 3' non-coding sequences [Mazur et al., (1987) Plant Physiol. 85:1110-1117] that was mutated so that it encodes a SU-resistant form of ALS. The mutation, introduced by site-directed mutagenesis, are those present in the tobacco SU-resistant Hra gene described by Lee et al., (1988) EMBO J. 5:1241-1248. The resulting plasmid was designated pAW25.

The binary vector pAW25 containing the chimeric embryo-specific  $\beta$ -conglycinin promoter:sense SEQ ID NO:1 gene was transformed by the freeze/thaw method [Holsters et al., (1978) Mol. Gen. Genet. 163:181-187] into the avirulent Agrobacterium strain LBA4404/pAL4404 [Hoekema et al., (1983) Nature 303:179-180].

Arabidopsis root cultures were transformed by co-cultivation with Agrobacterium using standard aseptic techniques for the manipulation of sterile media and axenic plant/bacterial cultures were followed, including the use of a laminar flow hood for all transfers. Compositions of the culture media are listed in Table 8. Unless otherwise indicated, 25x100 mm petri plates were used for plant tissue cultures. Incubation of plant tissue cultures was at 23°C under constant illumination with mixed fluorescent and "Gro and Sho" plant lights (General Electric) unless otherwise noted. To initiate in vitro root cultures of the T-DNA homogyzous mutant line 3707 (Arabidopsis thaliana (L.) Heynh, geographic race Wassilewshija) seeds of the mutant line were sterilized for 10 min in a solution of 50% Chlorox with 0.1% SDS, rinsed 3 to 5 times with sterile dH<sub>2</sub>O, dried thoroughly on sterile filter paper, and then 2-3 seeds

0021935-122398

were sown in liquid B5 medium in 250 mL Belco flasks. The flasks were capped, placed on a rotary shaker at 70-80 rpm, and incubated for 3-4 weeks. Prior to inoculation with Agrobacterium, root tissues were

5 cultured on callus induction medium (MSKig). Roots were harvested by removing the root mass from the Belco flask, placing it in a petri dish, and, using forceps, pulling small bundles of roots from the root mass and placing them on MSKig medium. Petri dishes were sealed

10 with filter tape and incubated for four days.

Agrobacterium strain LBA4404 carrying the plasmids pAL4404 and pAW50 were grown in 5 mL of YEB broth containing 25 mg/L kanamycin and 100 mg/L rifampicin. The culture was grown for approximately 17-20 h in glass

15 culture tubes in a New Brunswick platform shaker (225 rpm) maintained at 28°C. Pre-cultured roots were cut into 0.5 cm segments and placed in a 100  $\mu$ m filter, made from a Tri-Pour beaker (VWR Scientific, San Francisco, CA USA) and wire mesh, which is set in a petri dish.

20 Root segments were inoculated for several min in 30-50 mL of a 1:20 dilution of the overnight Agrobacterium culture with periodic gentle mixing. Inoculated roots were transferred to sterile filter paper to draw off most of the liquid. Small bundles of roots, consisting

25 of several root segments, were placed on MSKig medium containing 100  $\mu$ m acetosyringone (3',5'-Dimethoxy-4'-hydroxyaceto-phenone, Aldrich Chemical Co., Milwaukee, WI, USA). Petri plates were sealed with parafilm or filter tape and incubated for 2 to 3 days.

30 After infection, root segments were rinsed and transferred to shoot induction medium with antibiotics. Root bundles were placed in a 100  $\mu$ m filter unit (described above) and rinsed with 30-50 mL liquid MSKig medium. The filter was vigorously shaken in the

35 solution to help remove the Agrobacterium, transferred

to a clean petri dish, and rinsed again. Roots were blotted on sterile filter paper and bundles of roots were placed on MSg medium containing 500 mg/l vancomycin and either 10 or 20 ppb chlorsulfuron. Plates were  
 5 sealed with filter tape and incubated for 12 to 14 days.

Green nodules and small shoot primordia were visible at about 2-3 weeks. The explants were either left intact or were broken into numerous pieces and placed on GM medium containing 200-300 mg/L vancomycin  
 10 and either 10 or 20 ppb chlorsulfuron for further shoot development. Plates were either sealed with two pieces of tape or with filter tape. As they developed, individual shoots were isolated from the callus and were placed on MSrg medium containing 100 mg/L vancomycin and  
 15 either 10 or 20 ppb chlorsulfuron. Dishes were sealed as described above and incubated for seven to 10 days. Shoots were then transferred to GM medium containing 100-200 mg/L vancomycin in 25x100 petri dishes or Magenta G7 vessels. Many primary transformants (T1)  
 20 which were transferred to individual containers set seed (T2).

T2 seed was harvested from selected putative transformants and sown on GM medium containing 10ppb chlorsulfuron. Plates were sealed with filter tape,  
 25 cold treated for 2 or more days at 4°C, and then incubated for 10 to 20 days at 23°C under constant illumination as described above. Seedlings were scored as resistant (green, true leaves develop) and sensitive (no true leaves develop).

30 Selected chlorsulfuron resistant T2 seedlings were transplanted to soil and were grown to maturity at 23°C daytime (16 h) 18°C nighttime (8 h) at 65-80% relative humidity.

T2 seeds from two plants were harvested at maturity  
 35 and analysed individually for fatty acid composition by



gas chromatography of the fatty acyl methyl esters essentially as described by Browse et al., (Anal. Biochem. (1986) 152:141-145) except that 2.5% H<sub>2</sub>SO<sub>4</sub> in methanol was used as the methylation reagent and samples  
 5 were heated for 1.5 h at 80°C to effect the methanolysis of the seed triglycerides. The results are shown in Table 7.

TABLE 7  
 Percent Fatty Acid in Seeds of  
 Transgenic Mutant 3707

Seed Sample	16:0	18:0	18:1	18:2	18:3
wildtype(6)	6	4	14	30	19
mutant 3707(6)	6	4	14	44	3
1-1	10	4	22	9	55
1-2	11	6	22	14	48
1-3	12	7	16	6	57
1-4	10	4	30	52	4
1-5	10	4	18	17	48
1-6	10	5	15	15	53
2-1	11	5	19	60	4
2-2	10	5	19	9	56
2-3	9	4	27	8	52
2-4	10	5	17	10	56
2-5	10	5	19	9	56
2-6	10	5	17	17	48

The fatty acid composition of the wild-type and mutant line 3707 represents the average of 6 single  
 10 seeds each. Seeds from plant 1 are designated 1-1 to 1-6 and those from plant 2 are designated 2-1 to 2-6. The 20:1 and 20:2 amounts are not shown. The data shows that the one out of six seeds in each plant show the mutant fatty acid phenotype, while the remaining seeds  
 15 show more than 10-fold increase in 18:3 to ca.55%. While most of the increase occurs at the expense of 18:2, some of it also occurs at the expense of 18:1.

00240325-120394

Such high levels are of linolenic acid in vegetable oils are observed in specialty oil crops, such as linseed. Thus, overexpression of this gene in other oilscrops, especially canola, which is a close relative of *Arabidopsis*, is also expected to result in such high levels of 18:3.

TABLE 8

## Medium Composition

YEP MEDIUM		BASIC MEDIUM	
Bacto Beef Extract	5.0 g	1 Pkg. Murashige and Skoog Minimal Organics Medium without Sucrose (Gibco #510-3118 or Sigma #M6899)	
Bacto Yeast Extract	1.0 g		
Peptone	5.0 g	10 mL Vitamin Supplement	
Sucrose	5.0 g		
MgSO <sub>4</sub> ·7H <sub>2</sub> O	0.5 g	0.05% MES	0.5 g/L
Agar (optional)	15.0 g	0.8% agar	8 g/L
pH		pH	
VITAMIN SUPPLEMENT		GM = Germination Medium	
10 mg/L thiamine		Basic Medium	
50 mg/L pyridoxine		1% sucrose	10 g/L
50 mg/L nicotinic acid			
MSKig = Callus Induction Medium		MSg = Shoot Induction Medium	
Basic Medium		Basic Medium	
2% glucose	20 g/L	2% glucose	20 g/L
0.5 mg/L 2,4-D	2.3 µL	0.15 mg/L IAA	0.86 µM
0.3 mg/L Kinetin	1.4 µM	5.0 mg/L 2iP	24.6 µM
5 mg/L IAA	28.5 µM		
MSRg = Shoot Induction Medium			
Basic Medium			
2% glucose	20 g/L		
12 mg/L IBA	58.8 µM		
0.1 mg/L Kinetin	0.46 µM		

EXAMPLE 10

Construction of Vectors for Transformation  
of Brassica napus for Reduced Expression  
of Delta-15 Desaturases in Developing Seeds

5 Detailed procedures for manipulation of DNA  
fragments by restriction endonuclease digestion, size  
separation by agarose gel electrophoresis, isolation of  
DNA fragments from agarose gels, ligation of DNA  
fragments, modification of cut ends of DNA fragments and  
10 transformation of E. coli cells with circular DNA  
plasmids are all described in Sambrook et al.,  
(Molecular Cloning, A Laboratory Manual, 2nd ed (1989)  
Cold Spring Harbor Laboratory Press) and Ausubel et al.,  
Current Protocols in Molecular Biology (1989) John Wiley  
15 & Sons).

Sequences of the cDNA's encoding the B. napus  
cytoplasmic delta-15 desaturase and the Brassica napus  
plastid delta-15 desaturase were placed in the antisense  
orientation behind the promoter region from the a'  
20 subunit of the soybean storage protein  $\beta$ -conglycinin to  
provide embryo specific expression and high expression  
levels.

An embryo-specific expression cassette was  
constructed to serve as the basis for chimeric gene  
25 constructs for anti-sense expression of the nucleotide  
sequences of delta-15 desaturase cDNAs. The vector  
pCW109 was produced by the insertion of 555 base pairs  
of the  $\beta$ -conglycinin (a' subunit of the 7s seed storage  
protein) promoter from soybean (Glycine max), the  
30  $\beta$ -conglycinin 5' untranslated region followed by a  
multiple cloning sequence containing the restriction  
endonuclease sites for Nco I, Sma I, Kpn I and Xba I,  
then 1174 base pairs of the common bean phaseolin 3'  
untranslated region into the Hind III site in the  
35 cloning vector pUC18 (BRL). The  $\beta$ -conglycinin promoter

5

1

1

2

2

(3)

orientation. The antisense construction was given the name pCCFdR1.

The transcription unit [ $\beta$ -conglycinin promoter:antisense delta-15 desaturase:phaseolin 3'end] was released from pCCFdR1 by Hind III digestion, isolated, and ligated into pBluescript which had also been Hind III digested to give plasmid pCCFdR2. This construct has unique BamH I and Sal I sites which were digested. The 3 kB transcriptional unit was isolated and cloned into the Bam HI and Sal I sites in pZ199 described below to give the binary vector pZCC3FdR. The orientation given by this directional cloning is with transcription of both the selectable marker gene and the delta-15 antisense gene in the same direction and toward the right border tDNA sequence.

An antisense construction based on the plastid delta-15 desaturase was made with the 425 most 3' bases of SEQ ID NO:8 which is contained in the plasmid pBNSFD-8. pBNSFD-8 represents a cDNA of the plastid delta-15 desaturase in pBluescript. The cDNA insert was removed from pBNSFD-8 by digestion with Xho I and Sma I, the fragments were blunted, and the 425 base insert isolated by gel purification. The isolated fragment was cloned into the Sma I site of pCW109A and the antisense orientation of the chosen clone confirmed by digestion of the plasmid with Pst I. Pst I cuts in the plastid delta-15 sequence and in the pCW109A vector 5' to the  $\beta$ -conglycinin promoter to release a 1.2 kB fragment indicative of the antisense orientation. The plasmid containing this construction was called pCCdFdR1.

Digestion of pCCdFdR1 with Hind III removes a 2.3 kB fragment containing the transcriptional unit [ $\beta$ -conglycinin promoter:plastid delta-15 antisense:3'-phaseolin sequence]. The fragment was gel isolated and cloned into Hind III digested pBluescript. The

orientation of the fragment was relative to the Bam HI site in the cloning region of pBluescript was determined by digestion with Pst I as described above. A clone oriented with the promoter toward the Sal I containing end was chosen and given the name pCCdFdr2.

pCCdFdr2 was digested with Bam HI and Sal I, the released fragment was gel isolated and ligated into pZ199 which had been digested with Bam HI and Sal I to give the binary vector pZCCdFdr.

Vectors for transformation of the antisense delta-15 desaturase constructions under control of the  $\beta$ -conglycinin promoter into plants using Agrobacterium tumefaciens were produced by constructing a binary Ti plasmid vector system (Bevan, (1984) Nucl. Acids Res. 12:8711-8720). The starting vector used for these systems (pZS199) is based on a vector which contains: (1) the chimeric gene nopaline synthase/neomycin phosphotransferase as a selectable marker for transformed plant cells (Bevan et al., (1984) Nature 304:184-186), (2) the left and right borders of the T-DNA of the Ti plasmid (Bevan et al., (1984) Nucl. Acids Res. 12:8711-8720), (3) the E. coli lacZ a-complementing segment (Vieria and Messing (1982) Gene 19:259-267) with unique restriction endonuclease sites for Eco RI, Kpn I, Bam HI, Hin DIII, and Sal I, (4) the bacterial replication origin from the Pseudomonas plasmid pVS1 (Itoh et al., (1984) Plasmid 11:206-220), and (5) the bacterial neomycin phosphotransferase gene from Tn5 (Berg et al., (1975) Proc. Natnl. Acad. Sci. U.S.A. 72:3628-3632) as a selectable marker for transformed A. tumefaciens. The nopaline synthase promoter in the plant selectable marker was replaced by the 35S promoter (Odell et al. (1985) Nature, 313:810-813) by a standard restriction endonuclease digestion and ligation strategy. The 35S promoter is

required for efficient Brassica napus transformation as described below.

#### EXAMPLE 11

##### AGROBACTERIUM MEDIATED TRANSFORMATION

##### OF BRASSICA NAPUS

5           The binary vectors pZCC3FdR abd pZCCdFdR were transferred by a freeze/thaw method (Holsters et al., (1978) Mol Gen Genet 163:181-187) to the Agrobacterium strain LBA4404/pAL4404 (Hoekema et al., (1983), Nature 10 303:179-180).

Brassica napus cultivar "Westar" was transformed by co-cultivation of seedling pieces with disarmed Agrobacterium tumefaciens strain LBA4404 carrying the the appropriate binary vector.

15           B. napus seeds were sterilized by stirring in 10% Chlorox, 0.1% SDS for thirty min, and then rinsed thoroughly with sterile distilled water. The seeds were germinated on sterile medium containing 30 mM CaCl<sub>2</sub> and 1.5% agar, and grown for six days in the dark at 24°C.

20           Liquid cultures of Agrobacterium for plant transformation were grown overnight at 28°C in Minimal A medium containing 100 mg/L kanamycin. The bacterial cells are pelleted by centrifugation and resuspended at a concentration of 10<sup>8</sup> cells/mL in liquid Murashige and 25 Skoog Minimal Organic medium containing 100 µM acetosyringone.

B. napus seedling hypocotyls were cut into 5 mm segments which were immediately placed into the bacterial suspension. After 30 min, the hypocotyl 30 pieces were removed from the bacterial suspension and placed onto BC-12 callus medium containing 100 µM acetosyringone. The plant tissue and Agrobacteria were co-cultivated for three days at 24°C in dim light.

          The co-cultivation was terminated by transferring 35 the hypocotyl pieces to BC-12 callus medium containing

09210525-12330

200 mg/L carbenicillin to kill the Agrobacteria, and 25 mg/L kanamycin to select for transformed plant cell growth. The seedling pieces were incubated on this medium for three weeks at 24°C under continuous light.

5       After three weeks, the segments were transferred to BS-48 regeneration medium containing 200 mg/L carbenicillin and 25 mg/L kanamycin. Plant tissue was subcultured every two weeks onto fresh selective regeneration medium, under the same culture conditions  
10       described for the callus medium. Putatively transformed calli grow rapidly on regeneration medium; as calli reached a diameter of about 2 mm, they were removed from the hypocotyl pieces and placed on the same medium lacking kanamycin.

15       Shoots began to appear within several weeks after transfer to BS-48 regeneration medium. As soon as the shoots formed discernable stems, they were excised from the calli, transferred to MSV-1A elongation medium, and moved to a 16:8 h day/night photoperiod at 24°C.

20       Once shoots had elongated several internodes, they were cut above the agar surface and the cut ends were dipped in Rootone. Treated shoots were planted directly into wet Metro-Mix 350 soilless potting medium. The pots were covered with plastic bags which were removed when  
25       the plants were clearly growing -- after about 10 days.

      Plants were grown under a 16:8 h day/night photoperiod, with a daytime temperature of 23°C and a nighttime temperature of 17°C. When the primary flowering stem began to elongate, it was covered with a  
30       mesh pollen-containment bag to prevent outcrossing. Self-pollination was facilitated by shaking the plants several times each day. Seeds derived from self-pollinations were harvested about three months after planting.



TABLE 9

Minimal A Bacterial Growth Medium	Brassica Callus Medium BC-12
Dissolve in distilled water:	Per liter:
10.5 g potassium phosphate, dibasic	Murashige and Skoog Minimal
4.5 g potassium phosphate, monobasic	Organic Medium (MS salts, 100
1.0 g ammonium sulfate	mg/L i-inositol, 0.4 mg/L
0.5 g sodium citrate, dihydrate	thiamine; GIBCO #510-3118)
Make up to 979 mLs with distilled water	30 sucrose
Autoclave	18 g mannitol
Add 20 mLs filter-sterilized 10% sucrose	1.0 mg/L 2,4-D
Add 1 mL filter-sterilized 1 M MgSO <sub>4</sub>	3.0 mg/L kinetin
	0.6% agarose
	pH 5.8
Brassica Regeneration Medium BS-48	Brassica Shoot Elongation
Murashige and Skoog Minimal	Medium MSV-1A
Organic Medium Gamborg B5 Vitamins (SIGMA #1019)	Murashige and Skoog Minimal
10 g glucose	Organic Medium Gamborg B5
250 mg xylose	Vitamins
600 mg MES	10 g sucrose
0.4% agarose	0.6% agarose
pH 5.7	pH 5.8
Filter-sterilize and add after autoclaving:	
2.0 mg/L zeatin	
0.1 mg/L IAA	

EXAMPLE 12ANALYSIS OF TRANSGENIC BRASSICA NAPUS PLANTS

- 5 Insertion of the intact antisense transcriptional unit was verified by Southern analysis using transgenic plant leaf tissue as the source of DNA as described in Example 5. Ten micrograms of leaf DNA was digested to completion with a mixture of Bam HI and Sal I

restriction endonucleases and then separated by agarose gel electrophoresis. The separated DNA was transferred to Hybond H<sup>+</sup> membrane and hybridized with radiolabeled insert from pBNSF3-2. An estimate of the number of  
5 copies of the inserted transgene was made by calibrating each Southern blot with standard amounts of pBNSF3-2 corresponding to 1 and 5 copies per genome and comparing intensities of the autoradiographic signal from the standards, the endogenous delta-15 desaturase signals  
10 and the inserted gene signal. To date, 38 independent transformants have been analyzed for presence of the gene and 36 were found to be positive.

The relative content of the 5 most abundant fatty acids in canola seeds was determined either by direct  
15 trans-esterification of individual seeds in 0.5 mL of methanolic H<sub>2</sub>SO<sub>4</sub> (2.5%) or by hexane extraction of bulk seed samples followed by trans-esterification of an aliquot in 0.8 mL of 1% sodium methoxide in methanol. Fatty acid methyl esters were extracted from the  
20 methanolic solutions into hexane after the addition of an equal volume of water.

The relative content of 18:3 fatty acid varies significantly during seed development. To a lesser extent, the ratio of 18:3 to 18:2 varies also. Thus  
25 meaningful data can be obtained only from seeds after maturation and drydown. Additionally, the ratio of 18:3 to total fatty acid content and to 18:0 varies significantly due to environmental factors, primarily temperature. In this circumstance, the most appropriate  
30 controls are the transformed plants which by Southern analysis do not contain the antisense delta-15 transgene. Analysis from the first 5 transformants to reach dry seed are given in Table 10 below. Seeds were harvested using a hand thresher, bulked and a 1.5 g  
35 (about 300 seeds) sample was taken. Seed from each

- transformant was crushed with a mortar and pestel, extracted 4 times with 8 mL hexane at about 50°C. The combined extracts were reduced in volume to 5 mL and two 50 microliter aliquots were taken for esterification as described above. Separation of the fatty acid methyl esters was done by gas-liquid chromatography using an Omegawax 320 column (Supelco Inc., 0.32 mm ID X 30M) run isothermally at 220° and cycled to 260° between each injection.

TABLE 10

Transformant No.	% 18:3	%18:3/18:2	Antisense delta-15 Copy No.
pZCC3FdR-91	6.2	0.39	0
pZCC3FdR-81	5.9	0.33	1
pZCC3FdR-15	6.0	0.38	2
pZCC3FdR-11	5.6	0.34	1
pZCC3FdR-148	8.2	0.40	2

- The differences between the 4 transformed lines and line 92 are very small, however to test the significance of the difference in the 18:3/18:2 ratio between line 81 and 91, 25 individual seeds from each line were transesterified and their fatty acid composition determined.
- The average ratio for line 81 was 0.345 with a coefficient of variation of 11.6% while the average for line 91 was 0.375 with a coefficient of variation of 8.0%. The sample means are significantly different at the 0.01% level using Student's t test.

## EXAMPLE 13

CONSTRUCTION OF VECTORS FOR TRANSFORMATION OF  
GLYCINE MAX FOR REDUCED EXPRESSION OF DELTA-15  
DESATURASES IN DEVELOPING SEEDS

- The antisense G. max plastid delta-15 desaturase cDNA sequence under control of the  $\beta$ -conglycinin promoter was constructed using the vector pCW109A described in Example 10 above. For use in the soybean

transformation system described below, the transcriptional unit was placed in a vector along with an appropriate selectable marker expression system. The starting vector was pML45, which consists of the non-tissue specific and constitutive promoter designated 508D and described in Hershey (WO 9011361) driving expression of the neomycin phosphotransferase gene described in (Beck et al. (1982) Gene 19:327-336) followed by the 3' end of the nopaline synthase gene including nucleotides 848 to 1550 described by (Depicker et al. (1982) J. Appl. Genet. 1:561-574). This transcriptional unit was inserted into the commercial cloning vector pGEM9Z (BRL) and is flanked at the 5' end of the 508D promoter by the restriction sites Sal I, Xba I, Bam HI and Sma I in that order. An additional Sal I site is present at the 3' end of the NOS 3' sequence and the Xba I, Bam HI and Sal I sites are unique.

Removal of the unit[ $\beta$ -conglycinin promoter:cloning region:phaseolin 3' end] from pCW109A by digestion with Hind III, blunting the ends and isolating the 1.8 kb fragment afforded the expression cassette pCST by ligating the above isolated fragment into the Sma I site of pML45. A clone with the  $\beta$ -conglycinin promoter in the same orientation as the 508D promoter were chosen by digestion with Xba I. The correct orientation releases a 700 bp fragment. This vector cassette was called pCST.

The 2.2 kb insert encoding the soybean, plastid delta-15 desaturase was subcloned from the plasmid pXF1 by digestion with HinP I to remove about 1 kb of unrelated cDNA. HinP I cuts within the cDNA insert very near the 5' end of the cDNA for the delta-15 desaturase and about 300 bp from the 3' end of that cDNA. The Cla I compatible ends were cloned into Cla I digested pBluescript and a clone with the 5' end of the cDNA

5        The delta-15 encoding sequence was removed from  
pS3Fd1 by digestion with Hinc II and Eco RV, the 2.2 kB  
fragment was gel isolated and cloned into the opened Sma  
I site in pCST1. A clone with the delta-15 sequence in  
the antisense orientation to the  $\beta$ -conglycinin promoter  
10 was selected by digestion with Xba I. The antisense  
construct releases a 400 bp piece and that clone was  
designated pCS3FdST1R.

## TRANSFORMATION OF SOMATIC SOYBEAN EMBRYO CULTURES

Soybean embryogenic suspension cultures were transformed with pCS3dST1R by the method of particle gun bombardment (see Kline et al. (1987) Nature (London) 327:70). A Du Pont Biolistic PDS1000/HE instrument (helium retrofit) was used for these transformations.

To 50 mL of a 60 mg/mL 1 mm gold particle suspension was added (in order); 5  $\mu$ L DNA (1  $\mu$ g/ $\mu$ L), 20  $\mu$ L spermidine (0.1M), and 50  $\mu$ L  $\text{CaCl}_2$  (2.5 M). The particle preparation was agitated for 3 min, spun in a microfuge for 10 sec and the supernatant removed. The DNA-coated particles were then washed once in 400  $\mu$ L 70% ethanol and resuspended in 40  $\mu$ L of anhydrous ethanol. The DNA/particle suspension was sonicated three times

for 1 sec each. Five  $\mu$ L of the DNA-coated gold particles were then loaded on each macro carrier disk.

Approximately 300-400 mg of a four week old suspension culture was placed in an empty 60x15 mm petri dish and the residual liquid removed from the tissue with a pipette. For each transformation experiment, approximately 5-10 plates of tissue were normally bombarded. Membrane rupture pressure was set at 1000 psi and the chamber was evacuated to a vacuum of 28 inches of mercury. The tissue was placed approximately 3.5 inches away from the retaining screen and bombarded three times. Following bombardment, the tissue was placed back into liquid and cultured as described above.

Eleven days post bombardment, the liquid media was exchanged with fresh SB55 containing 50 mg/mL hygromycin. The selective media was refreshed weekly. Seven weeks post bombardment, green, transformed tissue was observed growing from untransformed, necrotic embryogenic clusters. Isolated green tissue was removed and inoculated into individual flasks to generate new, clonally propagated, transformed embryogenic suspension cultures. Thus each new line was treated as independent transformation event. These suspensions can then be maintained as suspensions of embryos clustered in an immature developmental stage through subculture or regenerated into whole plants by maturation and germination of individual somatic embryos.

Transformed embryogenic clusters were removed from liquid culture and placed on a solid agar media (SB103) containing no hormones or antibiotics. Embryos were cultured for eight weeks at 26°C with mixed florescent and incandescent lights on a 16:8 h day/night schedule. During this period, individual embryos were removed from the clusters and analyzed at various stages of embryo

development After eight weeks the embryos become suitable for germination.

TABLE 11

Media:		B5 Vitamin Stock
SB55 and SBP6 Stock Solutions (g/L):		10 g m-inositol
		100 mg nicotinic acid
MS Sulfate 100X Stock		100 mg pyridoxine HCl
MgSO <sub>4</sub> 7H <sub>2</sub> O	37.0	1 g thiamine
MnSO <sub>4</sub> H <sub>2</sub> O	1.69	SB55 (per Liter)
ZnSO <sub>4</sub> 7H <sub>2</sub> O	0.86	10 mL each MS stocks
CuSO <sub>4</sub> 5H <sub>2</sub> O	0.0025	1 mL B5 Vitamin stock
MS Halides 100X Stock		0.8 g NH <sub>4</sub> NO <sub>3</sub>
CaCl <sub>2</sub> 2H <sub>2</sub> O	44.0	3.033 g KNO <sub>3</sub>
KI	0.083	1 mL 2,4-D (10mg/mL stock)
CoCl <sub>2</sub> 6H <sub>2</sub> O	0.00125	60 g sucrose
KH <sub>2</sub> PO <sub>4</sub>	17.0	0.667 g asparagine
H <sub>3</sub> BO <sub>3</sub>	0.62	pH 5.7
Na <sub>2</sub> MoO <sub>4</sub> 2H <sub>2</sub> O	0.025	For SBP6- substitute 0.5 mL 2,4-D
MS FeEDTA 100X Stock		SB103 (per Liter)
Na <sub>2</sub> EDTA	3.724	MS Salts
FeSO <sub>4</sub> 7H <sub>2</sub> O	2.784	6% maltose
		750 mg MgCl <sub>2</sub>
		0.2% Gelrite
		pH 5.7

EXAMPLE 15ANALYSIS OF TRANSGENIC GLYCINE MAX PLANTS

- 5 While in the globular embryo state in liquid culture as described in Example 14, somatic soybean embryos contain very low amounts of triacylglycerol or storage proteins typical of maturing, zygotic soybean embryos. At this developmental stage, the ratio of
- 10 total triacylglyceride to total polar lipid (phospholipids and glycolipid) is about 1:4, as is typical of zygotic soybean embryos at the developmental

stage from which the somatic embryo culture was initiated. At the globular stage as well, the mRNAs for the prominent seed proteins ( $\alpha'$  subunit of  $\beta$ -conglycinin, Kunitz Trypsin Inhibitor III and Soybean Seed Lectin) are essentially absent. Upon transfer to hormone free media to allow differentiation to the maturing somatic embryo state as described in Example 14, triacylglycerol becomes the most abundant lipid class. As well, mRNAs for  $\alpha'$ -subunit of  $\beta$ -conglycinin, Kunitz Trypsin Inhibitor III and Soybean Seed Lectin become very abundant messages in the total mRNA population. In these respects the somatic soybean embryo system behaves very similarly to maturing zygotic soybean embryos *in vivo*, and is therefore a good and rapid model system for analyzing the phenotypic effects of modifying the expression of genes in the fatty acid biosynthesis pathway. Similar somatic embryo culture systems have been documented and used in another oilseed crop, rapeseed (Taylor et al. (1990) *Planta* 181:18-26). Fatty acid analysis was performed as described in Example 12 using single embryos as the tissue source. A number of embryos from line 2872 (control tissue transformed with pCST) and lines 299,303,306 and 307 (line 2872 transformed with plasmid pCS3FdST1R) were analyzed for fatty acid content. The relative fatty-acid composition of embryos taken from tissue transformed with pCS3FdST1R was compared with control tissue, transformed with pCST. The results of this analysis are shown in Table 12.

TABLE 12

Line	Embryo	16:0	18:0	18:1	18:2	18:3
2872	1	17.7	4.1	11.3	52.8	14.1
	2	17.3	4.3	10.9	49.5	18.0
	3	16.1	4.1	13.8	48.2	17.3
	4	17.5	3.6	11.7	52.0	14.1





<u>Line</u>	<u>Embryo</u>	<u>16:0</u>	<u>18:0</u>	<u>18:1</u>	<u>18:2</u>	<u>18:3</u>
307-1-3	1	14.8	3.1	9.4	60.5	12.2
	2	18.0	3.0	5.3	56.2	15.2
	3	18.0	3.4	2.5	58.6	15.4
307-1-4	1	15.0	2.7	13.8	61.7	6.9
	2	15.9	2.7	9.8	62.0	9.6
	3	14.6	3.2	13.4	61.4	6.7
307-1-5	1	15.9	3.5	7.6	61.7	11.2
	2	14.6	3.5	10.0	61.3	10.6
	3	18.7	2.6	6.8	53.0	19.0
307-1-7	1	15.3	3.5	12.5	60.3	8.5
	2	16.2	2.2	13.9	57.1	10.6
	3	14.9	3.1	12.2	58.0	11.8
307-1-9	1	16.4	2.9	23.2	47.9	9.6
	2	19.6	0.0	20.4	51.3	8.8
	3	16.8	3.3	24.6	49.6	5.7
307-1-11	1	18.1	3.6	5.7	52.9	19.7
	2	14.7	3.7	9.9	58.7	13.0
	3	15.1	3.7	11.3	55.8	14.1

The average 18:3 content of control embryos was 14.5% with a range from 11.1% to 18.0%. The average 18:3 content of transformed embryos was 11.5% with a range of 6.3% to 19.9%. Almost 80% of the transformed embryos (38/48) had an 18:3 content below that of the control mean. About 44% had an 18:3 content less than the lowest observed control value and 12.5% had an 18:3 content less than half of the control mean value (i.e., less than 7.5%). The lowest 18:3 content observed in transformed tissue was 6.3% (299-1-3, 307-1-2 #1) compared with the control low of 11.1%. In all cases in transformed tissue, a decrease in 18:3 content was reflected by an equivalent increase in 18:2 content indicating that the desaturation of 18:2 to 18:3 had

been reduced. The relative content of the the other fatty acids remained unchanged.

Southern analysis for the presence of the intact, introduced antisense construction was performed, as described in Example 12 using Bam HI cut gDNA, on a number of the transformed lines listed below using groups of embryos from a single transformation event. The approximate intact antisense copy number was estimated from the number and intensity of hybridizing bands on the autoradiograms and is shown in Table 13.

TABLE 13

<u>Line No.</u>	<u>Antisense copy No.</u>	<u>18:3 (low)</u>	<u>18:3 (average)</u>	<u>18:2/18:3 ratio</u>
2872	0	11.1	14.5	3.6
303-7/1	1	11.4	12.6	4.7
307-1/2	3	12.2	16.0	3.5
306-4/8	3	10.8	12.2	4.3
307-1/7	4	8.5	10.3	5.7
306-4/5	6	7.4	10.4	5.8
307-1/1	6	6.3	9.6	6.3
299-15/1	7	8.1	9.7	6.1
307-1/4	8	6.7	7.7	8.0

There was a reasonable correlation between intact antisense copy number and 18:3 content, an increase in copy number correlating with a decreased 18:3 content and a consequent increase in the 18:2/18:3 ratio. The average 18:2/18:3 ratio of line 307-1/4, which had at least 8 copies of the antisense cDNA, was more than twice that of the control.

## SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANTS: Browse, John, Kinney, Anthony J.,  
Pierce, John, Wierzbicki, Anna M.,  
Yadav, Narendra S., Perez-Grau, Luis
- (ii) TITLE OF INVENTION: Fatty Acid Desaturase Genes  
from Plants
- (iii) NUMBER OF SEQUENCES: 32
- (iv) CORRESPONDENCE ADDRESS:
- (A) ADDRESSEE: E. I. du Pont de Nemours and Company  
(B) STREET: 1007 Market Street  
(C) CITY: Wilmington  
(D) STATE: Delaware  
(E) COUNTRY: U.S.A.  
(F) ZIP: 19898
- (v) COMPUTER READABLE FORM:
- (A) MEDIUM TYPE: Floppy disk  
(B) COMPUTER: Macintosh  
(C) OPERATING SYSTEM: Macintosh System, 6.0  
(D) SOFTWARE: Microsoft Word, 4.0
- (vi) CURRENT APPLICATION DATA:
- (A) APPLICATION NUMBER:  
(B) FILING DATE:  
(C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:
- (A) APPLICATION NUMBER: 07/804,259  
(B) FILING DATE: 4 DECEMBER 1991
- (viii) ATTORNEY/AGENT INFORMATION:
- (A) NAME: Floyd, Linda A.  
(B) REGISTRATION NUMBER: 33,692  
(C) REFERENCE/DOCKET NUMBER: BB-1036-A
- (ix) TELECOMMUNICATION INFORMATION:
- (A) TELEPHONE: (302) 992-4929  
(B) TELEFAX: (302) 892-7949  
(C) TELEX: 835420

CTCTCTCTCT	CTCTCTTCTC	TCITTTCTCTC	CCCTCTCTCT	CGGCG	ATG	GTT	GTT		54							
					Met	Val	Val									
					1											
GCT	ATG	GAC	CAA	CGC	ACC	AAT	GTG	AAC	GGA	GAT	CCC	GGC	GCC	GGA	GAC	102
Ala	Met	Asp	Gln	Arg	Thr	Asn	Val	Asn	Gly	Asp	Pro	Gly	Ala	Gly	Asp	
	5					10					15					
CGG	AAG	AAA	GAA	GAA	AGG	TTT	GAT	CCG	AGT	GCA	CAA	CCA	CCG	TTC	AAG	150
Arg	Lys	Lys	Glu	Glu	Arg	Phe	Asp	Pro	Ser	Ala	Gln	Pro	Pro	Phe	Lys	
20					25					30					35	
ATC	GGA	GAT	ATA	AGG	GCG	GCG	ATT	CCT	AAG	CAC	TGT	TGG	GTT	AAG	AGT	198
Ile	Gly	Asp	Ile	Arg	Ala	Ala	Ile	Pro	Lys	His	Cys	Trp	Val	Lys	Ser	
				40					45				50			
CCT	TTG	AGA	TCA	ATG	AGT	TAC	GTC	AGA	GAC	ATT	ATC	GCC	GTC	GCG		246
Pro	Leu	Arg	Ser	Met	Ser	Tyr	Val	Val	Arg	Asp	Ile	Ile	Ala	Val	Ala	
			55					60					65			
GCT	TTG	GCC	ATC	GCT	GCC	GTG	TAT	GTT	GAT	AGC	TGG	TTC	CTT	TGG	CCT	294
Ala	Leu	Ala	Ile	Ala	Ala	Val	Tyr	Val	Asp	Ser	Trp	Phe	Leu	Trp	Pro	
		70				75						80				
CTT	TAT	TGG	GCC	GCC	CAA	GGA	ACA	CTT	TTC	TGG	GCC	ATC	TTT	GTT	CTC	342
Leu	Tyr	Trp	Ala	Ala	Gln	Gly	Thr	Leu	Phe	Trp	Ala	Ile	Phe	Val	Leu	
	85					90					95					
GGC	CAC	GAC	TGT	GGA	CAT	GGG	AGT	TTC	TCA	GAC	ATT	CCT	CTA	CTG	AAT	390
Gly	His	Asp	Cys	Gly	His	Gly	Ser	Phe	Ser	Asp	Ile	Pro	Leu	Leu	Asn	
100					105					110					115	

AGT GTG GTT GGT CAC ATT CTT CAT TCT TTC ATC CTC GTT CCT TAC CAT Ser Val Val Gly His Ile Leu His Ser Phe Ile Leu Val Pro Tyr His 120 125 130	438
GGT TGG AGA ATA AGC CAC CGG ACA CAC CAG AAC CAT GGC CAT GTT Gly Trp Arg Ile Ser His Arg Thr His His Gln Asn His Gly His Val 135 140 145	486
GAA AAC GAC GAG TCA TGG GTT CCG TTA CCA GAA AGS GTG TAC AAG AAA Glu Asn Asp Glu Ser Trp Val Pro Leu Pro Glu Arg Val Tyr Lys Lys 150 155 160	534
TTG CCC CAC AGT ACT CGG ATG CTC AGA TAC ACT GTC CCT CTC CCC ATG Leu Pro His Ser Thr Arg Met Leu Arg Tyr Thr Val Pro Leu Pro Met 165 170 175	582
CTC GCA TAT CCT CTC TAT TTG TGC TAC AGA AGT CCT GGA AAA GAA GGA Leu Ala Tyr Pro Leu Tyr Leu Cys Tyr Arg Ser Pro Gly Lys Glu Gly 180 185 190 195	630
TCA CAT TTT AAC CCA TAC AGT AGT TTA TTT GCT CCA AGC GAG AGA AAG Ser His Phe Asn Pro Tyr Ser Ser Leu Phe Ala Pro Ser Glu Arg Lys 200 205 210	678
CTT ATT GCA ACT TCA ACT ACT TGT TGG TCC ATA ATG TTC GTC AGT CTT Leu Ile Ala Thr Ser Thr Thr Cys Trp Ser Ile Met Phe Val Ser Leu 215 220 225	726
ATC GCT CTA TCT TTC GTC TTC GGT CCA CTC GCG GTT CTT AAA GTC TAC Ile Ala Leu Ser Phe Val Phe Gly Pro Leu Ala Val Leu Lys Val Tyr 230 235 240	774
GGT GTA CCG TAC ATT ATC TTT GTG ATG TGG TTG GAT GCT GTC ACG TAT Gly Val Pro Tyr Ile Ile Phe Val Met Trp Leu Asp Ala Val Thr Tyr 245 250 255	822
TTG CAT CAT CAT GGT CAC GAT GAG AAG TTG CCT TGG TAT AGA GGC AAG Leu His His His Gly His Asp Glu Lys Leu Pro Trp Tyr Arg Gly Lys 260 265 270 275	870
GAA TGG AGT TAT CTA CGT GGA GGA TTA ACA ACA ATT GAT AGA GAT TAC Glu Trp Ser Tyr Leu Arg Gly Gly Leu Thr Thr Ile Asp Arg Asp Tyr 280 285 290	918
GGA ATC TTT AAC AAC ATT CAT CAC GAC ATT GGA ACT CAC GTG ATC CAT Gly Ile Phe Asn Asn Ile His His Asp Ile Gly Thr His Val Ile His 295 300 305	966
CAT CTC TTC CCA CAA ATC CCT CAC TAT CAC TTG GTC GAC GCC ACG AAA His Leu Phe Pro Gln Ile Pro His Tyr His Leu Val Asp Ala Thr Lys 310 315 320	1014
GCA GCT AAA CAT GTG TTG GGA AGA TAC TAC AGA GAA CCA AAG ACG TCA Ala Ala Lys His Val Leu Gly Arg Tyr Tyr Arg Glu Pro Lys Thr Ser 325 330 335	1062

(2) INFORMATION FOR SEQ ID NO:2:

(A) LENGTH: 386 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met	Val	Val	Ala	Met	Asp	Gln	Arg	Thr	Asn	Val	Asn	Gly	Asp	Pro	Gly
1				5					10					15	
Ala	Gly	Asp	Arg	Lys	Lys	Glu	Glu	Arg	Phe	Asp	Pro	Ser	Ala	Gln	Pro
			20					25					30		
Pro	Phe	Lys	Ile	Gly	Asp	Ile	Arg	Ala	Ala	Ile	Pro	Lys	His	Cys	Trp
		35					40					45			
Val	Lys	Ser	Pro	Leu	Arg	Ser	Met	Ser	Tyr	Val	Val	Arg	Asp	Ile	Ile
	50					55					60				
Ala	Val	Ala	Ala	Leu	Ala	Ile	Ala	Ala	Val	Tyr	Val	Asp	Ser	Trp	Phe
	65				70					75					80
Leu	Trp	Pro	Leu	Tyr	Trp	Ala	Ala	Gln	Gly	Thr	Leu	Phe	Trp	Ala	Ile
			85						90					95	
Phe	Val	Leu	Gly	His	Asp	Cys	Gly	His	Gly	Ser	Phe	Ser	Asp	Ile	Pro
		100						105					110		
Leu	Leu	Asn	Ser	Val	Val	Gly	His	Ile	Leu	His	Ser	Phe	Ile	Leu	Val
	115						120					125			
Pro	Tyr	His	Gly	Trp	Arg	Ile	Ser	His	Arg	Thr	His	His	Gln	Asn	His
	130					135					140				

Gly His Val Glu Asn Asp Glu Ser Trp Val Pro Leu Pro Glu Arg Val  
 145 150 155 160  
 Tyr Lys Lys Leu Pro His Ser Thr Arg Met Leu Arg Tyr Thr Val Pro  
 165 170 175  
 Leu Pro Met Leu Ala Tyr Pro Leu Tyr Leu Cys Tyr Arg Ser Pro Gly  
 180 185 190  
 Lys Glu Gly Ser His Phe Asn Pro Tyr Ser Ser Leu Phe Ala Pro Ser  
 195 200 205  
 Glu Arg Lys Leu Ile Ala Thr Ser Thr Thr Cys Trp Ser Ile Met Phe  
 210 215 220  
 Val Ser Leu Ile Ala Leu Ser Phe Val Phe Gly Pro Leu Ala Val Leu  
 225 230 235 240  
 Lys Val Tyr Gly Val Pro Tyr Ile Ile Phe Val Met Trp Leu Asp Ala  
 245 250 255  
 Val Thr Tyr Leu His His His Gly His Asp Glu Lys Leu Pro Trp Tyr  
 260 265 270  
 Arg Gly Lys Glu Trp Ser Tyr Leu Arg Gly Gly Leu Thr Thr Ile Asp  
 275 280 285  
 Arg Asp Tyr Gly Ile Phe Asn Asn Ile His His Asp Ile Gly Thr His  
 290 295 300  
 Val Ile His His Leu Phe Pro Gln Ile Pro His Tyr His Leu Val Asp  
 305 310 315 320  
 Ala Thr Lys Ala Ala Lys His Val Leu Gly Arg Tyr Tyr Arg Glu Pro  
 325 330 335  
 Lys Thr Ser Gly Ala Ile Pro Ile His Leu Val Glu Ser Leu Val Ala  
 340 345 350  
 Ser Ile Lys Lys Asp His Tyr Val Ser Asp Thr Gly Asp Ile Val Phe  
 355 360 365  
 Tyr Glu Thr Asp Pro Asp Leu Tyr Val Tyr Ala Ser Asp Lys Ser Lys  
 370 375 380  
 Ile Asn  
 385

(2) INFORMATION FOR SEQ ID NO:3:

(1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 255 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear



(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

(2) INFORMATION FOR SEQ ID NO:4:

```

(i) SEQUENCE CHARACTERISTICS:
      (A) LENGTH: 1525 base pairs
      (B) TYPE: nucleic acid
      (C) STRANDEDNESS: single
      (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(vi) ORIGINAL SOURCE:
      (A) ORGANISM: Arabidopsis thaliana

(vii) IMMEDIATE SOURCE:
      (B) CLONE: pACF2-2

(ix) FEATURE:
      (A) NAME/KEY: CDS
      (B) LOCATION: 10..1350

```

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

CAAGTTCTCA	ATG	GCG	AAC	TTG	GTC	TTA	TCA	GAA	TGT	GGT	ATA	CGA	CCT		48	
	Met	Ala	Asn	Leu	Val	Leu	Ser	Glu	Cys	Gly	Ile	Arg	Pro			
	1				5					10						
CTC	CCC	AGA	ATC	TAC	ACA	ACA	CCC	AGA	TCC	AAT	TTC	CTC	TCC	AAC	AAC	96
Leu	Pro	Arg	Ile	Tyr	Thr	Thr	Pro	Arg	Ser	Asn	Phe	Leu	Ser	Asn	Asn	
	15					20					25					
AAC	AAA	TTC	AGA	CCA	TCA	CTT	TCT	TCT	TCT	TCT	TAC	AAA	ACA	TCA	TCA	144
Asn	Lys	Phe	Arg	Pro	Ser	Leu	Ser	Ser	Ser	Ser	Tyr	Lys	Thr	Ser	Ser	
	30				35					40					45	
TCT	CCT	CTG	TCT	TTT	GGT	CTG	AAT	TCA	CGA	GAT	GGG	TTC	ACG	AGG	AAT	192
Ser	Pro	Leu	Ser	Phe	Gly	Leu	Asn	Ser	Arg	Asp	Gly	Phe	Thr	Arg	Asn	
				50					55					60		
TGG	GCG	TTG	AAT	GTG	AGC	ACA	CCA	TTA	ACG	ACA	CCA	ATA	TTT	GAG	GAG	240
Trp	Ala	Leu	Asn	Val	Ser	Thr	Pro	Leu	Thr	Thr	Pro	Ile	Phe	Glu	Glu	
			65					70					75			
TCT	CCA	TTG	GAG	GAA	GAT	AAT	AAA	CAG	AGA	TTC	GAT	CCA	GGT	GCG	CCT	288
Ser	Pro	Leu	Glu	Glu	Asp	Asn	Lys	Gln	Arg	Phe	Asp	Pro	Gly	Ala	Pro	
		80					85					90				
CCT	CCG	TTC	AAT	TTA	GCT	GAT	ATT	AGA	GCA	GCT	ATA	CCT	AAG	CAT	TGT	336
Pro	Pro	Phe	Asn	Leu	Ala	Asp	Ile	Arg	Ala	Ala	Ile	Pro	Lys	His	Cys	
		95				100					105					
TGG	GTT	AAG	AAT	CCA	TGG	AAG	TCT	TTG	AGT	TAT	GTC	GTC	AGA	GAC	GTC	384
Trp	Val	Lys	Asn	Pro	Trp	Lys	Ser	Leu	Ser	Tyr	Val	Val	Arg	Asp	Val	
	110				115					120				125		
GCT	ATC	GTC	TTT	GCA	TTG	GCT	GCT	GGA	GCT	GCT	TAC	CTC	AAC	AAT	TGG	432
Ala	Ile	Val	Phe	Ala	Leu	Ala	Ala	Gly	Ala	Ala	Tyr	Leu	Asn	Asn	Trp	
				130				135					140			
ATT	GTT	TGG	CCT	CTC	TAT	TGG	CTC	GCT	CAA	GGA	ACC	ATG	TTT	TGG	GCT	480
Ile	Val	Trp	Pro	Leu	Tyr	Trp	Leu	Ala	Gln	Gly	Thr	Met	Phe	Trp	Ala	
			145				150						155			
CTC	TTT	GTT	CTT	GGT	CAT	GAC	TGT	GGA	CAT	GGT	AGT	TTC	TCA	AAT	GAT	528
Leu	Phe	Val	Leu	Gly	His	Asp	Cys	Gly	His	Gly	Ser	Phe	Ser	Asn	Asp	
		160					165					170				
CCG	AAG	TTG	AAC	AGT	GTG	GTC	GGT	CAT	CTT	CTT	CAT	TCC	TCA	ATT	CTG	576
Pro	Lys	Leu	Asn	Ser	Val	Val	Gly	His	Leu	Leu	His	Ser	Ser	Ile	Leu	
		175				180					185					
GTC	CCA	TAC	CAT	GGC	TGG	AGA	ATT	AGT	CAC	AGA	ACT	CAC	CAC	CAG	AAC	624
Val	Pro	Tyr	His	Gly	Trp	Arg	Ile	Ser	His	Arg	Thr	His	His	Gln	Asn	
	190				195					200				205		
CAT	GGA	CAT	GTT	GAG	AAT	GAC	GAA	TCT	TGG	CAT	CCT					

CCT Ile	TAC Tyr	AAT Asn	ACT Thr	TTG Leu	GAC Asp	AAG Lys	CCG Pro	ACT Thr	AGA Arg	TTC Phe	TTT Phe	AGA Arg	TTT Phe	ACA Thr	CTG Leu	720
CCT Pro	CTC Leu	GTG Val	ATG Met	CTT Leu	GCA Ala	TAC Tyr	CCT Pro	TTC Phe	TAC Tyr	TTG Leu	TGG Trp	GCT Ala	CGA Arg	AGT Ser	CCG Pro	768
GGG Gly	AAA Lys	AAG Lys	GGT Gly	TCT Ser	CAT His	TAC Tyr	CAT His	CCA Pro	GAC Asp	AGT Ser	GAC Asp	TTG Leu	TTC Phe	CTC Leu	CCT Pro	816
AAA 270	GAG Glu	AGA Arg	AAG Lys	GAT Asp	GTC Val	CTC Leu	ACT Thr	TCT Ser	ACT Thr	GCT Ala	TGT Cys	TGG Trp	ACT Thr	GCA Ala	ATG Met	864
GCT Ala	GCT Ala	CTG Leu	CTT Leu	GTT Val	TGT Cys	CTC Leu	AAC Asn	TTC Phe	ACA Thr	ATC Ile	GGT Gly	CCA Pro	ATT Ile	CAA Gln	ATG Met	912
CTC Leu	AAA Lys	CTT Leu	TAT Thr	GGA Gly	ATT Ile	CCT Pro	TAC Tyr	TGG Trp	ATA Ile	AAT Asn	GTA Val	ATG Met	TGG Trp	TTG Leu	GAC Asp	960
TTT Phe	GTG Val	ACT Thr	TAC Thr	CTG Leu	CAT His	CAC His	CAT His	GGT Gly	CAT His	GAA Glu	GAT Asp	AAG Lys	CTT Leu	CCT Pro	TGG Trp	1008
TAC Tyr	CGT Arg	GGC Gly	AAG Lys	GAG Glu	TGG Trp	AGT Ser	TAC Tyr	CTG Leu	AGA Arg	GGA Gly	GGA Gly	CTT Leu	ACA Thr	ACA Thr	TTG Leu	1056
GAT Asp	CGT Arg	GAC Asp	TAC Asp	GGA Gly	TTG Leu	ATC Ile	AAT Asn	AAC Asn	ATC Ile	CAT His	CAT His	GAT Asp	ATT Ile	GGA Gly	ACT Thr	1104
CAT His	GTG Val	ATA Ile	CAT His	CAT His	CTT Leu	TTC Phe	CCG Pro	CAG Gln	ATC Ile	CCA Pro	CAT His	TAT Tyr	CAT His	CTA His	GTA Val	1152
GAA Glu	GCA Ala	ACA Thr	GAA Ala	GCA Ala	GCT Ala	AAA Lys	CCA Pro	GTA Pro	TTA Leu	GGG Gly	AAG Lys	TAT Tyr	TAC Tyr	AGG Arg	GAG Glu	1200
CCT Pro	GAT Asp	AAG Lys	TCT Ser	GGA Gly	CCG Pro	TTG Leu	CCA Pro	TTA Leu	CAT His	TTA Leu	CTG Leu	GAA Glu	ATT Ile	CTA Leu	GCG Al	1248
AAA Lys	AGT Ser	ATA Ile	AAA Lys	GAA Glu	GAT Asp	CAT His	TAC Tyr	GTG Val	AGC Ser	GAC Asp	GAA Glu	GGA Gly	GAA Glu	GTT Val	GTA Val	1296
TAC 430	TAT Tyr	AAA Lys	GCA Ala	GAT Asp	CCA Pro	AAT Asn	CTC Leu	TAT Tyr	GGA Glu	GAG Val	GTC Lys	AAA Lys	GTA Val	AGA Arg	GCA Ala	1344
GAT Asp	TGAAATGAAG CAGGCTTGAG ATTGAAGTTT TTTCTATTTC AGACCAGCTG															1397

ATTTTGTGCT TACTGTATCA ATTATTGTG TCACCCACCA GAGAGTTAGT ATCTCTGAAT 1457  
 ACGATCGATC AGATGGAAAC AACAAATTG TTTGCGATAC TGAAGCTATA TATACCATAC 1517  
 ATTGCATT 1525

## (2) INFORMATION FOR SEQ ID NO:5:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 446 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Met Ala Asn Leu Val Leu Ser Glu Cys Gly Ile Arg Pro Leu Pro Arg  
 1 5 10 15  
 Ile Tyr Thr Thr Pro Arg Ser Asn Phe Leu Ser Asn Asn Asn Lys Phe  
 20 25 30  
 Arg Pro Ser Leu Ser Ser Ser Ser Tyr Lys Thr Ser Ser Ser Pro Leu  
 35 40 45  
 Ser Phe Gly Leu Asn Ser Arg Asp Gly Phe Thr Arg Asn Trp Ala Leu  
 50 55 60  
 Asn Val Ser Thr Pro Leu Thr Thr Pro Ile Phe Glu Glu Ser Pro Leu  
 65 70 75 80  
 Glu Glu Asp Asn Lys Gln Arg Phe Asp Pro Gly Ala Pro Pro Pro Phe  
 85 90 95  
 Asn Leu Ala Asp Ile Arg Ala Ala Ile Pro Lys His Cys Trp Val Lys  
 100 105 110  
 Asn Pro Trp Lys Ser Leu Ser Tyr Val Val Arg Asp Val Ala Ile Val  
 115 120 125  
 Phe Ala Leu Ala Ala Gly Ala Ala Tyr Leu Asn Asn Trp Ile Val Trp  
 130 135 140  
 Pro Leu Tyr Trp Leu Ala Gln Gly Thr Met Phe Trp Ala Leu Phe Val  
 145 150 155 160  
 Leu Gly His Asp Cys Gly His Gly Ser Phe Ser Asn Asp Pro Lys Leu  
 165 170 175  
 Asn Ser Val Val Gly His Leu Leu His Ser Ser Ile Leu Val Pro Tyr  
 180 185 190  
 His Gly Trp Arg Ile Ser His Arg Thr His His Gln Asn His Gly His  
 195 200 205

Val Glu Asn Asp Glu Ser Trp His Pro Met Ser Glu Lys Ile Tyr Asn  
 210 215 220  
 Thr Leu Asp Lys Pro Thr Arg Phe Phe Arg Phe Thr Leu Pro Leu Val  
 225 230 235 240  
 Met Leu Ala Tyr Pro Phe Tyr Leu Trp Ala Arg Ser Pro Gly Lys Lys  
 245 250 255  
 Gly Ser His Tyr His Pro Asp Ser Asp Leu Phe Leu Pro Lys Glu Arg  
 260 265 270  
 Lys Asp Val Leu Thr Ser Thr Ala Cys Trp Thr Ala Met Ala Ala Leu  
 275 280 285  
 Leu Val Cys Leu Asn Phe Thr Ile Gly Pro Ile Gln Met Leu Lys Leu  
 290 295 300  
 Tyr Gly Ile Pro Tyr Trp Ile Asn Val Met Trp Leu Asp Phe Val Thr  
 305 310 315 320  
 Tyr Leu His His His Gly His Glu Asp Lys Leu Pro Trp Tyr Arg Gly  
 325 330 335  
 Lys Glu Trp Ser Tyr Leu Arg Gly Gly Leu Thr Thr Leu Asp Arg Asp  
 340 345 350  
 Tyr Gly Leu Ile Asn Asn Ile His His Asp Ile Gly Thr His Val Ile  
 355 360 365  
 His His Leu Phe Pro Gln Ile Pro His Tyr His Leu Val Glu Ala Thr  
 370 375 380  
 Glu Ala Ala Lys Pro Val Leu Gly Lys Tyr Tyr Arg Glu Pro Asp Lys  
 385 390 395 400  
 Ser Gly Pro Leu Pro Leu His Leu Leu Glu Ile Leu Ala Lys Ser Ile  
 405 410 415  
 Lys Glu Asp His Tyr Val Ser Asp Glu Gly Glu Val Val Tyr Tyr Lys  
 420 425 430  
 Ala Asp Pro Asn Leu Tyr Gly Glu Val Lys Val Arg Ala Asp  
 435 440 445

## (2) INFORMATION FOR SEQ ID NO:6:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1429 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Brassica napus

(vii) IMMEDIATE SOURCE:

(B) CLONE: pBNSF3-f2

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 79..1212

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

TTCAAATTC	A	GACAATCCCC	TTCTTCTCCC	CGGTTTCGTC	TGAACTCTCG	AAACTGGGCG	60
TTGAATGTAA	CCACACCT	CTA	ACA	GTC	GAC	TCC	111
	Leu	Thr	Val	Asp	Ser	Ser	
	1			5			
GAG	GAA	GAA	CCC	AAA	ACG	CAG	159
Glu	Glu	Glu	Pro	Lys	Thr	Gln	
	15					20	
TTC	AAC	CTA	GCT	GAC	ATC	AGA	
Phe	Asn	Leu	Ala	Asp	Ile	Arg	207
	30					35	
AAG	AAT	CCA	TGG	AAG	TCT	ATG	
Lys	Asn	Pro	Trp	Lys	Ser	Met	255
	45					50	
GTG	TTC	GCA	CTA	GCT	GCT	GGA	
Val	Phe	Ala	Leu	Ala	Ala	Gly	303
	60					65	
TGG	CCT	CTC	TAT	TGG	ATT	GCT	
Trp	Pro	Leu	Tyr	Trp	Ile	Ala	351
						80	
GTT	CTT	GGC	CAT	GAC	TGT	GGA	
Val	Leu	Gly	His	Asp	Cys	Gly	399
						95	
TTG	AAC	AGT	GTG	GTG	GGT	CAC	
Leu	Asn	Ser	Val	Val	Gly	His	447
	110					115	
TAC	CAT	GGC	TGG	AGA	ATT	AGC	
Tyr	His	Gly	Trp	Arg	Ile	Ser	495
	125					130	
CAT	GTT	GAG	AAC	GAT	GAA	TCT	
His	Val	Glu	Asn	Asp	Glu	Ser	543
	140					145	

AAG AGT TTG GAC AAA CCC ACT CGG TTC TTT AGA TTT ACA TTG CCT CTC	591
Lys Ser Leu Asp Lys Pro Thr Arg Phe Phe Arg Phe Thr Leu Pro Leu	
160 165 170	
GTG ATG CTC GCT TAC CCT TTC TAC TTG TGG GCA AGA AGT CCA GGG AAG	639
Val Met Leu Ala Tyr Pro Phe Tyr Leu Trp Ala Arg Ser Pro Gly Lys	
175 180 185	
AAG GGT TCT CAT TAC CAT CCA GAC AGC GAC TTG TTC CTT CCT AAA GAG	687
Lys Gly Ser His Tyr His Pro Asp Ser Asp Leu Phe Leu Pro Lys Glu	
190 195 200	
AGA AAC GAT GTT CTC ACT TCT ACC GCT TGT TGG ACT GCA ATG GCT GTT	735
Arg Asn Asp Val Leu Thr Ser Thr Ala Cys Trp Thr Ala Met Ala Val	
205 210 215	
CTG CTT GTC TGT CTC AAC TTC GTG ATG GGT CCA ATG CAA ATG CTC AAA	783
Leu Leu Val Cys Leu Asn Phe Val Met Gly Pro Met Gln Met Leu Lys	
220 225 230 235	
CTT TAT GTC ATT CCT TAC TGG ATA AAT GTA ATG TGG TTG GAC TTT GTG	831
Leu Tyr Val Ile Pro Tyr Trp Ile Asn Val Met Trp Leu Asp Phe Val	
240 245 250	
ACT TAC CTG CAT CAC CAT GGT CAT GAA GAT AAG CTC CCT TGG TAC CGT	879
Thr Tyr Leu His His His Gly His Glu Asp Lys Leu Pro Trp Tyr Arg	
255 260 265	
GGG AAG GAA TGG AGT TAC TTG AGA GGA GGA CTT ACA ACA TTG GAC CGG	927
Gly Lys Glu Trp Ser Tyr Leu Arg Gly Gly Leu Thr Leu Asp Arg	
270 275 280	
GAC TAC GGA TTG ATC AAC AAC ATC CAT CAC GAC ATT GGA ACT CAT GTG	975
Asp Tyr Gly Leu Ile Asn Asn Ile His His Asp Ile Gly Thr His Val	
285 290 295	
ATA CAT CAT CTT TTC CCT CAG ATC CCA CAT TAT CAT CTA GTA GAA GCA	1023
Ile His His Leu Phe Pro Gln Ile Pro His Tyr His Leu Val Glu Ala	
300 305 310 315	
ACA GAA GCA GCT AAA CCA GTA TTA GGG AAG TAT TAT AGG GAG CCT GAT	1071
Thr Glu Ala Ala Lys Pro Val Leu Gly Lys Tyr Tyr Arg Glu Pro Asp	
320 325 330	
AAG TCT GGA CCT TTG CCA TTA CAT TTA CTG GGA ATC TTA GCA AAA AGT	1119
Lys Ser Gly Pro Leu Pro Leu His Leu Leu Gly Ile Leu Ala Lys Ser	
335 340 345	
ATT AAA GAA GAT CAT TTT GTG AGC GAT GAA GGA GAT GTT GTA TAC TAT	1167
Ile Lys Glu Asp His Phe Val Ser Asp Glu Gly Asp Val Val Tyr Tyr	
350 355 360	
GAA GCA GAC CCT AAT CTC TAT GGA GAG ATC AAG GTA ACA GCA GAG	1212
Glu Ala Asp Pro Asn Leu Tyr Gly Glu Ile Lys Val Thr Ala Glu	
365 370 375	
TGAATGAAG CTGTCAGATT TATCTATTTT TGACCAGCTG ATTTTITTTTG CTTATTAATG	1272

TCAATTCATT GTGTTACCAT TATCTCTGAA TACAATCAGA TGGAAACCCC AACTTTGTTT 1332  
TCAATACTTG AAGCTATATA TATATATATA TATGTAAGAT ACATTGTATT GTCATTAGAT 1392  
TCACCATTCT CAAGGTTCTT ATACAAAAAA AAAAAAA 1429

## (2) INFORMATION FOR SEQ ID NO:7:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 378 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

Leu Thr Val Asp Ser Ser Ser Ser Pro Pro Ile Glu Glu Glu Pro Lys  
1 5 10 15  
Thr Gln Arg Phe Asp Pro Gly Ala Pro Pro Pro Phe Asn Leu Ala Asp  
20 25 30  
Ile Arg Ala Ala Ile Pro Lys His Cys Trp Val Lys Asn Pro Trp Lys  
35 40 45  
Ser Met Ser Tyr Val Val Arg Glu Leu Ala Ile Val Phe Ala Leu Ala  
50 55 60  
Ala Gly Ala Ala Tyr Leu Asn Asn Trp Leu Val Trp Pro Leu Tyr Trp  
65 70 75 80  
Ile Ala Gln Gly Thr Met Phe Trp Ala Leu Phe Val Leu Gly His Asp  
85 90 95  
Cys Gly His Gly Ser Phe Ser Asn Asp Pro Arg Leu Asn Ser Val Val  
100 105 110  
Gly His Leu Leu His Ser Ser Ile Leu Val Pro Tyr His Gly Trp Arg  
115 120 125  
Ile Ser His Arg Thr His His Gln Asn His Gly His Val Glu Asn Asp  
130 135 140  
Glu Ser Trp His Pro Met Ser Glu Lys Ile Tyr Lys Ser Leu Asp Lys  
145 150 155 160  
Pro Thr Arg Phe Phe Arg Phe Thr Leu Pro Leu Val Met Leu Ala Tyr  
165 170 175  
Pro Phe Tyr Leu Trp Ala Arg Ser Pro Gly Lys Lys Gly Ser His Tyr  
180 185 190  
His Pro Asp Ser Asp Leu Phe Leu Pro Lys Glu Arg Asn Asp Val Leu  
195 200 205

00110005.122500



Thr Ser Thr Ala Cys Trp Thr Ala Met Ala Val Leu Leu Val Cys Leu  
 210 215 220  
 Asn Phe Val Met Gly Pro Met Gln Met Leu Lys Leu Tyr Val Ile Pro  
 225 230 235 240  
 Tyr Trp Ile Asn Val Met Trp Leu Asp Phe Val Thr Tyr Leu His His  
 245 250 255  
 His Gly His Glu Asp Lys Leu Pro Trp Tyr Arg Gly Lys Glu Trp Ser  
 260 265 270  
 Tyr Leu Arg Gly Gly Leu Thr Thr Leu Asp Arg Asp Tyr Gly Leu Ile  
 275 280 285  
 Asn Asn Ile His His Asp Ile Gly Thr His Val Ile His His Leu Phe  
 290 295 300  
 Pro Gln Ile Pro His Tyr His Leu Val Glu Ala Thr Glu Ala Ala Lys  
 305 310 315 320  
 Pro Val Leu Gly Lys Tyr Tyr Arg Glu Pro Asp Lys Ser Gly Pro Leu  
 325 330 335  
 Pro Leu His Leu Leu Gly Ile Leu Ala Lys Ser Ile Lys Glu Asp His  
 340 345 350  
 Phe Val Ser Asp Glu Gly Asp Val Val Tyr Tyr Glu Ala Asp Pro Asn  
 355 360 365  
 Leu Tyr Gly Glu Ile Lys Val Thr Ala Glu  
 370 375

## (2) INFORMATION FOR SEQ ID NO:8:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1429 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: cDNA

## (iii) HYPOTHETICAL: NO

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: *Brassica napus*

## (vii) IMMEDIATE SOURCE:

- (B) CLONE: pBNSfd-2

## (ix) FEATURE:

(A) NAME/KEY: CDS  
 (B) LOCATION: 1..1215

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

TTC AAA TTC AGA CAA TCC CCT TCT TCT CCC CGG TTT CGT CTG AAC TCT Phe Lys Phe Arg Gln Ser Pro Ser Ser Pro Arg Phe Arg Leu Asn Ser 1 5 10 15	48
CGA AAC TGG GCG TTG AAT GTA ACC ACA CCT CTA ACA GTC GAC TCC TCA Arg Asn Trp Ala Leu Asn Val Thr Pro Leu Thr Val Asp Ser Ser 20 25 30	96
TCA TCT CCT CCA ATC GAG GAA GAA CCC AAA ACG CAG AGA TTC GAC CCA Ser Ser Pro Pro Ile Glu Glu Glu Pro Lys Thr Gln Arg Phe Asp Pro 35 40 45	144
GGC GCT CCT CCG TTC AAC CTA GCT GAC ATC AGA GCG GCG ATA CCT Gly Ala Pro Pro Pro Phe Asn Leu Ala Asp Ile Arg Ala Ala Ile Pro 50 55 60	192
AAG CAT TGC TGG GTT AAG AAT CCA TGG AAG TCT ATG AGT TAC GTC GTC Lys His Cys Trp Val Lys Asn Pro Trp Lys Ser Met Ser Tyr Val Val 65 70 75 80	240
AGA GAG CTA GCC ATC GTG TTC GCA CTA GCT GCT GGA GCT GCT TAC CTC Arg Glu Leu Ala Ile Val Phe Ala Leu Ala Ala Gly Ala Ala Tyr Leu 85 90 95	288
AAC AAT TGG CTT GTT TGG CCT CTC TAT TGG ATT GCT CAA GGA ACC ATG Asn Asn Trp Leu Val Trp Pro Leu Tyr Trp Ile Ala Gln Gly Thr Met 100 105 110	336
TTC TGG GCT CTC TTT GTT CTT GGC CAT GAC TGT GGA CAT GGA AGC TTC Phe Trp Ala Leu Phe Val Leu Gly His Asp Cys Gly His Gly Ser Phe 115 120 125	384
TCA AAT GAT CCG AGG TTG AAC AGT GTG GTG GGT CAC CTT CTT CAT TCC Ser Asn Asp Pro Arg Leu Asn Ser Val Val Gly His Leu Leu His Ser 130 135 140	432
TCT ATT CTA GTC CCT TAC CAT GGC TGG AGA ATT AGC CAC AGA ACT CAC Ser Ile Leu Val Pro Tyr His Gly Trp Arg Ile Ser His Arg Thr His 145 150 155 160	480
CAC CAG AAC CAT GGA CAT GTT GAG AAC GAT GAA TCT TGG CAT CCT ATG His Gln Asn His Gly His Val Glu Asn Asp Glu Ser Trp His Pro Met 165 170 175	528
TCT GAG AAA ATC TAC AAG AGT TTG GAC AAA CCC ACT CGG TTC TTT AGA Ser Glu Lys Ile Tyr Lys Ser Leu Asp Lys Pro Thr Arg Phe Phe Arg 180 185 190	576
TTT ACA TTG CCT CTC GTG ATG CTC GCT TAC CCT TTC TAC TTG TGG GCA Phe Thr Leu Pro Leu Val Met Leu Ala Tyr Pro Phe Tyr Leu Trp Ala 195 200 205	624

0010035-100000

AGA AGT CCA GGG AAG AAG GGT TCT CAT TAC CAT CCA GAC AGC GAC TTG Arg Ser Pro.Gly Lys Lys Gly Ser His Tyr His Pro Asp Ser Asp Leu 210 215 220	672
TTC CTT CCT AAA GAG AGA AAC GAT GTT CTC ACT TCT ACC GCT TGT TGG Phe Leu Pro Lys Glu Arg Asn Asp Val Leu Thr Ser Thr Ala Cys Trp 225 230 235 240	720
ACT GCA ATG GCT GTT CTG CTT GTC TGT CTC AAC TTC GTG ATG GGT CCA Thr Ala Met Ala Val Leu Leu Val Cys Leu Asn Phe Val Met Gly Pro 245 250 255	768
ATG CAA ATG CTC AAA CTT TAT GTC ATT CCT TAC TGG ATA AAT GTA ATG Met Gln Met Leu Lys Leu Tyr Val Ile Pro Tyr Trp Ile Asn Val Met 260 265 270	816
TGG TTG GAC TTT GTG ACT TAC CTG CAT CAC CAT GGT CAT GAA GAT AAG Trp Leu Asp Phe Val Thr Tyr Leu His His His Gly His Glu Asp Lys 275 280 285	864
CTC CCT TGG TAC CGT GGG AAG GAA TGG AGT TAC TTG AGA GGA GGA CTT Leu Pro Trp Tyr Arg Gly Lys Glu Trp Ser Tyr Leu Arg Gly Gly Leu 290 295 300	912
ACA ACA TTG GAC CGG GAC TAC GGA TTG ATC AAC AAC ATC CAT CAC GAC Thr Thr Leu Asp Arg Asp Tyr Gly Leu Ile Asn Asn Ile His His Asp 305 310 315 320	960
ATT GGA ACT CAT GTG ATA CAT CAT CTT TTC CCT CAG ATC CCA CAT TAT Ile Gly Thr His Val Ile His His Leu Phe Pro Gln Ile Pro His Tyr 325 330 335	1008
CAT CTA GTA GAA GCA ACA GAA GCA GCT AAA CCA GTA TTA GGG AAG TAT His Leu Val Glu Ala Thr Glu Ala Ala Lys Pro Val Leu Gly Lys Tyr 340 345 350	1056
TAT AGG GAG CCT GAT AAG TCT GGA CCT TTG CCA TTA CAT TTA CTG GGA Tyr Arg Glu Pro Asp Lys Ser Gly Pro Leu Pro Leu His Leu Leu Gly 355 360 365	1104
ATC TTA GCA AAA AGT ATT AAA GAA GAT CAT TTT GTG AGC GAT GAA GGA Ile Leu Ala Lys Ser Ile Lys Glu Asp His Phe Val Ser Asp Glu Gly 370 375 380	1152
GAT GTT GTA TAC TAT GAA GCA GAC CCT AAT CTC TAT GGA GAG ATC AAG Asp Val Val Tyr Tyr Glu Ala Asp Pro Asn Leu Tyr Gly Glu Ile Lys 385 390 395 400	1200
GTA ACA GCA GAG TGAATGAAG CTGTCAGATT TATCTATTTT TGACCAGCTG Val Thr Ala Glu 405	1252
ATTTTTTTTG CTTATTAATG TCAATTCATT GTGTTACCAT TATCTGTGAA TACAATCAGA	1312
TGGAAACCCC AACTTTGTTT TCAATACTTG AAGCTATATA TATATATATA TATGTAAGAT	1372
ACATTGTATT GTCATTAGAT TCACCATTCT CAAGGTTCTT ATACAAAAAA AAAAAA	1429

## (2) INFORMATION FOR SEQ ID NO:9:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 404 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

Phe Lys Phe Arg Gln Ser Pro Ser Ser Pro Arg Phe Arg Leu Asn Ser  
 1 5 10 15  
 Arg Asn Trp Ala Leu Asn Val Thr Thr Pro Leu Thr Val Asp Ser Ser  
 20 25 30  
 Ser Ser Pro Pro Ile Glu Glu Glu Pro Lys Thr Gln Arg Phe Asp Pro  
 35 40 45  
 Gly Ala Pro Pro Pro Phe Asn Leu Ala Asp Ile Arg Ala Ala Ile Pro  
 50 55 60  
 Lys His Cys Trp Val Lys Asn Pro Trp Lys Ser Met Ser Tyr Val Val  
 65 70 75 80  
 Arg Glu Leu Ala Ile Val Phe Ala Leu Ala Ala Gly Ala Ala Tyr Leu  
 85 90 95  
 Asn Asn Trp Leu Val Trp Pro Leu Tyr Trp Ile Ala Gln Gly Thr Met  
 100 105 110  
 Phe Trp Ala Leu Phe Val Leu Gly His Asp Cys Gly His Gly Ser Phe  
 115 120 125  
 Ser Asn Asp Pro Arg Leu Asn Ser Val Val Gly His Leu Leu His Ser  
 130 135 140  
 Ser Ile Leu Val Pro Tyr His Gly Trp Arg Ile Ser His Arg Thr His  
 145 150 155 160  
 His Gln Asn His Gly His Val Glu Asn Asp Glu Ser Trp His Pro Met  
 165 170 175  
 Ser Glu Lys Ile Tyr Lys Ser Leu Asp Lys Pro Thr Arg Phe Phe Arg  
 180 185 190  
 Phe Thr Leu Pro Leu Val Met Leu Ala Tyr Pro Phe Tyr Leu Trp Ala  
 195 200 205  
 Arg Ser Pro Gly Lys Lys Gly Ser His Tyr His Pro Asp Ser Asp Leu  
 210 215 220  
 Phe Leu Pro Lys Glu Arg Asn Asp Val Leu Thr Ser Thr Ala Cys Trp  
 225 230 235 240

002105025-10000000

Thr Ala Met Ala Val Leu Leu Val Cys Leu Asn Phe Val Met Gly Pro  
 245 250 255  
 Met Gln Met Leu Lys Leu Tyr Val Ile Pro Tyr Trp Ile Asn Val Met  
 260 265 270  
 Trp Leu Asp Phe Val Thr Tyr Leu His His His Gly His Glu Asp Lys  
 275 280 285  
 Leu Pro Trp Tyr Arg Gly Lys Glu Trp Ser Tyr Leu Arg Gly Gly Leu  
 290 295 300  
 Thr Thr Leu Asp Arg Asp Tyr Gly Leu Ile Asn Asn Ile His His Asp  
 305 310 315 320  
 Ile Gly Thr His Val Ile His His Leu Phe Pro Gln Ile Pro His Tyr  
 325 330 335  
 His Leu Val Glu Ala Thr Glu Ala Ala Lys Pro Val Leu Gly Lys Tyr  
 340 345 350  
 Tyr Arg Glu Pro Asp Lys Ser Gly Pro Leu Pro Leu His Leu Leu Gly  
 355 360 365  
 Ile Leu Ala Lys Ser Ile Lys Glu Asp His Phe Val Ser Asp Glu Gly  
 370 375 380  
 Asp Val Val Tyr Tyr Glu Ala Asp Pro Asn Leu Tyr Gly Glu Ile Lys  
 385 390 395 400  
 Val Thr Ala Glu

## (2) INFORMATION FOR SEQ ID NO:10:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2181 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: cDNA

## (iii) HYPOTHETICAL: NO

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Glycine max

## (vii) IMMEDIATE SOURCE:

- (B) CLONE: pXF1

## (ix) FEATURE:

- (A) NAME/KEY: CDS

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:



GGT Trp	GCT Ala	CTC Leu	TTT Phe	GTG Val	CTT Leu	GGA Gly	CAT His	GAT Asp	TGT Cys	GGC Gly	CAT His	GGA Gly	AGC Ser	TTT Phe	TCA Ser	1178
95																
GAT Asp	AGC Ser	CCT Pro	TTG Leu	CTG Leu	AAT Asn	AGC Ser	CTG Leu	GTG Val	GGA Gly	CAC His	ATC Ile	TTG Leu	CAT His	TCC Ser	TCA Ser	1226
110																
ATT Ile	CTT Leu	GTG Val	CCA Pro	TAC Tyr	CAT His	GGA Gly	TGG Trp	AGA Arg	ATT Ile	AGC Ser	CAC His	AGA Arg	ACT Thr	CAC His	CAT His	1274
125																
CAA Gln	AAC Asn	CAT His	GGA Gly	CAC His	ATT Ile	GAG Glu	AAG Lys	GAT Asp	GAG Glu	TCA Ser	TGG Trp	GTT Val	CCA Pro	TTA Leu	ACA Thr	1322
145																
GAG Glu	AAG Lys	ATT Ile	TAC Tyr	AAG Lys	AAT Asn	CTA Leu	GAC Asp	AGC Ser	ATG Met	ACA Thr	AGA Arg	CTC Leu	ATT Ile	AGA Arg	TTC Phe	1370
160																
ACT Thr	GTG Val	CCA Pro	TTT Phe	CCA Pro	TTG Leu	TTT Phe	GTG Val	TAT Tyr	CCA Pro	ATT Ile	TAT Tyr	TTG Phe	TTT Phe	TCA Ser	AGA Arg	1418
175																
AGC Ser	CCC Pro	GGA Gly	AAG Lys	GAA Glu	GGC Gly	TCT Ser	CAC His	TTC Phe	AAT Asn	CCC Pro	TAC Tyr	AGC Ser	AAT Asn	CTG Leu	TTC Phe	1466
190																
CCA Pro	CCC Pro	AGT Ser	GAG Glu	AGA Arg	AAA Lys	GGA Gly	ATA Ile	GCA Ala	ATA Ile	TCA Ser	ACA Thr	CTG Leu	TGT Cys	TGG Trp	GCT Ala	1514
205																
ACC Thr	ATG Met	TTT Phe	TCT Ser	CTG Leu	CTT Leu	ATC Ile	TAT Tyr	CTC Leu	TCA Ser	TTC Phe	ATA Ile	ACT Thr	AGT Ser	CCA Pro	CTT Leu	1562
225																
CTA Leu	GTG Val	CTC Leu	AAG Lys	CTC Leu	TAT Tyr	GGA Gly	ATT Ile	CCA Pro	TAT Tyr	TGG Trp	ATA Ile	TTT Phe	GTT Val	ATG Met	TGG Trp	1610
240																
CTG Leu	GAC Asp	TTT Phe	GTC Val	ACA Thr	TAC Tyr	TTG Leu	CAT His	CAC His	CAT His	GGT Gly	CAC His	CAC His	CAG Gln	AAA Lys	CTG Leu	1658
255																
CCT Pro	TGG Trp	TAC Tyr	CGC Arg	GGC Gly	AAG Lys	GAA Glu	TGG Trp	AGT Ser	TAT Tyr	TTA Leu	AGA Gly	GGT Gly	GGC Gly	CTC Leu	ACC Thr	1706
270																
ACT Thr	GTG Val	GAT Asp	CGT Arg	GAC Asp	TAT Tyr	GGT Gly	TGG Trp	ATC Ile	TAT Tyr	AAC Asn	ATT Ile	CAC His	CAT His	GAC Asp	ATT Ile	1754
285																
GGC Gly	ACC Thr	CAT His	GTT Val	ATC Ile	CAC His	CAT His	CTT Leu	TTC Phe	CCC Pro	CAA Gln	ATT Ile	CCT Pro	CAT His	TAT His	CAC His	1802
305																
310																
315																

CTC GTT GAA GCG ACA CAA GCA GCA AAA CCA GTT CTT GGA GAT TAC TAC 1850  
 Leu Val Glu Ala Thr Gln Ala Ala Lys Pro Val Leu Gly Asp Tyr Tyr  
 320 325 330

CGT GAG CCA GAA AGA TCT GCG CCA TTA CCA TTT CAT CTA ATA AAG TAT 1898  
 Arg Glu Pro Glu Arg Ser Ala Pro Leu Pro Phe His Leu Ile Lys Tyr  
 335 340 345

TTA ATT CAG AGT ATG AGA CAA GAC CAC TTC GTA AGT GAC ACT GGA GAT 1946  
 Leu Ile Gln Ser Met Arg Gln Asp His Phe Val Ser Asp Thr Gly Asp  
 350 355 360

GTT GTT TAT TAT CAG ACT GAT TCT CTG CTC CTC CAC TCG CAA CGA GAC 1994  
 Val Val Tyr Tyr Gln Thr Asp Ser Leu Leu Leu His Ser Gln Arg Asp  
 365 370 375 380

TGAGTTTCAA ACTTTTGGG TTATTATTTA TTGGATTCTA GCTACTCAAA TTACTTTTTT 2054  
 TTTAATGTGA TGTTTTTGG AGTTTAACGT TTTCGAACA ACTTGCAAA TACTTGCTA 2114  
 GAGAGACATG GAATATTTAT TTGAAATTAG TAAGTAGTA ATAATAAATT TTGAATTGTC 2174  
 AGTTTCA 2181

## (2) INFORMATION FOR SEQ ID NO:11:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 380 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

Met Val Lys Asp Thr Lys Pro Leu Ala Tyr Ala Ala Asn Asn Gly Tyr  
 1 5 10 15

Gln Gln Lys Gly Ser Ser Phe Asp Phe Asp Pro Ser Ala Pro Pro Pro  
 20 25 30

Phe Lys Ile Ala Glu Ile Arg Ala Ser Ile Pro Lys His Cys Trp Val  
 35 40 45

Lys Asn Pro Trp Arg Ser Leu Ser Tyr Val Leu Arg Asp Val Leu Val  
 50 55 60

Ile Ala Ala Leu Val Ala Ala Ala Ile His Phe Asp Asn Trp Leu Leu  
 65 70 75 80

Trp Leu Ile Tyr Cys Pro Ile Gln Gly Thr Met Phe Trp Ala Leu Phe  
 85 90 95

Val Leu Gly His Asp Cys Gly His Gly Ser Phe Ser Asp Ser Pro Leu  
 100 105 110



Leu Asn Ser Leu Val Gly His Ile Leu His Ser Ser Ile Leu Val Pro  
 115 120 125  
 Tyr His Gly Trp Arg Ile Ser His Arg Thr His His Gln Asn His Gly  
 130 135 140  
 His Ile Glu Lys Asp Glu Ser Trp Val Pro Leu Thr Glu Lys Ile Tyr  
 145 150 155 160  
 Lys Asn Leu Asp Ser Met Thr Arg Leu Ile Arg Phe Thr Val Pro Phe  
 165 170 175  
 Pro Leu Phe Val Tyr Pro Ile Tyr Leu Phe Ser Arg Ser Pro Gly Lys  
 180 185 190  
 Glu Gly Ser His Phe Asn Pro Tyr Ser Asn Leu Phe Pro Pro Ser Glu  
 195 200 205  
 Arg Lys Gly Ile Ala Ile Ser Thr Leu Cys Trp Ala Thr Met Phe Ser  
 210 215 220  
 Leu Leu Ile Tyr Leu Ser Phe Ile Thr Ser Pro Leu Leu Val Leu Lys  
 225 230 235 240  
 Leu Tyr Gly Ile Pro Tyr Trp Ile Phe Val Met Trp Leu Asp Phe Val  
 245 250 255  
 Thr Tyr Leu His His His Gly His His Gln Lys Leu Pro Trp Tyr Arg  
 260 265 270  
 Gly Lys Glu Trp Ser Tyr Leu Arg Gly Gly Leu Thr Thr Val Asp Arg  
 275 280 285  
 Asp Tyr Gly Trp Ile Tyr Asn Ile His His Asp Ile Gly Thr His Val  
 290 295 300  
 Ile His His Leu Phe Pro Gln Ile Pro His Tyr His Leu Val Glu Ala  
 305 310 315 320  
 Thr Gln Ala Ala Lys Pro Val Leu Gly Asp Tyr Tyr Arg Glu Pro Glu  
 325 330 335  
 Arg Ser Ala Pro Leu Pro Phe His Leu Ile Lys Tyr Leu Ile Gln Ser  
 340 345 350  
 Met Arg Gln Asp His Phe Val Ser Asp Thr Gly Asp Val Val Tyr Tyr  
 355 360 365  
 Gln Thr Asp Ser Leu Leu Leu His Ser Gln Arg Asp  
 370 375 380

## (2) INFORMATION FOR SEQ ID NO:12:

## (1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1675 base pairs  
 (B) TYPE: nucleic acid

(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Glycine max

(vii) IMMEDIATE SOURCE:

(B) CLONE: pSFD-118bwp

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 169..1530

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

CTGTGGCAAT TTTTCTCTTC TCCTTCTGGT TCTCATCTTT GTGTTCTTCT TTGTTTCTCA	60
CCTTCTGAG GATTTTTCOA TCTTAGTTCC TGGAGGCACC AGGAACCTGA CCAAATAAAT	120
AAACCTTTTT TTTCTTCTAA TTTTCTGAA GTTTCATTTT TAGTCCA ATG GCA ACT	177
Met Ala Thr	
1	
TGG TAT CAT CAG AAA TGT GGC TTG AAG CCT CTT GCT CCA GTA ATT CCT	225
Trp Tyr His Gln Lys Cys Gly Leu Lys Pro Leu Ala Pro Val Ile Pro	
5 10 15	
AGA CCT AGA ACT GGG GCT GCT TTG TCC AGC ACC TCA AGG GTT GAA TTT	273
Arg Pro Arg Thr Gly Ala Ala Leu Ser Ser Thr Ser Arg Val Glu Phe	
20 25 30 35	
TTG GAC ACA AAC AAG GTA GTG GCA GGT CCT AAG TTT CAA CCT TTG AGG	321
Leu Asp Thr Asn Lys Val Val Ala Gly Pro Lys Phe Gln Pro Leu Arg	
40 45 50	
TGC AAC CTC AGG GAG AGG AAT TGG GGG CTG AAA GTG AGT GCC CCT TTG	369
Cys Asn Leu Arg Glu Arg Asn Trp Gly Leu Lys Val Ser Ala Pro Leu	
55 60 65	
AGG GTT GCT TCC ATT GAA GAG GAG CAA AAG AGT GTT GAT TTA ACC AAT	417
Arg Val Ala Ser Ile Glu Glu Glu Gln Lys Ser Val Asp Leu Thr Asn	
70 75 80	
GGG ACT AAT GGG GTT GAG CAT GAG AAG CTT CCA GAA TTT GAC CCT GGT	465
Gly Thr Asn Gly Val Glu His Glu Lys Leu Pro Glu Phe Asp Pro Gly	
85 90 95	
GCT CCG CCA CCA TTC AAC TTG GCT GAT ATT AGA GCA GCC ATT CCA AAG	513
Ala Pro Pro Pro Phe Asn Leu Ala Asp Ile Arg Ala Ala Ile Pro Lys	
100 105 110 115	

CAT His	TGC Cys	TGG Trp	GTG Val	AAG Lys 120	GAC Asp	CCT Pro	TGG Trp	AGG Arg	TCC Ser 125	ATG Met	AGC Ser	TAT Tyr	GTG Val	GTG Val	AGG Arg	561
GAT Asp	GTG Val	ATT Ile	GCT Ala 135	GTC Val	TTT Phe	GGT Gly	TTG Leu 140	GCT Ala 140	GCT Ala	GCT Ala	GCG Ala	TAT Tyr 145	CTC Leu	AAT Asn		609
AAT Asn	TGG Trp	TTG Leu 150	GTT Val	TGG Trp	CCT Pro	CTC Leu	TAT Tyr 155	TGG Trp	GCT Ala	GCT Ala	CAA Gln	GGC Gly 160	ACT Thr	ATG Met	TTC Phe	657
TGG Trp	GCT Ala 165	CTG Leu	TTT Phe	GTT Val	CTT Leu	GGT Gly 170	CAT His	GAT Asp	TGT Cys	GGT Gly	CAT His 175	GGA Gly	AGC Ser	TTT Phe	TCA Ser	705
AAC 180	AAC Asn	TCC Ser	AAA Lys	TTG Leu	AAC Asn 185	AGT Ser	GTT Val	GTT Val	GGA Gly	CAT His 190	CTG Leu	CTG Leu	CAT His	TCT Ser	TCA Ser 195	753
ATT Ile	CTA Leu	GTG Val	CCA Pro	TAT Tyr 200	CAT His	GGA Gly	TGG Trp	AGA Arg	ATC Ile 205	AGT Ser	CAT His	AGG Arg	ACT Thr	CAT His 210	CAC His	801
CAA Gln	CAT His	CAT His	GGT Gly 215	CAT His	GCT Ala	GAA Glu	AAT Asn	GAT Asp 220	GAA Glu	TCA Ser	TGG Trp	CAT His 225	CCG Pro	TTG Leu	CCT Pro	849
GAA Glu	AAA Lys	TTG Leu 230	TTC Phe	AGA Arg	AGC Ser	TTG Leu	GAC Asp 235	ACT Thr	GTA Val	ACT Thr	CGT Arg	ATG Met 240	TTA Leu	AGA Arg	TTC Phe	897
ACA Thr	GCA Ala	CCT Pro	TTT Phe	CCA Pro	CTT Leu 250	CTT Leu	GCA Ala	TTT Phe	CCT Pro	GTG Val	TAC Tyr 255	CTT Leu	TTT Phe	AGT Ser	AGG Arg	945
AGT Ser 260	CCT Pro	GGG Gly	ACT Lys	ACT Thr	GGT Gly 265	TCT Ser	CAC His	TTT Phe	GAC Asp	CCC Pro 270	AGC Ser	AGT Ser	GAC Asp	TTG Leu 275	TTC Phe	993
GTT Val	CCC Pro	AAT Asn	GAA Glu	AGA Arg 280	AAA Lys	GAT Asp	GTT Val	ATT Ile	ACT Thr 285	TCC Ser	ACA Thr	GCT Ala	TGT Cys	TGG Trp 290	GCT Ala	1041
GCT Ala	ATG Met	TTG Leu	GGA Gly 295	TTG Leu	CTT Leu	GTT Val	GGA Gly 300	TTG Gly	GGG Phe	TTT Phe	GTA Val	ATG Met	GGT Gly 305	CCA Pro	ATT Ile	1089
CAA Gln	CTT Leu	CTT Leu	AAG Lys 310	CTT Leu	TAT Tyr	GGT Gly	GTT Val 315	CCC Pro	TAT Tyr	GTT Val	ATA Ile	TTC Phe 320	GTT Val	ATG Met	TGG Trp	1137
TTG Leu	GAT Asp 325	TTG Leu	GTG Val	ACT Thr	TAT Tyr	TTG Leu 330	CAC His	CAT His	CAT His	GGC Gly 335	CAT His	GAA Glu	GAC Asp	AAA Lys	TTA Leu	1185

CCT TGG TAC CGT GGA AAG GAA TGG AGC TAC CTC AGG GGT GGT CTA ACT 1233  
 Pro Trp Tyr Arg Gly Lys Glu Trp Ser Tyr Leu Arg Gly Gly Leu Thr 355  
 340 345 350

ACT CTT GAT CGT GAT TAT GGA TGG ATC AAT AAC ATT CAC CAT GAC ATT 1281  
 Thr Leu Asp Arg Asp Tyr Gly Trp Ile Asn Asn Ile His His Asp Ile 370  
 360 365

GGC ACT CAT GTC ATT CAT CAC CTA TTT CCT CAA ATT CCA CAC TAT CAC 1329  
 Gly Thr His Val Ile His His Leu Phe Pro Gln Ile Pro His Tyr His 385  
 375 380

TTA GTT GAG GCT ACT GAG GCT GCT AAG CCA GTG TTT GGA AAA TAT TAT 1377  
 Leu Val Glu Ala Thr Glu Ala Lys Pro Val Phe Gly Lys Tyr Tyr 400  
 390 395

AGA GAA CCA AAG AAA TCA GCA GCA CCT CTT CCT TTT CAC CTT ATT GGG 1425  
 Arg Glu Pro Lys Lys Ser Ala Ala Pro Leu Pro Phe His Leu Ile Gly 415  
 405 410

GAA ATA ATA AGG AGC TTC AAG ACT GAC CAT TTT GTT AGT GAC ACG GGG 1473  
 Glu Ile Ile Arg Ser Phe Lys Thr Asp His Phe Val Ser Asp Thr Gly 435  
 420 425 430

GAT GTT GTG TAC TAT CAA ACC GAC TCT AAG ATT AAT GGC TCT TCC AAA 1521  
 Asp Val Val Tyr Tyr Gln Thr Asp Ser Lys Ile Asn Gly Ser Ser Lys 450  
 440 445

TTA GAG TGAATATTAA AATTCTTTTC TATATAGACA AGAGAGGCTT ATACACAATT 1577  
 Leu Glu

CTTATTGCTT TAAAGATTGT CTGAGTTTC TCCGAAAGTT ACTGCACCTA CTGGAGTTG 1637

AATCCTTCAT TAATAAGGG ATGGATGGAT CATATAAA 1675

## (2) INFORMATION FOR SEQ ID NO:13:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 453 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

Met Ala Thr Trp Tyr His Gln Lys Cys Gly Leu Lys Pro Leu Ala Pro  
 1 5 10 15

Val Ile Pro Arg Pro Arg Thr Gly Ala Ala Leu Ser Ser Thr Ser Arg  
 20 25 30

Val Glu Phe Leu Asp Thr Asn Lys Val Val Ala Gly Pro Lys Phe Gln  
 35 40 45

Pro Leu Arg Cys Asn Leu Arg Glu Arg Asn Trp Gly Leu Lys Val Ser  
 50 55 60  
 Ala Pro Leu Arg Val Ala Ser Ile Glu Glu Glu Gln Lys Ser Val Asp  
 65 70 75 80  
 Leu Thr Asn Gly Thr Asn Gly Val Glu His Glu Lys Leu Pro Glu Phe  
 85 90 95  
 Asp Pro Gly Ala Pro Pro Pro Phe Asn Leu Ala Asp Ile Arg Ala Ala  
 100 105 110  
 Ile Pro Lys His Cys Trp Val Lys Asp Pro Trp Arg Ser Met Ser Tyr  
 115 120 125  
 Val Val Arg Asp Val Ile Ala Val Phe Gly Leu Ala Ala Ala Ala Ala  
 130 135 140  
 Tyr Leu Asn Asn Trp Leu Val Trp Pro Leu Tyr Trp Ala Ala Gln Gly  
 145 150 155 160  
 Thr Met Phe Trp Ala Leu Phe Val Leu Gly His Asp Cys Gly His Gly  
 165 170 175  
 Ser Phe Ser Asn Asn Ser Lys Leu Asn Ser Val Val Gly His Leu Leu  
 180 185 190  
 His Ser Ser Ile Leu Val Pro Tyr His Gly Trp Arg Ile Ser His Arg  
 195 200 205  
 Thr His His Gln His His Gly His Ala Glu Asn Asp Glu Ser Trp His  
 210 215 220  
 Pro Leu Pro Glu Lys Leu Phe Arg Ser Leu Asp Thr Val Thr Arg Met  
 225 230 235 240  
 Leu Arg Phe Thr Ala Pro Phe Pro Leu Leu Ala Phe Pro Val Tyr Leu  
 245 250 255  
 Phe Ser Arg Ser Pro Gly Lys Thr Gly Ser His Phe Asp Pro Ser Ser  
 260 265 270  
 Asp Leu Phe Val Pro Asn Glu Arg Lys Asp Val Ile Thr Ser Thr Ala  
 275 280 285  
 Cys Trp Ala Ala Met Leu Gly Leu Leu Val Gly Leu Gly Phe Val Met  
 290 295 300  
 Gly Pro Ile Gln Leu Leu Lys Leu Tyr Gly Val Pro Tyr Val Ile Phe  
 305 310 315 320  
 Val Met Trp Leu Asp Leu Val Thr Tyr Leu His His His Gly His Glu  
 325 330 335  
 Asp Lys Leu Pro Trp Tyr Arg Gly Lys Glu Trp Ser Tyr Leu Arg Gly  
 340 345 350

1  
 2  
 3  
 4  
 5  
 6  
 7  
 8  
 9  
 10  
 11  
 12  
 13  
 14  
 15  
 16  
 17  
 18  
 19  
 20  
 21  
 22  
 23  
 24  
 25  
 26  
 27  
 28  
 29  
 30  
 31  
 32  
 33  
 34  
 35  
 36  
 37  
 38  
 39  
 40  
 41  
 42  
 43  
 44  
 45  
 46  
 47  
 48  
 49  
 50  
 51  
 52  
 53  
 54  
 55  
 56  
 57  
 58  
 59  
 60  
 61  
 62  
 63  
 64  
 65  
 66  
 67  
 68  
 69  
 70  
 71  
 72  
 73  
 74  
 75  
 76  
 77  
 78  
 79  
 80  
 81  
 82  
 83  
 84  
 85  
 86  
 87  
 88  
 89  
 90  
 91  
 92  
 93  
 94  
 95  
 96  
 97  
 98  
 99  
 100

Gly Leu Thr Thr Leu Asp Arg Asp Tyr Gly Trp Ile Asn Asn Ile His  
 355 360 365  
 His Asp Ile Gly Thr His Val Ile His His Leu Phe Pro Gln Ile Pro  
 370 375 380  
 His Tyr His Leu Val Glu Ala Thr Glu Ala Ala Lys Pro Val Phe Gly  
 385 390 395 400  
 Lys Tyr Tyr Arg Glu Pro Lys Lys Ser Ala Ala Pro Leu Pro Phe His  
 405 410 415  
 Leu Ile Gly Glu Ile Ile Arg Ser Phe Lys Thr Asp His Phe Val Ser  
 420 425 430  
 Asp Thr Gly Asp Val Val Tyr Tyr Gln Thr Asp Ser Lys Ile Asn Gly  
 435 440 445  
 Ser Ser Lys Leu Glu  
 450

## (2) INFORMATION FOR SEQ ID NO:14:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 396 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: cDNA

## (iii) HYPOTHETICAL: NO

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Zea mays

## (vii) IMMEDIATE SOURCE:

- (B) CLONE: pPCR20

## (ix) FEATURE:

- (A) NAME/KEY: exon  
 (B) LOCATION: 31..363

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

GGATCCACGC ATCATCAGAA TCACGGTCAC ATCCACAGGG ACGAGTCATG GCACCCGATC	60
ACGGAGAAGC TGTACCGGCA ACTAGAGCCA CGCACCAAGA AGCTGAGATT CACGGTGCCC	120
TTCCCCCTGC TCGCATTCCC CGTCTACCTC TTGTACAGGA GCCCCGGCAA GCTCGGCTCC	180
CACTTCCTTC CCAGCAGCGA CCTGTTCAGC CCCAAGGAGA AGAGCGACGT CATGGTGTCA	240

001005-1200

ACCACCTGCT GGTGCATCAT GCTCGCCTCC CTCCTCGCCA TGGCGTGCGC GTTCGGCCCA	300
CTCCAGGTGC TCAAGATGTA CGGCATCCCA TACCTGGTGT TCGTGATGTG GCTTGACCTG	360
GTGACGTACT TACATCACCA CGGCCACGAT GGATCC	396

## (2) INFORMATION FOR SEQ ID NO:15:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 126 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: unknown
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: protein

(iii) HYPOTHETICAL: YES

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Zea mays

(vii) IMMEDIATE SOURCE:

(B) CLONE: pPCR20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

His	His	Gln	Asn	His	Gly	His	Ile	His	Arg	Asp	Glu	Ser	Trp	His	Pro
1					5				10					15	
Ile	Thr	Glu	Lys	Leu	Tyr	Arg	Gln	Leu	Glu	Pro	Arg	Thr	Lys	Lys	Leu
			20				25						30		
Arg	Phe	Thr	Val	Pro	Phe	Pro	Leu	Leu	Ala	Phe	Pro	Val	Tyr	Leu	Leu
			35				40					45			
Tyr	Arg	Ser	Pro	Gly	Lys	Leu	Gly	Ser	His	Phe	Leu	Pro	Ser	Ser	Asp
			50			55					60				
Leu	Phe	Ser	Pro	Lys	Glu	Lys	Ser	Asp	Val	Met	Val	Ser	Thr	Thr	Cys
			65			70			75					80	
Trp	Cys	Ile	Met	Leu	Ala	Ser	Leu	Leu	Ala	Met	Ala	Cys	Ala	Phe	Gly
			85					90						95	
Pro	Leu	Gln	Val	Leu	Lys	Met	Tyr	Gly	Ile	Pro	Tyr	Leu	Val	Phe	Val
			100					105					110		
Met	Trp	Leu	Asp	Leu	Val	Thr	Tyr	Leu	His	His	His	Gly	His		
			115				120					125			

20250315.122300





(vii) IMMEDIATE SOURCE:

(B) CLONE: pFadx-2 and pYacp7

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

Ser	Ser	Tyr	Val	Arg	Ala	Lys	Thr	Arg	Asn	Trp	Ala	Leu	Asn	Val	Ala
1				5					10					15	
Thr	Pro	Leu	Thr	Thr	Leu	Gln	Ser	Pro	Ser	Glu	Glu	Asp	Arg	Glu	Arg
			20					25					30		
Phe	Asp	Pro	Gly	Ala	Pro	Pro	Pro	Phe	Asn	Leu	Ala	Asp	Ile	Arg	Ala
		35					40					45			
Ala	Ile	Pro	Lys	His	Cys	Trp	Val	Lys	Asn	Pro	Trp	Met	Ser	Met	Ser
		50				55					60				
Tyr	Val	Val	Arg	Asp	Val	Ala	Ile	Val	Phe	Gly	Leu	Ala	Ala	Val	Ala
					70					75					80
Ala	Tyr	Phe	Asn	Asn	Trp	Leu	Leu	Trp	Pro	Leu	Tyr	Trp	Phe	Ala	Gln
				85					90					95	
Gly	Thr	Met	Phe	Trp	Ala	Leu	Phe	Val	Leu	Gly	His	Asp	Cys	Gly	His
			100					105					110		
Gly	Ser	Phe	Ser	Asn	Asp	Pro	Arg	Leu	Asn	Ser	Val	Ala	Gly	His	Leu
		115				120						125			
Leu	His	Ser	Ser	Ile	Leu	Val	Pro	Tyr	His	Gly	Trp	Arg	Ile	Ser	His
						135						140			
Arg	Thr	His	His	Gln	Asn	His	Gly	His	Val	Glu	Asn				
145					150					155					

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 31 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ix) FEATURE:

(A) NAME/KEY: misc feature  
(B) LOCATION: 1..11  
(D) OTHER INFORMATION: /note= "N= INOSINE"

(ix) FEATURE:

(A) NAME/KEY: misc feature  
(B) LOCATION: 12..31  
(D) OTHER INFORMATION: /note= "N= A OR T OR G OR C"

CGGGATCCAC NCAYCAYCAR AAYCAYGGNC A

31

(2) INFORMATION FOR SEQ ID NO:19:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 35 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ix) FEATURE:

(A) NAME/KEY: misc feature  
(B) LOCATION: 1..15  
(D) OTHER INFORMATION: /note= "N= INOSINE"

(ix) **FEATURE:**

```
(A) NAME/KEY:  misc feature
(B) LOCATION:  16.35
(D) OTHER INFORMATION:  /note= "N= A OR T OR G OR C"
```

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

CGGGATCCRT CRTGNCCRTG RTGRTGNARR TANGT

35

(2) INFORMATION FOR SEQ ID NO:20:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 42 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ix) FEATURE:

(A) NAME/KEY: misc feature  
(B) LOCATION: 1..36  
(D) OTHER INFORMATION: /note= "N= INOSINE"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

TTCGTNNTNG GNCAYGAYTG YGGNCAYGGN CAYGGNAGNT TC

42

(2) INFORMATION FOR SEQ ID NO:21:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 36 base pairs  
(B) TYPE: nucleic acid

- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ix) FEATURE:

- (A) NAME/KEY: misc feature
- (B) LOCATION: 1..36
- (D) OTHER INFORMATION: /note= "N= INOSINE"

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

TTCGTNNTNG GNCAYGAYTG YGGNCAYGGN TCNTTC

36

## (2) INFORMATION FOR SEQ ID NO:22:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 18 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

GGHCAYGAYT GYGHCAC

18

## (2) INFORMATION FOR SEQ ID NO:23:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 18 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

GGHCAYGAYT GYGHCAT

18

## (2) INFORMATION FOR SEQ ID NO:24:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 18 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

GTACTRTARC CDTGDGTR

18

## (2) INFORMATION FOR SEQ ID NO:25:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 18 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

GTGCTRTARC CDTGDGTR

18

## (2) INFORMATION FOR SEQ ID NO:26:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 23 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

GTRCANTARG TRGTRAAAYAA YGG

23

## (2) INFORMATION FOR SEQ ID NO:27:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 23 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

GTRCANTADG TRGTRGADAA YGG

23

## (2) INFORMATION FOR SEQ ID NO:28:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 36 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ix) FEATURE:

- (A) NAME/KEY: misc feature
- (B) LOCATION: 1..36
- (D) OTHER INFORMATION: /note= "N= INOSINE"

00210035.102300

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

TTCGTNNTNG GNCAYGAYTG YGNCAYGGN AGNTTT

36

(2) INFORMATION FOR SEQ ID NO:29:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 36 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ix) FEATURE:

- (A) NAME/KEY: misc feature
- (B) LOCATION: 1..36
- (D) OTHER INFORMATION: /note= "N= INOSINE"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

TTCGTNNTNG GNCAYGAYTG YGNCAYGGN TCNTTT

36

(2) INFORMATION FOR SEQ ID NO:30:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 38 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ix) FEATURE:

- (A) NAME/KEY: misc feature
- (B) LOCATION: 1..38
- (D) OTHER INFORMATION: /note= "N= INOSINE"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

GTRCTRTANC CNTGNGTNC NTANGTAGT RANAAGGG

38

(2) INFORMATION FOR SEQ ID NO:31:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 38 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ix) FEATURE:

- (A) NAME/KEY: misc feature
- (B) LOCATION: 1..38

(D) OTHER INFORMATION: /note= "N= INOSINE"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

GTRCTRTRTANC CNTGNGTNCA NTANGTGGTG RANAAGGG

38

(2) INFORMATION FOR SEQ ID NO:32:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 138 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ix) FEATURE:

(A) NAME/KEY: misc feature  
(B) LOCATION: 1..135  
(D) OTHER INFORMATION: /note= "N= INOSINE"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

GTGGTGNGTN CNGTGNGANA NNCKCCANCC GTGGTANGGN ACNANNANGA ANGANGAGTG

60

NANNANGTGN CCNACNANNG AGTTNANNAN NGGNATNTCN GAGAANGANC CGTGNC CGCA

120

NTCGTGNC CN ANNACGAA

138

09210035.122308

CLAIMS

1. An isolated nucleic acid fragment comprising a nucleic acid sequence encoding a fatty acid desaturase or a fatty acid desaturase-related enzyme with an amino acid identity of 50% or greater to the polypeptide encoded by SEQ ID NOS:1, 4, 6, 8, 10, 12, 14 or 16.
2. The isolated nucleic acid fragment of Claim 1 wherein the amino acid identity is 65% or greater to the polypeptide encoded by SEQ ID NOS:1, 4, 6, 8, 10, 12, 14 or 16.
3. The isolated nucleic acid fragment of Claim 1 wherein the nucleic acid identity is 90% or greater to SEQ ID NOS:1, 4, 6, 8, 10, 12, 14 or 16.
4. An isolated nucleic acid fragment of Claim 1 wherein said fragment is isolated from a plant selected from the group consisting of soybean, oilseed Brassica species, Arabidopsis thaliana and corn.
5. A chimeric gene capable of causing altered levels of linolenic acid in a transformed plant cell, the gene comprising a nucleic acid fragment of any of Claims 1, 2, or 3, the fragment operably linked to suitable regulatory sequences.
6. Plants containing the chimeric genes of Claim 5.
7. Oil obtained from seeds of the plants containing the chimeric genes of Claim 5.
8. A method of producing seed oil containing altered levels of linolenic (18:3) acid comprising:
- (a) transforming a plant cell of an oil-producing species with a chimeric gene of Claim 5;
  - (b) growing fertile plants from the transformed plant cells of step (a);
  - (c) screening progeny seeds from the fertile plants of step (b) for the desired levels of linolenic (18:3) acid; and

09210935.1.22308

(d) processing the progeny seed of step (c) to obtain seed oil containing altered levels of linolenic (18:3) acid.

9 The product of the method of Claim 8.

5 10. A method of Claim 8 wherein said plant cell of an oil-producing species is selected from the group consisting of Arabidopsis thaliana, soybean, oilseed Brassica species, sunflower, cotton, cocoa, peanut, safflower, and corn.

10 11. A method of breeding plant species producing altered levels of linolenic acid in the seed oil of oil-producing plant species comprising:

15 (a) making a cross between two varieties of oil-producing species differing in the linolenic acid trait;

(b) making a Southern blot of restriction enzyme digested genomic DNA isolated from several progeny plants resulting from the cross of step (a); and

20 (c) hybridizing the Southern blot with a radiolabelled nucleic acid fragment of Claim 1.

12. The product of the method of Claim 11.

13. A method of RFLP mapping in a genomic RFLP marker comprising:

25 (a) making a cross between two varieties of plants;

(b) making a Southern blot of restriction enzyme digested genomic DNA isolated from several progeny plants resulting from the cross of step (a); and

30 (c) hybridizing the Southern blot with a radiolabelled nucleic acid fragments of Claim 1.

14. A method to isolate nucleic acid fragments encoding fatty acid desaturases and fatty acid desaturase-related enzymes, comprising:



(a) comparing SEQ ID NOS:2, 5, 7, 9, 11, 13, 15 and 17 with other fatty acid desaturase polypeptide sequences;

(b) identifying the conserved sequence(s) of  
5 4 or more amino acids obtained in step (a);

(c) making region-specific nucleotide probe(s) or oligomer(s) based on the conserved sequences identified in step b; and

(d) using the nucleotide probe(s) or  
10 oligomers(s) of step c to isolate sequences encoding fatty acid desaturases and fatty-acid desaturase-related enzymes by sequence-dependent protocols.

15. The product of the method of Claim 14.

16. The isolated genomic DNA of Arabidopsis  
15 thaliana identified by accession number ATCC 75167.

17. An isolated cDNA clone which encodes for soybean delta-15 desaturase, the clone designated pXF1 comprising the DNA sequence of SEQ ID NO 10 and identified by accession number ATCC 68874.

20 18. An isolated cDNA clone which encodes for oilseed Brassica species delta-15 desaturase, the clone designated pBNSF3 comprising the DNA sequence of SEQ ID NO:6 and identified by accession number ATCC 68854.

25 19. An isolated Polymerase Chain Reaction Product for Zea mays delta-15 desaturase, the clone designated pcr20 comprising the DNA sequence of SEQ ID NO:14.

TITLE

FATTY ACID DESATURASE GENES FROM PLANTS

ABSTRACT OF THE DISCLOSURE

5 The preparation and use of nucleic acid fragments  
encoding fatty acid desaturase enzymes are described.  
The invention permits alteration of plant lipid  
composition. Chimeric genes incorporating such nucleic  
acid fragments with suitable regulatory sequences may be  
used to create transgenic plants with altered levels of  
10 unsaturated fatty acids.

15

20

25

30

35 LAF/bjm

I hereby claim the benefit under Title 35, United States Code, §120 of any United States application(s) or PCT international application(s) designating the United States of America that is/are listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in that/those prior application(s) in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, §1.56(a) which occurred between the filing date of the prior application(s) and the national or PCT international filing date of this application:

**PRIOR U.S. APPLICATIONS OR PCT INTERNATIONAL APPLICATIONS DESIGNATING THE U.S. FOR BENEFIT UNDER 35 U.S.C. 120:**

U.S. APPLICATIONS		STATUS (Check one)		
U.S. APPLICATION NUMBER	U.S. FILING DATE	PATENTED	PENDING	ABANDONED
07/804,259	04 December 1991			X
PCT APPLICATIONS DESIGNATING THE U.S.				
PCT APPLICATION NO.	PCT FILING DATE	U.S. SERIAL NUMBERS ASSIGNED (if any)		
PCT/US92/10284	03 December 1992		X	

**POWER OF ATTORNEY:** As a named inventor, I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and transact all business in the Patent and Trademark Office connected therewith. (List name and registration number)

Barbara C. Siegell, Reg. No. 30,684

Hoge T. Sutherland, Reg. No. 20,010

Send Correspondence to: Barbara C. Siegell

Legal/Patent Records Center  
 E. I. du Pont de Nemours and Company  
 1007 Market Street  
 Wilmington, Delaware 19898

Direct Telephone Calls to:  
 (name and telephone number)

Barbara C. Siegell  
 (302) 992-4931

201	FULL NAME OF INVENTOR	FAMILY NAME	FIRST GIVEN NAME	SECOND GIVEN NAME
	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP
	POST OFFICE ADDRESS	CITY	CITY	STATE & ZIP CODE COUNTRY
202	FULL NAME OF INVENTOR	FAMILY NAME	FIRST GIVEN NAME	SECOND GIVEN NAME
	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP
	POST OFFICE ADDRESS	CITY	CITY	STATE & ZIP CODE COUNTRY
203	FULL NAME OF INVENTOR	FAMILY NAME	FIRST GIVEN NAME	SECOND GIVEN NAME
	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP
	POST OFFICE ADDRESS	CITY	CITY	STATE & ZIP CODE COUNTRY

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

SIGNATURE OF INVENTOR 201	SIGNATURE OF INVENTOR 202	SIGNATURE OF INVENTOR 203
John Browne	Luis Perez-Grau	A Skinner
DATE	DATE	DATE
August 17, 1994	June 10, 94	July 11, 94

204	FULL NAME OF INVENTOR	FAMILY NAME	FIRST GIVEN NAME	SECOND GIVEN NAME
	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP
	POST OFFICE ADDRESS	CITY	STATE & ZIP CODE COUNTRY	
205	FULL NAME OF INVENTOR	FAMILY NAME	FIRST GIVEN NAME	SECOND GIVEN NAME
	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP
	POST OFFICE ADDRESS	CITY	STATE & ZIP CODE COUNTRY	
206	FULL NAME OF INVENTOR	FAMILY NAME	FIRST GIVEN NAME	SECOND GIVEN NAME
	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP
	POST OFFICE ADDRESS	CITY	STATE & ZIP CODE COUNTRY	

207	FULL NAME OF INVENTOR	FAMILY NAME	FIRST GIVEN NAME	SECOND GIVEN NAME
	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP
	POST OFFICE ADDRESS	CITY	STATE & ZIP CODE COUNTRY	
208	FULL NAME OF INVENTOR	FAMILY NAME	FIRST GIVEN NAME	SECOND GIVEN NAME
	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP
	POST OFFICE ADDRESS	CITY	STATE & ZIP CODE COUNTRY	
209	FULL NAME OF INVENTOR	FAMILY NAME	FIRST GIVEN NAME	SECOND GIVEN NAME
	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP
	POST OFFICE ADDRESS	CITY	STATE & ZIP CODE COUNTRY	

SIGNATURE OF INVENTOR 204	SIGNATURE OF INVENTOR 205	SIGNATURE OF INVENTOR 206
DATE	DATE	DATE
SIGNATURE OF INVENTOR 207	SIGNATURE OF INVENTOR 208	SIGNATURE OF INVENTOR 209
DATE	DATE	DATE